

Stefano Frixione

QCD at NLO: Monte Carlo developments

DIS04, Strbske Pleso, 14/4/2004

Higher orders versus MC's

	<i>Good</i>	<i>Bad</i>	<i>Users</i>
HO	Hard emissions Total rates	Soft&coll emissions Hadronization No events	Theorists
MC	Soft&coll emissions Hadronization Outputs events	Hard emissions Total rates	Experimentalists

In other words: $HO \cap MC = \emptyset$

A formalism incorporating HO *and* MC should combine their *Good* features, avoiding the *Bad* ones. However, the radical differences between the two approaches made QCDists wonder whether such a combination was possible

Motivations for matching HO and MC

A formalism with all the **Good** features is certainly desirable, and its definition is a challenging theoretical problem. But, are there compelling physical motivations?

- It is not unlikely that new physics signals will emerge from counting experiments, which require firm control on SM signal and background simulations
- The high-energy regime of the Tevatron and the LHC implies the relevance of **multi-jet, multi-scale processes, with large K -factors**
- Standard MC's don't perform well in predicting multi-jet observables, and the practice of multiplying the results by inclusive K -factors is just wrong. This may lead to **major errors in the strategies for searches** (kind of new in HEP!)
- Multi-scale processes are badly predicted by fixed-order computations. Results matching these computations with resummed ones are mandatory (**a procedure largely successful at LEP**)
- The hadronization procedure in HO computations is extremely naive, and strictly speaking can be applied only at very large p_T 's

A less ambitious goal

The problem is much simplified if one selects only those Feynman diagrams contributing to HO, which correspond to the emissions of real particles

The improved MC is capable of simulating the emission of n_E extra hard partons

Implicit is the notion of Born \equiv LO level, the process(es) with the smallest number n_B of final-state partons which contributes to a given reaction (usually $n_B = 2$)

This procedure is called **Matrix Element Correction**


The strategy: generate the hard subprocesses with a standalone package (AcerMC, ALPGEN, AMEGIC++, CompHEP, Grace, MadEvent), and use the resulting kinematics as initial conditions for your favourite parton shower MC (HERWIG, PYTHIA)

- *Practical problems*: efficient ME generation for $(n_E + n_B)$ -parton final states, efficient phase-space sampling
- *Principle problem*: real ME's diverge in the soft/collinear regions, and a cut δ_{sep} must be introduced at the parton level to avoid divergences \implies physical observables will depend on the unphysical δ_{sep} cutoff (a 20–30% effect at best)

Getting rid of δ_{sep} dependence

In the context of e^+e^- physics, [Catani, Krauss, Kuhn & Webber](#) show that the problem **cannot be solved at fixed n_B** . Extended to colour dipoles by [Lönnblad](#); extended to hadronic collisions by [Krauss](#)

- **The problem:** δ_{sep} dependence \Leftrightarrow double counting




- **The solution:** separate the PS- and ME-dominated regions in an **arbitrary** manner; to compensate for the arbitrariness, the **shower must be modified** accordingly

- **The aim:** compute the observable at $\mathcal{O}(\alpha_S^{n-2})$, for any n , and resum to NLL accuracy (downstairs) where needed. By-product: the δ_{sep} dependence is reduced

$$\sigma_n \sim \alpha_S^{n-2} \sum_k a_k \alpha_S^k \log^{2k} \delta_{sep} \longrightarrow \alpha_S^{n-2} \left(\delta_{sep}^a + \sum_k b_k \alpha_S^k \log^{2k-2} \delta_{sep} \right)$$

Adding virtual corrections: NLOwPS

The problem: virtual corrections pose additional difficulties



The solution is unlikely to be unique, so start with some *definitions*

- ◆ Total rates are accurate to NLO
- ◆ Hard emissions are treated as in NLO computations
- ◆ Soft/collinear emissions are treated as in MC
- ◆ NLO results are recovered upon expansion of NLOwPS results in α_S .
In other words: there is no **double counting** in NLOwPS
- ◆ The matching between hard- and soft/collinear-emission regions is smooth
- ◆ The output is a set of events, which are fully exclusive
- ◆ MC hadronization models are adopted

NLO ^{n_E} wPS with $n_E \neq 1$ is unfeasible with our present understanding

NLOwPS versus MEC

■ Why is the definition of NLOwPS's much more difficult than MEC?

The problem is a fundamental one: **KLN cancellation** is achieved in standard MC's through **unitarity**, and embedded in Sudakovs. This is no longer possible: IR singularities **do appear in hard ME's**

IR singularities are avoided in MEC by cutting them off with δ_{sep} . This must be so, since only loop diagrams can cut off the divergences of real matrix elements

NLOwPS's are better than MEC since:

- + There is no δ_{sep} dependence
- + The computation of total rates is meaningful and reliable


NLOwPS's are worse than MEC since:

- The number of hard legs is smaller
- There are negative weights

A realistic goal: CKKW in NLOwPS's (i.e. multi-leg, NLO generators)

What does NLO mean?

Consider Higgs production:



$$\frac{d\sigma}{dp_T} = (A\alpha_S^2 + B\alpha_S^3) \delta(p_T) + C(p_T)\alpha_S^3$$

$$\int_{p_T^{min}}^{\infty} dp_T \frac{d\sigma}{dp_T} = \mathcal{C}_3 \alpha_S^3, \quad p_T^{min} > 0$$

$$= \mathcal{D}_2 \alpha_S^2 + \mathcal{D}_3 \alpha_S^3, \quad p_T^{min} = 0$$

$$p_T^{min} > 0 \Rightarrow \text{LO}, \quad p_T^{min} = 0 \Rightarrow \text{NLO}$$

The answer depends on the observable, and even on the kinematic range considered.
So this definition cannot be adopted in the context of event generators

■ N^k LO accuracy in event generators is defined by the number k of extra gluons (either virtual or real) wrt the LO contribution (hopefully we all agree on LO definition)

The actual NLOwPS's

- MC@NLO (Webber & SF; Nason, Webber & SF)
Based on NLO subtraction method
Formulated in general, interfaced to Herwig
Processes implemented: $H_1 H_2 \longrightarrow W^+ W^-, W^\pm Z, ZZ, b\bar{b}, t\bar{t}, H^0, W^\pm, Z/\gamma$
- Φ -veto (Dobbs & Lefebvre)
Based on NLO slicing method
Avoids negative weights, at the price of double counting
Processes implemented: $H_1 H_2 \longrightarrow Z$
- grcNLO (Kurihara *et al* – GRACE)
Based on NLO hybrid slicing method, computes ME's numerically
Double counts, if the parton shower is not built *ad hoc*
Process implemented: $H_1 H_2 \longrightarrow Z$

A proposal by Collins aims at including NLL effects in showers, but lacks gluon emission so far. Φ -veto is based on an old proposal by Baer&Reno; jets in DIS have been considered by Pötter&Schörner using a similar method. Soper&Krämer implemented $e^+ e^- \rightarrow 3$ jets (but without a realistic MC)

Fixed-order versus MC

■ NLO cross section (based on subtraction)

$$\left(\frac{d\sigma}{dO}\right)_{subt} = \sum_{ab} \int dx_1 dx_2 d\phi_3 f_a(x_1) f_b(x_2) \left[\delta(O - O(2 \rightarrow 3)) \mathcal{M}_{ab}^{(r)}(x_1, x_2, \phi_3) + \delta(O - O(2 \rightarrow 2)) \left(\mathcal{M}_{ab}^{(b,v,c)}(x_1, x_2, \phi_2) - \mathcal{M}_{ab}^{(c.t.)}(x_1, x_2, \phi_3) \right) \right]$$

■ MC

$$\mathcal{F}_{MC} = \sum_{ab} \int dx_1 dx_2 d\phi_2 f_a(x_1) f_b(x_2) \mathcal{F}_{MC}^{(2 \rightarrow 2)} \mathcal{M}_{ab}^{(b)}(x_1, x_2, \phi_2)$$

◆ Matrix elements \longrightarrow normalization, hard kinematic configurations

◆ δ -functions, $\mathcal{F}_{MC}^{(2 \rightarrow 2)} \equiv$ showers \longrightarrow kinematic “evolution”

$\Rightarrow \left(\delta(O - O(2 \rightarrow 2)), \delta(O - O(2 \rightarrow 3)) \right) \longrightarrow \left(\mathcal{F}_{MC}^{(2 \rightarrow 2)}, \mathcal{F}_{MC}^{(2 \rightarrow 3)} \right) ?$

MC@NLO is based on a modified subtraction

The naive prescription doesn't work: MC evolution results in spurious NLO terms
We eliminate them by hand

■ MC@NLO

$$\mathcal{F}_{\text{MC@NLO}} = \sum_{ab} \int dx_1 dx_2 d\phi_3 f_a(x_1) f_b(x_2) \left[\mathcal{F}_{\text{MC}}^{(2 \rightarrow 3)} \left(\mathcal{M}_{ab}^{(r)}(x_1, x_2, \phi_3) - \mathcal{M}_{ab}^{(\text{MC})}(x_1, x_2, \phi_3) \right) + \mathcal{F}_{\text{MC}}^{(2 \rightarrow 2)} \left(\mathcal{M}_{ab}^{(b,v,c)}(x_1, x_2, \phi_2) - \mathcal{M}_{ab}^{(c.t.)}(x_1, x_2, \phi_3) + \mathcal{M}_{ab}^{(\text{MC})}(x_1, x_2, \phi_3) \right) \right]$$

$$\mathcal{M}_{\mathcal{F}(ab)}^{(\text{MC})} = \mathcal{F}_{\text{MC}}^{(2 \rightarrow 2)} \mathcal{M}_{ab}^{(b)} + \mathcal{O}(\alpha_S^2 \alpha_S^b)$$

There are *two* MC-induced contributions: they eliminate the spurious NLO terms due to the **branching** of a final-state parton, and to the **non-branching** probability

Peculiarities of MC@NLO

Let's look at the weight functions

$$w_{\text{H}}(\phi_3) = \mathcal{M}_{ab}^{(r)}(x_1, x_2, \phi_3) - \mathcal{M}_{ab}^{(\text{MC})}(x_1, x_2, \phi_3)$$

$$w_{\text{S}}(\phi_3) = \mathcal{M}_{ab}^{(b,v,c)}(x_1, x_2, \phi_2) - \mathcal{M}_{ab}^{(c.t.)}(x_1, x_2, \phi_3) + \mathcal{M}_{ab}^{(\text{MC})}(x_1, x_2, \phi_3)$$

which are finite (i.e. don't diverge) for any ϕ_3

The MC provides local, observable-independent, counterterms \implies greater numerical stability, unweighting possible

MC@NLO can thus be minimally seen as a way to stabilize NLO computations, through the construction of a simplified MC whose only aim is to furnish the local counterterms. In this sense, the generalization to NNLO should not be too difficult

MC@NLO: summary

1. Choose your favourite MC (Herwig, Pythia), and compute analytically the “NLO cross section”, i.e., the first emission. This is an **observable-independent, process-independent** procedure, which is done once and for all
2. Combine the LO+NLO matrix elements of the process to be implemented according to the universal, **observable-independent, subtraction-based** formalism of SF, Kunszt, Signer for cancelling IR divergences. All counterterm, virtual, and LO contributions must have an unique kinematics (**achieved through a projection**)
3. Add and subtract the MC counterterms, computed in step 1, to the quantity computed in step 2. The resulting expression allows to generate the hard kinematic configurations, which are eventually fed into the MC showers as **initial conditions**

Negative weights don't mean negative cross sections. They arise from a different mechanism wrt those at the NLO, and their number is fairly limited

MC@NLO 2.31 [hep-ph/0402116]

IPROC	Process
-1350-IL	$H_1 H_2 \rightarrow (Z/\gamma^* \rightarrow) l_{\text{IL}} \bar{l}_{\text{IL}} + X$
-1360-IL	$H_1 H_2 \rightarrow (Z \rightarrow) l_{\text{IL}} \bar{l}_{\text{IL}} + X$
-1370-IL	$H_1 H_2 \rightarrow (\gamma^* \rightarrow) l_{\text{IL}} \bar{l}_{\text{IL}} + X$
-1460-IL	$H_1 H_2 \rightarrow (W^+ \rightarrow) l_{\text{IL}}^+ \nu_{\text{IL}} + X$
-1470-IL	$H_1 H_2 \rightarrow (W^- \rightarrow) l_{\text{IL}}^- \bar{\nu}_{\text{IL}} + X$
-1396	$H_1 H_2 \rightarrow \gamma^* (\rightarrow \sum_i f_i \bar{f}_i) + X$
-1397	$H_1 H_2 \rightarrow Z^0 + X$
-1497	$H_1 H_2 \rightarrow W^+ + X$
-1498	$H_1 H_2 \rightarrow W^- + X$
-1600-ID	$H_1 H_2 \rightarrow H^0 + X$
-1705	$H_1 H_2 \rightarrow b\bar{b} + X$
-1706	$H_1 H_2 \rightarrow t\bar{t} + X$
-2850	$H_1 H_2 \rightarrow W^+ W^- + X$
-2860	$H_1 H_2 \rightarrow Z^0 Z^0 + X$
-2870	$H_1 H_2 \rightarrow W^+ Z^0 + X$
-2880	$H_1 H_2 \rightarrow W^- Z^0 + X$

- Works identically to HERWIG: the very same analysis routines can be used
- Reads shower initial conditions from an event file (as in ME corrections)
- Exploits Les Houches accord for process information and common blocks
- Features a self contained library of PDFs with old and new sets alike
- LHAPDF will also be implemented

NLOwPS: Φ -veto

Exploit a proposal by Baer&Reno to get rid of the soft/collinear configurations:

$$\int_{\phi_0} d\phi_3 \left(\mathcal{M}_{ab}^{(b,v,c)} + \mathcal{M}_{ab}^{(r)} \right) = 0$$

Another (freely defined) phase-space region $\phi_H \subset \phi_0$ is populated by hard-emission events (Pötter, Schörner, Dobbs)

$$\begin{aligned} \mathcal{F}_{\Phi_{\text{veto}}} = & \sum_{ab} \int dx_1 dx_2 d\phi_3 f_a(x_1) f_b(x_2) \\ & \left[\mathcal{F}_{\text{MC}}^{(2 \rightarrow 3)} \mathcal{M}_{ab}^{(r)}(x_1, x_2, \phi_3) \Theta(\phi_3 \in \phi_H) + \right. \\ & \left. \mathcal{F}_{\text{MC}}^{(2 \rightarrow 2)} \mathcal{M}_{ab}^{(b,v,c)}(x_1, x_2, \phi_2) \Theta(\phi_3 \in \overline{\phi_0} \cap \overline{\phi_H}) + \right] \end{aligned}$$

- + Only positive weights
- + Doesn't need to know details of MC implementation
- **Double counting** for $\phi_3 \in \overline{\phi_H}$, and **discontinuity** at $\partial\phi_H$ imply dependence upon ϕ_H , which is hidden by integration over Bjorken x 's
- Strictly speaking, the (perturbative) result **is non-perturbative** ($\phi_0 \sim \exp(-1/\alpha_s)$)

NLOwPS: grcNLO

Partition the phase space as in standard slicing (i.e., define a non-soft, non collinear region ϕ_{NSC}), and subtract there the real counterterm:

$$\mathcal{F}_{\text{grcNLO}} = \sum_{ab} \int dx_1 dx_2 d\phi_3 f_a(x_1) f_b(x_2) \left[\mathcal{F}_{\text{MC}}^{(2 \rightarrow 3)} \left(\mathcal{M}_{ab}^{(r)}(x_1, x_2, \phi_3) - \mathcal{M}_{ab}^{(c.t.)}(x_1, x_2, \phi_3) \right) \Theta(\phi_3 \in \phi_{NSC}) + \mathcal{F}_{\text{MC}}^{(2 \rightarrow 2)} \mathcal{M}_{ab}^{(b,v,c)}(x_1, x_2, \phi_2) \right]$$



This formally coincides with MC@NLO, provided that ϕ_{NSC} is the full phase space, and

$$\mathcal{M}_{ab}^{(\text{MC})} \equiv \mathcal{M}_{ab}^{(c.t.)}$$

This condition cannot be imposed: it must result from the MC implementation



- + All matrix elements generated numerically
- Double counting if $\mathcal{M}_{ab}^{(\text{MC})}$ is not built ad hoc
- Condition on $\mathcal{M}_{ab}^{(\text{MC})}$ implies the construction of a new MC

What to expect from an NLOwPS (here MC@NLO)



- MC@NLO rate = NLO rate \implies K-factors are included **consistently**
- MC@NLO- and MC-predicted **shapes** are identical where MC does a good job
- $\mathcal{S}+0$ jet and $\mathcal{S}+1$ jet treated **exactly**, $\mathcal{S}+n$ jets ($n > 1$) better than in MC's
- No dependence on δ_{sep} \implies tuning is the same as in ordinary MC's
- Some **negative-weight events**, to be subtracted (rather than added) from histograms



Single-inclusive b at the Tevatron



No significant discrepancy with data

- No PTMIN dependence in MC@NLO \implies solid predictions down to $p_T = 0$, no “perturbative-parameter tuning” (more work on b hadronization parameters needed)
- Full agreement with NLL+NLO computation (FONLL, Cacciari&Nason), if the large dependence (at small p_T) on the hadronization scheme of the latter is taken into account

Is the agreement with the resummed result accidental?



The same happens with Higgs. The result of Bozzi, Catani, de Florian, Grazzini has a matching condition similar to MC@NLO, in that it conserves the total rate

- ◆ The agreement with the analytically-resummed result improves when the logarithmic accuracy of the latter is increased \longrightarrow Herwig has more logs than you expect
- ◆ We can now apply any cuts we like (decay products, recoiling system) – a fully realistic jet-veto analysis is doable
- ◆ **Beware:** vastly different from Pythia!

Conclusions

There has been substantial theoretical progress in MC's in the past three years or so. The timing is just right, since it's the Tevatron and the LHC that demand the construction of improved MC tools

MEC for multileg processes are firmly established

- Expect CKKW to become part of HERWIG, PYTHIA, and SHERPA releases
- Reliable estimates for many backgrounds to new physics

NLOwPS's improve NLO computations and MC simulations in several respects

- MC@NLO is numerically more stable than NLO computations
- Realistic final states, including hadronization, are part of NLO predictions
- NLOwPS's are the **only way** in which K -factors can be embedded into MC's
- Hard radiation is incorporated in MC's, without any kinematical distortion

NLOwPS/MEC work just like ordinary MC's, and must be seen as upgrades of the latter. So the answer to the question: **when do I have to use them?** is: **always**