

Filip Strobin

0-dimensional uncountable compact spaces as attractors of iterated function systems

(with Taras Banakh and Magdalena Nowak)

Institute of Mathematics,
Jan Kochanowski University in Kielce
and
Institute of Mathematics,
Łódź z University of Technology

Notation

Notation

Let (X, d) be a metric space.

- * By $\mathcal{K}(X)$ we denote the space of all nonempty and compact subsets of X .
- * We consider $\mathcal{K}(X)$ as a metric space with the Hausdorff metric H :

$$H(K, D) := \max \left\{ \sup_{x \in K} \left(\inf_{y \in D} d(x, y) \right), \sup_{y \in D} \left(\inf_{x \in K} d(x, y) \right) \right\}$$

- * If $S = (f_1, \dots, f_n)$ is a finite system of continuous selfmaps of X (i.e., $f_i : X \rightarrow X$), then by F_S we denote the function $F_S : \mathcal{K}(X) \rightarrow \mathcal{K}(X)$ defined by

$$F_S(K) := f_1(K) \cup \dots \cup f_n(K)$$

- * If $f : X \rightarrow X$, then by $Lip(f)$ we denote the Lipschitz constant of f .
- * If $Lip(f) < 1$, that is there exists $\alpha < 1$ such that

$$\forall x, y \in X \quad d(f(x), f(y)) \leq \alpha d(x, y),$$

then we say that f is a *Banach contraction*.

Notation

Notation

Let (X, d) be a metric space.

- * By $\mathcal{K}(X)$ we denote the space of all nonempty and compact subsets of X .
- * We consider $\mathcal{K}(X)$ as a metric space with the Hausdorff metric H :

$$H(K, D) := \max \left\{ \sup_{x \in K} \left(\inf_{y \in D} d(x, y) \right), \sup_{y \in D} \left(\inf_{x \in K} d(x, y) \right) \right\}$$

- * If $\mathcal{S} = (f_1, \dots, f_n)$ is a finite system of continuous selfmaps of X (i.e., $f_i : X \rightarrow X$), then by $F_{\mathcal{S}}$ we denote the function $F_{\mathcal{S}} : \mathcal{K}(X) \rightarrow \mathcal{K}(X)$ defined by

$$F_{\mathcal{S}}(K) := f_1(K) \cup \dots \cup f_n(K)$$

- * If $f : X \rightarrow X$, then by $Lip(f)$ we denote the Lipschitz constant of f .
- * If $Lip(f) < 1$, that is there exists $\alpha < 1$ such that

$$\forall x, y \in X \quad d(f(x), f(y)) \leq \alpha d(x, y),$$

then we say that f is a *Banach contraction*.

IFSs and their attractors

Theorem, Hutchinson, Barnsley, 1980's

Let X be a complete metric space and $\mathcal{S} = (f_1, \dots, f_n)$ be a finite system of Banach contractions of X . Then there exists a unique $A_{\mathcal{S}} \in \mathcal{K}(X)$ such that

$$A_{\mathcal{S}} = F_{\mathcal{S}}(A_{\mathcal{S}}) = f_1(A_{\mathcal{S}}) \cup \dots \cup f_n(A_{\mathcal{S}})$$

Moreover, for every $K \in \mathcal{K}(X)$, the sequence of iterates $F_{\mathcal{S}}^{(k)}(K) \rightarrow A_{\mathcal{S}}$ with respect to the Hausdorff metric.

Remark

In the above result it is enough to assume that each f_i is a weak contraction (Rakotch, Browder, Matkowski).

Fractals in the sense of Hutchinson and Barnsley

- * If a function system \mathcal{S} satisfies the thesis of the above theorem, then we say that \mathcal{S} *generates a fractal*.
- * A compact set $A_{\mathcal{S}}$ which is generated by a function system \mathcal{S} in this way is called *fractal* or *attractor in the sense of Hutchinson and Barnsley*.
- * In this setting, function systems are called *iterated function systems (IFSs)*.

IFSs and their attractors

Theorem, Hutchinson, Barnsley, 1980's

Let X be a complete metric space and $\mathcal{S} = (f_1, \dots, f_n)$ be a finite system of Banach contractions of X . Then there exists a unique $A_{\mathcal{S}} \in \mathcal{K}(X)$ such that

$$A_{\mathcal{S}} = F_{\mathcal{S}}(A_{\mathcal{S}}) = f_1(A_{\mathcal{S}}) \cup \dots \cup f_n(A_{\mathcal{S}})$$

Moreover, for every $K \in \mathcal{K}(X)$, the sequence of iterates $F_{\mathcal{S}}^{(k)}(K) \rightarrow A_{\mathcal{S}}$ with respect to the Hausdorff metric.

Remark

In the above result it is enough to assume that each f_i is a weak contraction (Rakotch, Browder, Matkowski).

Fractals in the sense of Hutchinson and Barnsley

- * If a function system \mathcal{S} satisfies the thesis of the above theorem, then we say that \mathcal{S} *generates a fractal*.
- * A compact set $A_{\mathcal{S}}$ which is generated by a function system \mathcal{S} in this way is called *fractal* or *attractor in the sense of Hutchinson and Barnsley*.
- * In this setting, function systems are called *iterated function systems* (IFSs).

IFSs and their attractors

Theorem, Hutchinson, Barnsley, 1980's

Let X be a complete metric space and $\mathcal{S} = (f_1, \dots, f_n)$ be a finite system of Banach contractions of X . Then there exists a unique $A_{\mathcal{S}} \in \mathcal{K}(X)$ such that

$$A_{\mathcal{S}} = F_{\mathcal{S}}(A_{\mathcal{S}}) = f_1(A_{\mathcal{S}}) \cup \dots \cup f_n(A_{\mathcal{S}})$$

Moreover, for every $K \in \mathcal{K}(X)$, the sequence of iterates $F_{\mathcal{S}}^{(k)}(K) \rightarrow A_{\mathcal{S}}$ with respect to the Hausdorff metric.

Remark

In the above result it is enough to assume that each f_i is a weak contraction (Rakotch, Browder, Matkowski).

Fractals in the sense of Hutchinson and Barnsley

- * If a function system \mathcal{S} satisfies the thesis of the above theorem, then we say that \mathcal{S} *generates a fractal*.
- * A compact set $A_{\mathcal{S}}$ which is generated by a function system \mathcal{S} in this way is called *fractal* or *attractor in the sense of Hutchinson and Barnsley*.
- * In this setting, function systems are called *iterated function systems* (IFSs).

IFSs and their attractors

Theorem, Hutchinson, Barnsley, 1980's

Let X be a complete metric space and $\mathcal{S} = (f_1, \dots, f_n)$ be a finite system of Banach contractions of X . Then there exists a unique $A_{\mathcal{S}} \in \mathcal{K}(X)$ such that

$$A_{\mathcal{S}} = F_{\mathcal{S}}(A_{\mathcal{S}}) = f_1(A_{\mathcal{S}}) \cup \dots \cup f_n(A_{\mathcal{S}})$$

Moreover, for every $K \in \mathcal{K}(X)$, the sequence of iterates $F_{\mathcal{S}}^{(k)}(K) \rightarrow A_{\mathcal{S}}$ with respect to the Hausdorff metric.

Remark

In the above result it is enough to assume that each f_i is a weak contraction (Rakotch, Browder, Matkowski).

Fractals in the sense of Hutchinson and Barnsley

- * If a function system \mathcal{S} satisfies the thesis of the above theorem, then we say that \mathcal{S} *generates a fractal*.
- * A compact set $A_{\mathcal{S}}$ which is generated by a function system \mathcal{S} in this way is called *fractal* or *attractor in the sense of Hutchinson and Barnsley*.
- * In this setting, function systems are called *iterated function systems (IFSs)*.

Major questions concerning Hutchinson-Barnsley theory

Major questions

- (Q1) Which iterated function systems generate fractals?
- (Q2) What is the structure of fractals generated by IFSs?
- (Q3) Which compact spaces are fractals generated by IFSs?

Variants of (Q3)

- (3a) Which compact metric spaces are fractals generated by IFSs consisting of Banach contractions?
- (3b) Which compact metric spaces are fractals generated by IFSs consisting of weak contractions?
- (3c) Which compact metric spaces are homeomorphic to fractals generated by IFSs?
- (3c') Which compact metric spaces X have the following property:
There is an IFS $\mathcal{S} = (f_1, \dots, f_n)$ such that $X = F_{\mathcal{S}}(X) = f_1(X) \cup \dots \cup f_n(X)$,
and an admissible metric d on X so that each f_i is a weak (or Banach) contraction with respect to d .

Major questions concerning Hutchinson-Barnsley theory

Major questions

- (Q1) Which iterated function systems generate fractals?
- (Q2) What is the structure of fractals generated by IFSs?
- (Q3) Which compact spaces are fractals generated by IFSs?

Variants of (Q3)

- (3a) Which compact metric spaces are fractals generated by IFSs consisting of Banach contractions?
- (3b) Which compact metric spaces are fractals generated by IFSs consisting of weak contractions?
- (3c) Which compact metric spaces are homeomorphic to fractals generated by IFSs?
- (3c') Which compact metric spaces X have the following property:
There is an IFS $\mathcal{S} = (f_1, \dots, f_n)$ such that $X = F_{\mathcal{S}}(X) = f_1(X) \cup \dots \cup f_n(X)$,
and an admissible metric d on X so that each f_i is a weak (or Banach) contraction with respect to d .

Major questions concerning Hutchinson-Barnsley theory

Major questions

- (Q1) Which iterated function systems generate fractals?
- (Q2) What is the structure of fractals generated by IFSs?
- (Q3) Which compact spaces are fractals generated by IFSs?

Variants of (Q3)

- (3a) Which compact metric spaces are fractals generated by IFSs consisting of Banach contractions?
- (3b) Which compact metric spaces are fractals generated by IFSs consisting of weak contractions?
- (3c) Which compact metric spaces are homeomorphic to fractals generated by IFSs?
- (3c') Which compact metric spaces X have the following property:
There is an IFS $\mathcal{S} = (f_1, \dots, f_n)$ such that $X = F_{\mathcal{S}}(X) = f_1(X) \cup \dots \cup f_n(X)$,
and an admissible metric d on X so that each f_i is a weak (or Banach) contraction with respect to d .

Major questions and known results

(Q3)

Which compact spaces are fractals generated by IFSs?

Known results

- * (Crovisier and Rams [CR] 2006, S. [S] 2013): There is a Cantor set $C \subset \mathbb{R}$ which is not an attractor of any IFS consisting of weak contractions.
- * (Sanders [Sa] 2002) An arc $A \subset \mathbb{R}^n$ is the attractor of an IFS consisting of Banach contractions on \mathbb{R}^n iff its variation $V(A) < \infty$.
- * (Banach, Nowak [BN] 2013): There is an 1-dimensional Peano continuum which not homeomorphic to the attractor of any IFS consisting of Banach contractions.
- * (Nowak [N] 2013): For a countable and compact space X , TFCAE:
 - (i) X is homeomorphic to the attractor of a certain IFS consisting of Banach contractions;
 - (ii) X is homeomorphic to the attractor of a certain IFS consisting of weak contractions;
 - (iii) the scattered height of X is a successor ordinal.

Major questions and known results

(Q3)

Which compact spaces are fractals generated by IFSs?

Known results

- * (Crovisier and Rams [CR] 2006, S. [S] 2013): There is a Cantor set $C \subset \mathbb{R}$ which is not an attractor of any IFS consisting of weak contractions.
- * (Sanders [Sa] 2002) An arc $A \subset \mathbb{R}^n$ is the attractor of an IFS consisting of Banach contractions on \mathbb{R}^n iff its variation $V(A) < \infty$.
- * (Banach, Nowak [BN] 2013): There is an 1-dimensional Peano continuum which not homeomorphic to the attractor of any IFS consisting of Banach contractions.
- * (Nowak [N] 2013): For a countable and compact space X , TFCAE:
 - (i) X is homeomorphic to the attractor of a certain IFS consisting of Banach contractions;
 - (ii) X is homeomorphic to the attractor of a certain IFS consisting of weak contractions;
 - (iii) the scattered height of X is a successor ordinal.

Major questions and known results

(Q3)

Which compact spaces are fractals generated by IFSs?

Known results

- * (Crovisier and Rams [CR] 2006, S. [S] 2013): There is a Cantor set $C \subset \mathbb{R}$ which is not an attractor of any IFS consisting of weak contractions.
- * (Sanders [Sa] 2002) An arc $A \subset \mathbb{R}^n$ is the attractor of an IFS consisting of Banach contractions on \mathbb{R}^n iff its variation $V(A) < \infty$.
- * (Banach, Nowak [BN] 2013): There is an 1-dimensional Peano continuum which not homeomorphic to the attractor of any IFS consisting of Banach contractions.
- * (Nowak [N] 2013): For a countable and compact space X , TFCAE:
 - (i) X is homeomorphic to the attractor of a certain IFS consisting of Banach contractions;
 - (ii) X is homeomorphic to the attractor of a certain IFS consisting of weak contractions;
 - (iii) the scattered height of X is a successor ordinal.

Main result

Theorem 1, Banach, Nowak, S., 2014

Let X be an uncountable, 0-dimensional compact space. Then there is an IFS $\mathcal{S} = (f, g)$ such that $X = F_{\mathcal{S}}(X) = f(X) \cup g(X)$, and for every $c \in (0, 1)$ there is an admissible metric d on X such that $Lip(f), Lip(g) \leq c$.

Remark

Recall that there is a Cantor set on \mathbb{R} which is not the attractor of an IFS on consisting of weak contractions.

Corollary

If X is an uncountable compact metrizable space, then TFCAE:

- (i) X is 0-dimensional;
- (ii) X satisfies the thesis of Theorem 1;
- (iii) X is homeomorphic to an IFS $\mathcal{S} = (f_1, \dots, f_n)$ with $\sum_{i=1}^n Lip(f_i) < 1$.

Theorem, Daniello, Steele, 2014

Each compact uncountable nowhere dense subset of \mathbb{R} is homeomorphic to the attractor of an IFS consisting of Banach contractions.

Main result

Theorem 1, Banach, Nowak, S., 2014

Let X be an uncountable, 0-dimensional compact space. Then there is an IFS $\mathcal{S} = (f, g)$ such that $X = F_{\mathcal{S}}(X) = f(X) \cup g(X)$, and for every $c \in (0, 1)$ there is an admissible metric d on X such that $Lip(f), Lip(g) \leq c$.

Remark

Recall that there is a Cantor set on \mathbb{R} which is not the attractor of an IFS on consisting of weak contractions.

Corollary

If X is an uncountable compact metrizable space, then TFCAE:

- (i) X is 0-dimensional;
- (ii) X satisfies the thesis of Theorem 1;
- (iii) X is homeomorphic to an IFS $\mathcal{S} = (f_1, \dots, f_n)$ with $\sum_{i=1}^n Lip(f_i) < 1$.

Theorem, Daniello, Steele, 2014

Each compact uncountable nowhere dense subset of \mathbb{R} is homeomorphic to the attractor of an IFS consisting of Banach contractions.

Main result

Theorem 1, Banach, Nowak, S., 2014

Let X be an uncountable, 0-dimensional compact space. Then there is an IFS $\mathcal{S} = (f, g)$ such that $X = F_{\mathcal{S}}(X) = f(X) \cup g(X)$, and for every $c \in (0, 1)$ there is an admissible metric d on X such that $Lip(f), Lip(g) \leq c$.

Remark

Recall that there is a Cantor set on \mathbb{R} which is not the attractor of an IFS on consisting of weak contractions.

Corollary

If X is an uncountable compact metrizable space, then TFCAE:

- (i) X is 0-dimensional;
- (ii) X satisfies the thesis of Theorem 1;
- (iii) X is homeomorphic to an IFS $\mathcal{S} = (f_1, \dots, f_n)$ with $\sum_{i=1}^n Lip(f_i) < 1$.

Theorem, Daniello, Steele, 2014

Each compact uncountable nowhere dense subset of \mathbb{R} is homeomorphic to the attractor of an IFS consisting of Banach contractions.

Main result

Theorem 1, Banach, Nowak, S., 2014

Let X be an uncountable, 0-dimensional compact space. Then there is an IFS $\mathcal{S} = (f, g)$ such that $X = F_{\mathcal{S}}(X) = f(X) \cup g(X)$, and for every $c \in (0, 1)$ there is an admissible metric d on X such that $Lip(f), Lip(g) \leq c$.

Remark

Recall that there is a Cantor set on \mathbb{R} which is not the attractor of an IFS on consisting of weak contractions.

Corollary

If X is an uncountable compact metrizable space, then TFCAE:

- (i) X is 0-dimensional;
- (ii) X satisfies the thesis of Theorem 1;
- (iii) X is homeomorphic to an IFS $\mathcal{S} = (f_1, \dots, f_n)$ with $\sum_{i=1}^n Lip(f_i) < 1$.

Theorem, Daniello, Steele, 2014

Each compact uncountable nowhere dense subset of \mathbb{R} is homeomorphic to the attractor of an IFS consisting of Banach contractions.

Main result

Theorem 1, Banach, Nowak, S., 2014

Let X be an uncountable, 0-dimensional compact space. Then there is an IFS $\mathcal{S} = (f, g)$ such that $X = F_{\mathcal{S}}(X) = f(X) \cup g(X)$, and for every $c \in (0, 1)$ there is an admissible metric d on X such that $Lip(f), Lip(g) \leq c$.

Remark

Recall that there is a Cantor set on \mathbb{R} which is not the attractor of an IFS on consisting of weak contractions.

Corollary

If X is an uncountable compact metrizable space, then TFCAE:

- (i) X is 0-dimensional;
- (ii) X satisfies the thesis of Theorem 1;
- (iii) X is homeomorphic to an IFS $\mathcal{S} = (f_1, \dots, f_n)$ with $\sum_{i=1}^n Lip(f_i) < 1$.

Theorem, Daniello, Steele, 2014

Each compact uncountable nowhere dense subset of \mathbb{R} is homeomorphic to the attractor of an IFS consisting of Banach contractions.

Thank you for your attention

Bibliography

- [BN] T. Banach, M. Nowak, *A 1-dimensional Peano continuum which is not an IFS attractor*, Proc. Amer. Math. Soc. **141**:3 (2013), 931–935.
- [B] M.F. Barnsley, *Fractals everywhere*. Academic Press Professional, Boston, MA, 1993.
- [CR] S. Crovisier, M. Rams, *IFS attractors and Cantor sets*, Topology Appl. **153** (2006), no. 11, 18491859.
- [H] J Hutchinson, *Fractals and self-similarity*. Indiana Univ. Math. J. **30**:5 (1981), 713–747.
- [Kw] M. Kwieciński, *A locally connected continuum which is not an IFS attractor*, Bull. Polish Acad. Sci. Math. **47** (1999), no. 2, 127132.
- [N] M. Nowak, *Topological classification of scattered IFS-attractors*, Topology Appl. **160** (2013), no. 14, 18891901.
- [NS] M. Nowak, T. Szarek *The shark teeth is a topological IFS-attractor*, submitted, arxiv: 1305.4492, 2013
- [Sa] M. Sanders, *An n -Cell in \mathbb{R}^{n+1} that is not the attractor of any IFS on \mathbb{R}^{n+1}* , Missouri J. Math. Sci. **21**, Issue 1 (2009), 13-20.
- [S] F. Strobil, *Attractors of generalized IFSs that are not attractors of IFSs*, Attractors of generalized IFSs that are not attractors of IFSs, DOI: 10.1016/j.jmaa.2014.08.029.