


Symmetrization and convexity I


Szymon Wąsowicz


(joint work with A. Witkowski)


University of Bielsko-Biała, Poland


The 28th International Summer Conference on Real
Functions Theory, Stara Lesna (Slovakia, 08.31–09.05, 2014)











Properties

Properties

- $g(x) = f(x) + f(a + b - x)$ – the symmetrization of f

Properties

- $g(x) = f(x) + f(a + b - x)$ – the symmetrization of f
- g is symmetric wrt. the midpoint $\frac{a+b}{2}$

Properties

- $g(x) = f(x) + f(a + b - x)$ – the symmetrization of f
- g is symmetric wrt. the midpoint $\frac{a+b}{2}$
- $g(a) = g(b) = f(a) + f(b)$ and $g(\frac{a+b}{2}) = 2f(\frac{a+b}{2})$

Properties

- $g(x) = f(x) + f(a + b - x)$ – the symmetrization of f
- g is symmetric wrt. the midpoint $\frac{a+b}{2}$
- $g(a) = g(b) = f(a) + f(b)$ and $g(\frac{a+b}{2}) = 2f(\frac{a+b}{2})$
- $\int_a^b g(x) dx = 2 \int_a^b f(x) dx$

Properties

- $g(x) = f(x) + f(a + b - x)$ – the symmetrization of f
- g is symmetric wrt. the midpoint $\frac{a+b}{2}$
- $g(a) = g(b) = f(a) + f(b)$ and $g(\frac{a+b}{2}) = 2f(\frac{a+b}{2})$
- $\int_a^b g(x) dx = 2 \int_a^b f(x) dx$

Symmetrization of a convex function

Suppose f is convex on $[a, b]$.

Properties

- $g(x) = f(x) + f(a + b - x)$ – the symmetrization of f
- g is symmetric wrt. the midpoint $\frac{a+b}{2}$
- $g(a) = g(b) = f(a) + f(b)$ and $g(\frac{a+b}{2}) = 2f(\frac{a+b}{2})$
- $\int_a^b g(x) dx = 2 \int_a^b f(x) dx$

Symmetrization of a convex function

Suppose f is convex on $[a, b]$.

- g is convex

Properties

- $g(x) = f(x) + f(a + b - x)$ – the symmetrization of f
- g is symmetric wrt. the midpoint $\frac{a+b}{2}$
- $g(a) = g(b) = f(a) + f(b)$ and $g(\frac{a+b}{2}) = 2f(\frac{a+b}{2})$
- $\int_a^b g(x) dx = 2 \int_a^b f(x) dx$

Symmetrization of a convex function

Suppose f is convex on $[a, b]$.

- g is convex
- $g(\frac{a+b}{2}) = 2f(\frac{a+b}{2}) \leq f(x) + f(a + b - x) = g(x) \leq g(a)$

Hermite–Hadamard inequality by symmetrization

Suppose f is convex on $[a, b]$.

Hermite–Hadamard inequality by symmetrization

Suppose f is convex on $[a, b]$.

- $g\left(\frac{a+b}{2}\right) \leq g(x) \leq g(a)$

Hermite–Hadamard inequality by symmetrization

Suppose f is convex on $[a, b]$.

- $g\left(\frac{a+b}{2}\right) \leq g(x) \leq g(a)$
- $(b-a)g\left(\frac{a+b}{2}\right) \leq \int_a^b g(x) dx \leq (b-a)g(a)$

Hermite–Hadamard inequality by symmetrization

Suppose f is convex on $[a, b]$.

- $g\left(\frac{a+b}{2}\right) \leq g(x) \leq g(a)$
- $(b-a)g\left(\frac{a+b}{2}\right) \leq \int_a^b g(x) dx \leq (b-a)g(a)$
- $2(b-a)f\left(\frac{a+b}{2}\right) \leq \int_a^b (f(x) + f(a+b-x)) dx$
 $\leq (b-a)(f(a) + f(b))$

Hermite–Hadamard inequality by symmetrization


Suppose f is convex on $[a, b]$.

- $g\left(\frac{a+b}{2}\right) \leq g(x) \leq g(a)$
- $(b-a)g\left(\frac{a+b}{2}\right) \leq \int_a^b g(x) dx \leq (b-a)g(a)$
- $2(b-a)f\left(\frac{a+b}{2}\right) \leq \int_a^b (f(x) + f(a+b-x)) dx$
 $\leq (b-a)(f(a) + f(b))$
- $2(b-a)f\left(\frac{a+b}{2}\right) \leq 2 \int_a^b f(x) dx \leq (b-a)(f(a) + f(b))$


Hermite–Hadamard inequality by symmetrization

Suppose f is convex on $[a, b]$.

- $g\left(\frac{a+b}{2}\right) \leq g(x) \leq g(a)$
- $(b-a)g\left(\frac{a+b}{2}\right) \leq \int_a^b g(x) dx \leq (b-a)g(a)$
- $2(b-a)f\left(\frac{a+b}{2}\right) \leq \int_a^b (f(x) + f(a+b-x)) dx \leq (b-a)(f(a) + f(b))$
- $2(b-a)f\left(\frac{a+b}{2}\right) \leq 2 \int_a^b f(x) dx \leq (b-a)(f(a) + f(b))$
- $f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}$



$$\int_a^b g(x) dx - (b-a)g\left(\frac{a+b}{2}\right) \leq S$$



$$\int_a^b g(x) dx - (b-a)g\left(\frac{a+b}{2}\right) \leq S \leq (b-a)g(a) - \int_a^b g(x) dx$$

LHH estimates the integral MV better than RHH

Suppose f is convex on $[a, b]$.

LHH estimates the integral MV better than RHH

Suppose f is convex on $[a, b]$.

- $f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}$ HH inequality

LHH estimates the integral MV better than RHH

Suppose f is convex on $[a, b]$.

- $f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}$ HH inequality
- $0 \leq \frac{1}{b-a} \int_a^b f(x) dx - f\left(\frac{a+b}{2}\right)$ LHH

LHH estimates the integral MV better than RHH

Suppose f is convex on $[a, b]$.

- $f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}$ HH inequality
- $0 \leq \frac{1}{b-a} \int_a^b f(x) dx - f\left(\frac{a+b}{2}\right)$ LHH
- $0 \leq \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx$ RHH

LHH estimates the integral MV better than RHH

Suppose f is convex on $[a, b]$.

- $f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a) + f(b)}{2}$ HH inequality
- $0 \leq \frac{1}{b-a} \int_a^b f(x) dx - f\left(\frac{a+b}{2}\right)$ LHH
- $0 \leq \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(x) dx$ RHH
- $f \text{ convex on } [a, b] \implies 0 \leq \text{LHH} \leq \text{RHH}$

Proof by symmetrization

Proof by symmetrization

- $0 \leq \int_a^b g(x) dx - (b-a)g\left(\frac{a+b}{2}\right) \leq (b-a)g(a) - \int_a^b g(x) dx$


Proof by symmetrization


- $0 \leq \int_a^b g(x) dx - (b-a)g\left(\frac{a+b}{2}\right) \leq (b-a)g(a) - \int_a^b g(x) dx$
- $0 \leq 2 \int_a^b f(x) dx - 2(b-a)f\left(\frac{a+b}{2}\right)$
 $\leq (b-a)(f(a) + f(b)) - 2 \int_a^b f(x) dx$


Proof by symmetrization


- $0 \leq \int_a^b g(x) dx - (b-a)g\left(\frac{a+b}{2}\right) \leq (b-a)g(a) - \int_a^b g(x) dx$
- $0 \leq 2 \int_a^b f(x) dx - 2(b-a)f\left(\frac{a+b}{2}\right)$
 $\leq (b-a)(f(a) + f(b)) - 2 \int_a^b f(x) dx$
- Divide both sides by $2(b-a)$.


The inequality $0 \leq \text{LHH} \leq \text{RHH}$ is optimal




















Properties

Properties

- $F(x) = f(\sigma_0(x)) + f(\sigma_1(x)) + f(\sigma_2(x))$ – symmetrization of f

Properties

- $F(x) = f(\sigma_0(x)) + f(\sigma_1(x)) + f(\sigma_2(x))$ – symmetrization of f
- F is symmetric wrt. the barycenter b , i.e. $F(\sigma_i(x)) = F(x)$, $x \in S$, $i = 0, 1, 2$.

Properties

- $F(x) = f(\sigma_0(x)) + f(\sigma_1(x)) + f(\sigma_2(x))$ – symmetrization of f
- F is symmetric wrt. the barycenter b , i.e. $F(\sigma_i(x)) = F(x)$, $x \in S$, $i = 0, 1, 2$.
- $F(v_i) = f(v_0) + f(v_1) + f(v_2)$, $i = 0, 1, 2$ and $F(b) = 3f(b)$

Properties

- $F(x) = f(\sigma_0(x)) + f(\sigma_1(x)) + f(\sigma_2(x))$ – symmetrization of f
- F is symmetric wrt. the barycenter b , i.e. $F(\sigma_i(x)) = F(x)$, $x \in S$, $i = 0, 1, 2$.
- $F(v_i) = f(v_0) + f(v_1) + f(v_2)$, $i = 0, 1, 2$ and $F(b) = 3f(b)$
- $\int_S f(\sigma_i(x)) dx = \int_S f(x) dx$, $i = 0, 1, 2$

Properties

- $F(x) = f(\sigma_0(x)) + f(\sigma_1(x)) + f(\sigma_2(x))$ – symmetrization of f
- F is symmetric wrt. the barycenter b , i.e. $F(\sigma_i(x)) = F(x)$, $x \in S$, $i = 0, 1, 2$.
- $F(v_i) = f(v_0) + f(v_1) + f(v_2)$, $i = 0, 1, 2$ and $F(b) = 3f(b)$
- $\int_S f(\sigma_i(x)) \, dx = \int_S f(x) \, dx$, $i = 0, 1, 2$
- $\int_S F(x) \, dx = 3 \int_S f(x) \, dx$

Properties

- $F(x) = f(\sigma_0(x)) + f(\sigma_1(x)) + f(\sigma_2(x))$ – symmetrization of f
- F is symmetric wrt. the barycenter b , i.e. $F(\sigma_i(x)) = F(x)$, $x \in S$, $i = 0, 1, 2$.
- $F(v_i) = f(v_0) + f(v_1) + f(v_2)$, $i = 0, 1, 2$ and $F(b) = 3f(b)$
- $\int_S f(\sigma_i(x)) \, dx = \int_S f(x) \, dx$, $i = 0, 1, 2$
- $\int_S F(x) \, dx = 3 \int_S f(x) \, dx$
- f is convex on $S \implies F$ is convex on S

Hermite–Hadamard inequality

$S \subset \mathbb{R}^2$ – a simplex (a triangle) with vertices v_0, v_1, v_2 and barycenter \mathbf{b} , $f : S \rightarrow \mathbb{R}$ – a convex function

$$f(\mathbf{b}) \leq \frac{1}{\text{vol}(S)} \int_S f(\mathbf{x}) \, d\mathbf{x} \leq \frac{f(v_0) + f(v_1) + f(v_2)}{3}$$

Hermite–Hadamard inequality

$S \subset \mathbb{R}^2$ – a simplex (a triangle) with vertices v_0, v_1, v_2 and barycenter \mathbf{b} , $f : S \rightarrow \mathbb{R}$ – a convex function

$$f(\mathbf{b}) \leq \frac{1}{\text{vol}(S)} \int_S f(\mathbf{x}) \, d\mathbf{x} \leq \frac{f(v_0) + f(v_1) + f(v_2)}{3}$$

The inequality between LHH and RHH terms

Hermite–Hadamard inequality

$S \subset \mathbb{R}^2$ – a simplex (a triangle) with vertices v_0, v_1, v_2 and barycenter \mathbf{b} , $f : S \rightarrow \mathbb{R}$ – a convex function

$$f(\mathbf{b}) \leq \frac{1}{\text{vol}(S)} \int_S f(\mathbf{x}) \, d\mathbf{x} \leq \frac{f(v_0) + f(v_1) + f(v_2)}{3}$$

The inequality between LHH and RHH terms

$$\bullet \quad \frac{1}{\text{vol}(S)} \int_S f(\mathbf{x}) \, d\mathbf{x} - f(\mathbf{b}) \qquad \text{LHH}$$

Hermite–Hadamard inequality

$S \subset \mathbb{R}^2$ – a simplex (a triangle) with vertices v_0, v_1, v_2 and barycenter \mathbf{b} , $f : S \rightarrow \mathbb{R}$ – a convex function

$$f(\mathbf{b}) \leq \frac{1}{\text{vol}(S)} \int_S f(\mathbf{x}) \, d\mathbf{x} \leq \frac{f(v_0) + f(v_1) + f(v_2)}{3}$$

The inequality between LHH and RHH terms

- $\frac{1}{\text{vol}(S)} \int_S f(\mathbf{x}) \, d\mathbf{x} - f(\mathbf{b})$ LHH
- $\frac{f(v_0) + f(v_1) + f(v_2)}{3} - \frac{1}{\text{vol}(S)} \int_S f(\mathbf{x}) \, d\mathbf{x}$ RHH

Hermite–Hadamard inequality

$S \subset \mathbb{R}^2$ – a simplex (a triangle) with vertices v_0, v_1, v_2 and barycenter \mathbf{b} , $f : S \rightarrow \mathbb{R}$ – a convex function


$$f(\mathbf{b}) \leq \frac{1}{\text{vol}(S)} \int_S f(\mathbf{x}) \, d\mathbf{x} \leq \frac{f(v_0) + f(v_1) + f(v_2)}{3}$$

The inequality between LHH and RHH terms


- $\frac{1}{\text{vol}(S)} \int_S f(\mathbf{x}) \, d\mathbf{x} - f(\mathbf{b})$ LHH
- $\frac{f(v_0) + f(v_1) + f(v_2)}{3} - \frac{1}{\text{vol}(S)} \int_S f(\mathbf{x}) \, d\mathbf{x}$ RHH
- $f : S \rightarrow \mathbb{R}$ convex $\implies 0 \leq \text{LHH} \leq 2 \text{RHH}$

The proof goes by
symmetrization F.

The proof goes by
symmetrization F.




The proof goes by
symmetrization F.



The proof goes by
symmetrization F .

- Volume between the base and the surface

$$\begin{aligned} 0 &\leq \int_S F(x) \, dx - F(\mathbf{b}) \cdot \text{area}(\text{base}) \\ &= \text{area}(\text{base}) \cdot \text{LHH}(F) \end{aligned}$$




The proof goes by
symmetrization F .


- Volume between the base and the surface

$$\begin{aligned} 0 &\leq \int_S F(x) \, dx - F(\mathbf{b}) \cdot \text{area}(\text{base}) \\ &= \text{area}(\text{base}) \cdot \text{LHH}(F) \end{aligned}$$

- $\text{area}(\text{base}) \cdot \text{LHH}(F) \leq 2 \, \text{vol}(\text{pyramid})$



The proof goes by
symmetrization F .




- Volume between the base and the surface

$$\begin{aligned} 0 &\leq \int_S F(x) \, dx - F(\mathbf{b}) \cdot \text{area}(\text{base}) \\ &= \text{area}(\text{base}) \cdot \text{LHH}(F) \end{aligned}$$

- $\text{area}(\text{base}) \cdot \text{LHH}(F) \leq 2 \, \text{vol}(\text{pyramid})$
- $\text{vol}(\text{pyramid}) \leq \text{volume between the surface and the upper base of a prism}$

The proof goes by
symmetrization F .




- Volume between the base and the surface

$$\begin{aligned} 0 &\leq \int_S F(x) \, dx - F(\mathbf{b}) \cdot \text{area}(\text{base}) \\ &= \text{area}(\text{base}) \cdot \text{LHH}(F) \end{aligned}$$

- $\text{area}(\text{base}) \cdot \text{LHH}(F) \leq 2 \, \text{vol}(\text{pyramid})$
- $\text{vol}(\text{pyramid}) \leq \text{volume between the surface and the upper base of a prism}$

$$\text{vol}(\text{pyramid}) \leq \text{vol}(\text{prism}) - \int_S F(x) \, dx$$

The proof goes by
symmetrization F .




- Volume between the base and the surface

$$\begin{aligned} 0 &\leq \int_S F(x) \, dx - F(\mathbf{b}) \cdot \text{area}(\text{base}) \\ &= \text{area}(\text{base}) \cdot \text{LHH}(F) \end{aligned}$$

- $\text{area}(\text{base}) \cdot \text{LHH}(F) \leq 2 \, \text{vol}(\text{pyramid})$
- $\text{vol}(\text{pyramid}) \leq \text{volume between the surface and the upper base of a prism}$

$$\begin{aligned} \text{vol}(\text{pyramid}) &\leq \text{vol}(\text{prism}) - \int_S F(x) \, dx \\ &= \text{area}(\text{base}) \cdot F(v_0) - \int_S F(x) \, dx \end{aligned}$$

The proof goes by
symmetrization F .




- Volume between the base and the surface

$$\begin{aligned} 0 &\leq \int_S F(x) \, dx - F(\mathbf{b}) \cdot \text{area}(\text{base}) \\ &= \text{area}(\text{base}) \cdot \text{LHH}(F) \end{aligned}$$

- $\text{area}(\text{base}) \cdot \text{LHH}(F) \leq 2 \, \text{vol}(\text{pyramid})$
- $\text{vol}(\text{pyramid}) \leq \text{volume between the surface and the upper base of a prism}$

$$\begin{aligned} \text{vol}(\text{pyramid}) &\leq \text{vol}(\text{prism}) - \int_S F(x) \, dx \\ &= \text{area}(\text{base}) \cdot F(v_0) - \int_S F(x) \, dx \\ &= \text{vol}(S) \left(F(v_0) - \frac{1}{\text{vol}(S)} \int_S F(x) \, dx \right) \end{aligned}$$

The proof goes by
symmetrization F .



- Volume between the base and the surface

$$\begin{aligned} 0 &\leq \int_S F(x) \, dx - F(\mathbf{b}) \cdot \text{area}(\text{base}) \\ &= \text{area}(\text{base}) \cdot \text{LHH}(F) \end{aligned}$$

- $\text{area}(\text{base}) \cdot \text{LHH}(F) \leq 2 \, \text{vol}(\text{pyramid})$
- $\text{vol}(\text{pyramid}) \leq \text{volume between the surface and the upper base of a prism}$

$$\begin{aligned} \text{vol}(\text{pyramid}) &\leq \text{vol}(\text{prism}) - \int_S F(x) \, dx \\ &= \text{area}(\text{base}) \cdot F(v_0) - \int_S F(x) \, dx \\ &= \text{vol}(S) \left(F(v_0) - \frac{1}{\text{vol}(S)} \int_S F(x) \, dx \right) \\ &= \text{vol}(S) \cdot \text{RHH}(F) \end{aligned}$$

- Hence $0 \leq \text{LHH}(F) \leq 2 \text{RHH}(F)$.

- Hence $0 \leq \text{LHH}(F) \leq 2 \text{RHH}(F)$.
- By the properties of symmetrization it is trivial to check that this implies $0 \leq \text{LHH}(f) \leq 2 \text{RHH}(f)$.

- Hence $0 \leq \text{LHH}(F) \leq 2 \text{RHH}(F)$.
- By the properties of symmetrization it is trivial to check that this implies $0 \leq \text{LHH}(f) \leq 2 \text{RHH}(f)$.
- The surface of a pyramid is a graph of a convex function.

- Hence $0 \leq \text{LHH}(F) \leq 2 \text{RHH}(F)$.
- By the properties of symmetrization it is trivial to check that this implies $0 \leq \text{LHH}(f) \leq 2 \text{RHH}(f)$.
- The surface of a pyramid is a graph of a convex function.
- Moreover, in this case we get equality above.

- Hence $0 \leq \text{LHH}(F) \leq 2 \text{RHH}(F)$.
- By the properties of symmetrization it is trivial to check that this implies $0 \leq \text{LHH}(f) \leq 2 \text{RHH}(f)$.
- The surface of a pyramid is a graph of a convex function.
- Moreover, in this case we get equality above.
- That is why the inequality in question is optimal.

- $v_0, \dots, v_n \in \mathbb{R}^n$ affinely independent

- $v_0, \dots, v_n \in \mathbb{R}^n$ affinely independent
- $S = \text{conv}\{v_0, \dots, v_n\}$ – a simplex with barycenter

$$b = \frac{1}{n+1} \sum_{i=0}^n v_i$$

- $v_0, \dots, v_n \in \mathbb{R}^n$ affinely independent
- $S = \text{conv}\{v_0, \dots, v_n\}$ – a simplex with barycenter

$$b = \frac{1}{n+1} \sum_{i=0}^n v_i$$

- $S \ni x = \sum_{i=0}^n t_i v_i$ uniquely with $t_i \geq 0$, $t_0 + \dots + t_n = 1$

- $v_0, \dots, v_n \in \mathbb{R}^n$ affinely independent
- $S = \text{conv}\{v_0, \dots, v_n\}$ – a simplex with barycenter

$$b = \frac{1}{n+1} \sum_{i=0}^n v_i$$

- $S \ni x = \sum_{i=0}^n t_i v_i$ uniquely with $t_i \geq 0$, $t_0 + \dots + t_n = 1$
- C – the set of all cyclic permutations of $\{0, \dots, n\}$

- $v_0, \dots, v_n \in \mathbb{R}^n$ affinely independent
- $S = \text{conv}\{v_0, \dots, v_n\}$ – a simplex with barycenter
$$b = \frac{1}{n+1} \sum_{i=0}^n v_i$$
- $S \ni x = \sum_{i=0}^n t_i v_i$ uniquely with $t_i \geq 0$, $t_0 + \dots + t_n = 1$
- C – the set of all cyclic permutations of $\{0, \dots, n\}$
- We identify $\sigma \in C$ with the affine transformation $\sigma : S \rightarrow S$ given by

$$\sigma\left(\sum_{i=0}^n t_i v_i\right) = \sum_{i=0}^n t_{\sigma(i)} v_i$$

Return for a while to $n = 2$.

$$\sigma_0 = \begin{pmatrix} 0 & 1 & 2 \\ 0 & 1 & 2 \end{pmatrix}, \quad \sigma_1 = \begin{pmatrix} 0 & 1 & 2 \\ 1 & 2 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & 1 & 2 \\ 2 & 0 & 1 \end{pmatrix}, \quad C = \{\sigma_0, \sigma_1, \sigma_2\}$$

Return for a while to $n = 2$.

$$\sigma_0 = \begin{pmatrix} 0 & 1 & 2 \\ 0 & 1 & 2 \end{pmatrix}, \quad \sigma_1 = \begin{pmatrix} 0 & 1 & 2 \\ 1 & 2 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & 1 & 2 \\ 2 & 0 & 1 \end{pmatrix}, \quad C = \{\sigma_0, \sigma_1, \sigma_2\}$$

$$\sigma_0(t_0v_0 + t_1v_1 + t_2v_2) = t_0v_0 + t_1v_1 + t_2v_2$$

Return for a while to $n = 2$.

$$\sigma_0 = \begin{pmatrix} 0 & 1 & 2 \\ 0 & 1 & 2 \end{pmatrix}, \quad \sigma_1 = \begin{pmatrix} 0 & 1 & 2 \\ 1 & 2 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & 1 & 2 \\ 2 & 0 & 1 \end{pmatrix}, \quad C = \{\sigma_0, \sigma_1, \sigma_2\}$$

$$\sigma_0(t_0v_0 + t_1v_1 + t_2v_2) = t_0v_0 + t_1v_1 + t_2v_2$$

$$\sigma_1(t_0v_0 + t_1v_1 + t_2v_2) = t_1v_0 + t_2v_1 + t_0v_2$$

Return for a while to $n = 2$.

$$\sigma_0 = \begin{pmatrix} 0 & 1 & 2 \\ 0 & 1 & 2 \end{pmatrix}, \quad \sigma_1 = \begin{pmatrix} 0 & 1 & 2 \\ 1 & 2 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & 1 & 2 \\ 2 & 0 & 1 \end{pmatrix}, \quad C = \{\sigma_0, \sigma_1, \sigma_2\}$$

$$\sigma_0(t_0v_0 + t_1v_1 + t_2v_2) = t_0v_0 + t_1v_1 + t_2v_2$$

$$\sigma_1(t_0v_0 + t_1v_1 + t_2v_2) = t_1v_0 + t_2v_1 + t_0v_2$$

$$\sigma_2(t_0v_0 + t_1v_1 + t_2v_2) = t_2v_0 + t_0v_1 + t_1v_2$$

For $f : S \rightarrow \mathbb{R}$ and $\sigma \in C$ put $f_\sigma(x) = f(\sigma(x))$

For $f : S \rightarrow \mathbb{R}$ and $\sigma \in C$ put $f_\sigma(x) = f(\sigma(x))$

Symmetrization and its properties

$$F : S \rightarrow \mathbb{R}, \quad F = \sum_{\sigma \in C} f_\sigma \quad \text{symmetrization of } f : S \rightarrow \mathbb{R}$$

For $f : S \rightarrow \mathbb{R}$ and $\sigma \in C$ put $f_\sigma(x) = f(\sigma(x))$

Symmetrization and its properties

$$F : S \rightarrow \mathbb{R}, \quad F = \sum_{\sigma \in C} f_\sigma \quad \text{symmetrization of } f : S \rightarrow \mathbb{R}$$

- F is symmetric wrt. the barycenter, i.e. $F(\sigma(x)) = F(x)$,
 $x \in S, \sigma \in C$

For $f : S \rightarrow \mathbb{R}$ and $\sigma \in C$ put $f_\sigma(x) = f(\sigma(x))$

Symmetrization and its properties

$$F : S \rightarrow \mathbb{R}, \quad F = \sum_{\sigma \in C} f_\sigma \quad \text{symmetrization of } f : S \rightarrow \mathbb{R}$$

- F is symmetric wrt. the barycenter, i.e. $F(\sigma(x)) = F(x)$,
 $x \in S$, $\sigma \in C$
- $F(v_i) = \sum_{k=0}^n f(v_k)$ and $F(\mathbf{b}) = (n+1)f(\mathbf{b})$

For $f : S \rightarrow \mathbb{R}$ and $\sigma \in C$ put $f_\sigma(x) = f(\sigma(x))$

Symmetrization and its properties

$$F : S \rightarrow \mathbb{R}, \quad F = \sum_{\sigma \in C} f_\sigma \quad \text{symmetrization of } f : S \rightarrow \mathbb{R}$$

- F is symmetric wrt. the barycenter, i.e. $F(\sigma(x)) = F(x)$, $x \in S$, $\sigma \in C$
- $F(v_i) = \sum_{k=0}^n f(v_k)$ and $F(\mathbf{b}) = (n+1)f(\mathbf{b})$
- $\int_S f_\sigma(x) \, dx = \int_S f(x) \, dx$, $\sigma \in C$

For $f : S \rightarrow \mathbb{R}$ and $\sigma \in C$ put $f_\sigma(x) = f(\sigma(x))$

Symmetrization and its properties

$$F : S \rightarrow \mathbb{R}, \quad F = \sum_{\sigma \in C} f_\sigma \quad \text{symmetrization of } f : S \rightarrow \mathbb{R}$$

- F is symmetric wrt. the barycenter, i.e. $F(\sigma(x)) = F(x)$, $x \in S$, $\sigma \in C$
- $F(v_i) = \sum_{k=0}^n f(v_k)$ and $F(\mathbf{b}) = (n+1)f(\mathbf{b})$
- $\int_S f_\sigma(x) \, dx = \int_S f(x) \, dx$, $\sigma \in C$
- $\int_S F(x) \, dx = (n+1) \int_S f(x) \, dx$

Symmetrization of a convex function

Assume that $f : S \rightarrow \mathbb{R}$ is convex.

Symmetrization of a convex function

Assume that $f : S \rightarrow \mathbb{R}$ is convex.

- f_σ is convex, $\sigma \in C$

Symmetrization of a convex function

Assume that $f : S \rightarrow \mathbb{R}$ is convex.

- f_σ is convex, $\sigma \in C$
- The symmetrization $F = \sum_{\sigma \in C} f_\sigma$ is convex

Hermite–Hadamard inequality

$S \subset \mathbb{R}^n$ – a simplex with vertices v_0, \dots, v_n and barycenter \mathbf{b} ,
 $f : S \rightarrow \mathbb{R}$ – a convex function

$$f(\mathbf{b}) \leq \frac{1}{\text{vol}(S)} \int_S f(\mathbf{x}) \, d\mathbf{x} \leq \frac{f(v_0) + \dots + f(v_n)}{n+1}$$

Hermite–Hadamard inequality

$S \subset \mathbb{R}^n$ – a simplex with vertices v_0, \dots, v_n and barycenter \mathbf{b} ,
 $f : S \rightarrow \mathbb{R}$ – a convex function

$$f(\mathbf{b}) \leq \frac{1}{\text{vol}(S)} \int_S f(\mathbf{x}) \, d\mathbf{x} \leq \frac{f(v_0) + \dots + f(v_n)}{n+1}$$

The inequality between LHH and RHH terms

Hermite–Hadamard inequality

$S \subset \mathbb{R}^n$ – a simplex with vertices v_0, \dots, v_n and barycenter \mathbf{b} ,
 $f : S \rightarrow \mathbb{R}$ – a convex function

$$f(\mathbf{b}) \leq \frac{1}{\text{vol}(S)} \int_S f(\mathbf{x}) \, d\mathbf{x} \leq \frac{f(v_0) + \dots + f(v_n)}{n+1}$$

The inequality between LHH and RHH terms

$$\bullet \quad \frac{1}{\text{vol}(S)} \int_S f(\mathbf{x}) \, d\mathbf{x} - f(\mathbf{b}) \qquad \text{LHH}$$

Hermite–Hadamard inequality

$S \subset \mathbb{R}^n$ – a simplex with vertices v_0, \dots, v_n and barycenter \mathbf{b} ,
 $f : S \rightarrow \mathbb{R}$ – a convex function

$$f(\mathbf{b}) \leq \frac{1}{\text{vol}(S)} \int_S f(\mathbf{x}) \, d\mathbf{x} \leq \frac{f(v_0) + \dots + f(v_n)}{n+1}$$

The inequality between LHH and RHH terms

- $\frac{1}{\text{vol}(S)} \int_S f(\mathbf{x}) \, d\mathbf{x} - f(\mathbf{b})$ LHH
- $\frac{f(v_0) + \dots + f(v_n)}{n+1} - \frac{1}{\text{vol}(S)} \int_S f(\mathbf{x}) \, d\mathbf{x}$ RHH

Hermite–Hadamard inequality

$S \subset \mathbb{R}^n$ – a simplex with vertices v_0, \dots, v_n and barycenter \mathbf{b} ,
 $f : S \rightarrow \mathbb{R}$ – a convex function

$$f(\mathbf{b}) \leq \frac{1}{\text{vol}(S)} \int_S f(\mathbf{x}) \, d\mathbf{x} \leq \frac{f(v_0) + \dots + f(v_n)}{n+1}$$

The inequality between LHH and RHH terms

- $\frac{1}{\text{vol}(S)} \int_S f(\mathbf{x}) \, d\mathbf{x} - f(\mathbf{b})$ LHH
- $\frac{f(v_0) + \dots + f(v_n)}{n+1} - \frac{1}{\text{vol}(S)} \int_S f(\mathbf{x}) \, d\mathbf{x}$ RHH
- $f : S \rightarrow \mathbb{R}$ convex $\implies 0 \leq \text{LHH} \leq n \text{ RHH}$



A. Witkowski, Sz. Wąsowicz, *On some inequality of Hermite–Hadamard type*, Opuscula Math. 32 (2012), 591–600.

Thank you very much
for your kind attention