

# Resolvability properties of similar topologies

Sebastian Lindner

University of Łódź

# Similarity

## Definition

Let  $\mathcal{F}, \mathcal{G} \subset 2^X \setminus \{\emptyset\}$ . We say, that  $\mathcal{F}$  is **coinitial** to  $\mathcal{G}$  if for every  $G \in \mathcal{G}$  there exists  $F \in \mathcal{F}$  such that  $F \subset G$ .

# Similarity

## Definition

Let  $\mathcal{F}, \mathcal{G} \subset 2^X \setminus \{\emptyset\}$ . We say, that  $\mathcal{F}$  is **coinitial** to  $\mathcal{G}$  if for every  $G \in \mathcal{G}$  there exists  $F \in \mathcal{F}$  such that  $F \subset G$ .

## Definition

Let us consider a non-empty space  $X$  with two topologies,  $\mathcal{T}_1$  and  $\mathcal{T}_2$ . We will say that  $\mathcal{T}_1$  and  $\mathcal{T}_2$  are **similar** ( $\mathcal{T}_2 \simeq_s \mathcal{T}_1$ ) iff for every  $A \subset X$  we have

$$\text{Int}_{\mathcal{T}_1}(A) \neq \emptyset \iff \text{Int}_{\mathcal{T}_2}(A) \neq \emptyset.$$

# Similarity

The relation of similarity was investigated among others in [BFKT]. In that note one can find the following useful characterization:

**Theorem, [Bartoszewicz, Filipczak, Kowalski, Terepeta 2014]**

*The following statements are equivalent*

- $\mathcal{T}_1 \simeq_s \mathcal{T}_2$
- $\mathcal{D}(X, \mathcal{T}_1) = \mathcal{D}(X, \mathcal{T}_2)$
- $(\mathcal{NB}(X, \mathcal{T}_1), \mathcal{ND}(X, \mathcal{T}_1)) = (\mathcal{NB}(X, \mathcal{T}_2), \mathcal{ND}(X, \mathcal{T}_2))$ .
- $\mathcal{T}_1 \setminus \{\emptyset\}$  and  $\mathcal{T}_2 \setminus \{\emptyset\}$  are mutually coinital.

# Similarity

The relation of similarity was investigated among others in [BFKT]. In that note one can find the following useful characterization:

**Theorem, [Bartoszewicz, Filipczak, Kowalski, Terepeta 2014]**

*The following statements are equivalent*

- $\mathcal{T}_1 \simeq_s \mathcal{T}_2$
- $\mathcal{D}(X, \mathcal{T}_1) = \mathcal{D}(X, \mathcal{T}_2)$
- $(\mathcal{NB}(X, \mathcal{T}_1), \mathcal{ND}(X, \mathcal{T}_1)) = (\mathcal{NB}(X, \mathcal{T}_2), \mathcal{ND}(X, \mathcal{T}_2))$ .
- $\mathcal{T}_1 \setminus \{\emptyset\}$  and  $\mathcal{T}_2 \setminus \{\emptyset\}$  are mutually coinital.

Moreover, for every topological space  $(X, \mathcal{T})$  the family  $\mathcal{NB}(X, \mathcal{T})$  forms an algebra of sets, and  $\mathcal{ND}(X, \mathcal{T})$  is an ideal contained in  $\mathcal{NB}(X, \mathcal{T})$ .

# Similarity

Let  $\Phi$  be an arbitrary property of topological spaces.

## Definition

We will say that  $\Phi$  is  $\simeq_s$ -**proof** iff for every topology  $\mathcal{T}_1$  having the property  $\Phi$  every similar topology  $\mathcal{T}_2$  also has the property  $\Phi$ .

# Resolvability

## Definition

Let  $(X, \mathcal{T})$  - an arbitrary topological space.

For any cardinal  $\kappa$  we say that  $(X, \mathcal{T})$  is  $\kappa$ -resolvable iff there exists a family of cardinality  $\kappa$  of pairwise disjoint dense subsets of  $X$ .

# Resolvability

## Definition

Let  $(X, \mathcal{T})$  - an arbitrary topological space.

For any cardinal  $\kappa$  we say that  $(X, \mathcal{T})$  is  **$\kappa$ -resolvable** iff there exists a family of cardinality  $\kappa$  of pairwise disjoint dense subsets of  $X$ .

The dense-in-itself space is called **maximally resolvable** if it is  $\Delta(X, \mathcal{T})$ -resolvable, where  $\Delta(X, \mathcal{T}) = \min\{|G| : G \in \mathcal{T}, G \neq \emptyset\}$ .



# Resolvability

## Definition

Let  $(X, \mathcal{T})$  - an arbitrary topological space.

For any cardinal  $\kappa$  we say that  $(X, \mathcal{T})$  is  **$\kappa$ -resolvable** iff there exists a family of cardinality  $\kappa$  of pairwise disjoint dense subsets of  $X$ .

The dense-in-itself space is called **maximally resolvable** if it is  $\Delta(X, \mathcal{T})$ -resolvable, where  $\Delta(X, \mathcal{T}) = \min\{|G| : G \in \mathcal{T}, G \neq \emptyset\}$ .

The space  $(X, \mathcal{T})$  is called **extra-resolvable**, if there exists the family  $\mathbf{D}$  of dense subsets of  $X$  such that  $|\mathbf{D}| > \Delta(X, \mathcal{T})$  and for every  $A, B \in \mathbf{D}$  we have  $A = B$  or  $A \cap B \in \mathcal{ND}(X, \mathcal{T})$ .

# Resolvability

## Definition

2-resolvable space is called simply **resolvable**. If the space is not resolvable, it is called **irresolvable**.

# Resolvability

## Definition

2-resolvable space is called simply **resolvable**. If the space is not resolvable, it is called **irresolvable**.

The space is called **open hereditary irresolvable** (OHI) if none of its open subspaces is resolvable.

# Resolvability

## Definition

2-resolvable space is called simply **resolvable**. If the space is not resolvable, it is called **irresolvable**.

The space is called **open hereditary irresolvable** (OHI) if none of its open subspaces is resolvable.

The space is called **hereditary irresolvable** (HI) if none of its subspaces is resolvable.

# Resolvability

## Definition

2-resolvable space is called simply **resolvable**. If the space is not resolvable, it is called **irresolvable**.

The space is called **open hereditary irresolvable** (OHI) if none of its open subspaces is resolvable.

The space is called **hereditary irresolvable** (HI) if none of its subspaces is resolvable.

The space is called **submaximal** if every dense set is open.

# Resolvability

## Definition

2-resolvable space is called simply **resolvable**. If the space is not resolvable, it is called **irresolvable**.

The space is called **open hereditary irresolvable** (OHI) if none of its open subspaces is resolvable.

The space is called **hereditary irresolvable** (HI) if none of its subspaces is resolvable.

The space is called **submaximal** if every dense set is open.

Theorem, [Rose, Sizemore, Thurston 2006]

$$\textit{submaximal} \implies \textit{HI} \implies \textit{OHI}$$

# Observation

The following properties and parameters are  $\simeq_s$ -proof:

- $\kappa$ -resolvability
- $\Delta(X, \mathcal{T})$
- maximal resolvability
- extra resolvability
- being dense-in-itself

# Hewitt's decomposition

## Theorem, [Hewitt 1943]

*Every topological space  $(X, \mathcal{T})$  can be uniquely represented as a disjoint union*

$$X = F_X \cup G_X,$$

*where  $F_X$  is closed and resolvable and  $G_X$  is open and hereditary irresolvable. This pair of sets is called the Hewitt decomposition of the space  $X$ .*



# Results

## Definition

Let  $\mathcal{A}$  be an arbitrary algebra. By  $\mathcal{H}(\mathcal{A})$  we shall denote the maximal hereditary subfamily of  $\mathcal{A}$  :

$$\mathcal{H}(\mathcal{A}) = \{A \in \mathcal{A} : \forall B \subset A B \in \mathcal{A}\}.$$

## Theorem

*Let  $(X, \mathcal{T})$  - an arbitrary topological space. Let  $(F_X, G_X)$  form the Hewitt decomposition of the space  $X$ . Then*

$$\mathcal{H}(\mathcal{NB}) = \{A \cup N : A \subset G_X \wedge N \in \mathcal{ND}\}.$$

# Results

## Corollary

$(X, \mathcal{T})$  is *resolvable* iff  $\mathcal{H}(\mathcal{NB}) = \mathcal{ND}$ .

Half of this equivalence can be found in [BFKT].

## Theorem

*The following statements are equivalent:*

- $(X, \mathcal{T})$  is OHI
- $(\mathcal{H}(\mathcal{NB}) \setminus \mathcal{ND})$  is coinital to  $(\mathcal{NB} \setminus \mathcal{ND})$ .
- $\mathcal{NB} = 2^X$

## Theorem

*The following statements are equivalent:*

- $(X, \mathcal{T})$  is OHI
- $(\mathcal{H}(\mathcal{NB}) \setminus \mathcal{ND})$  is coinital to  $(\mathcal{NB} \setminus \mathcal{ND})$ .
- $\mathcal{NB} = 2^X$

## Corollary

*OHI is  $\simeq_s$ -proof.*

# Submaximality and HI

Theorem, [Crossley 1974]

*For every topological space  $(X, \mathcal{T})$  the family*

$$\mathcal{F}(\mathcal{T}) = \{G \setminus N : G \in \mathcal{T}, N \in \mathcal{ND}\}$$

*forms a topology, similar to  $\mathcal{T}$  and NODEC.*

Theorem, [compare : Arhangel'ski, Collins 1995]

*submaximal = OHI + NODEC.*

# Submaximality and HI

Theorem, [Crossley 1974]

*For every topological space  $(X, \mathcal{T})$  the family*

$$\mathcal{F}(\mathcal{T}) = \{G \setminus N : G \in \mathcal{T}, N \in \mathcal{ND}\}$$

*forms a topology, similar to  $\mathcal{T}$  and NODEC.*

Theorem, [compare : Arhangel'ski, Collins 1995]

*submaximal = OHI + NODEC.*

Corollary

*For every OHI topology  $\mathcal{T}$  there exists the submaximal (hence : HI) similar topology  $\mathcal{F}(\mathcal{T})$ .*

# Submaximality and HI

Theorem, [Crossley 1974]

*For every topological space  $(X, \mathcal{T})$  the family*

$$\mathcal{F}(\mathcal{T}) = \{G \setminus N : G \in \mathcal{T}, N \in \mathcal{ND}\}$$

*forms a topology, similar to  $\mathcal{T}$  and NODEC.*

Theorem, [compare : Arhangel'ski, Collins 1995]

*submaximal = OHI + NODEC.*

Corollary

*For every OHI topology  $\mathcal{T}$  there exists the submaximal (hence : HI) similar topology  $\mathcal{F}(\mathcal{T})$ .*

Corollary

*HI and submaximality are not  $\simeq_s$ -proof.*

# Similarity is not hereditary

## Corollary

*The similarity  $(X, \mathcal{T}_1) \simeq_s (X, \mathcal{T}_2)$  does not necessarily imply the relation  $(A, \mathcal{T}_1|_A) \simeq_s (A, \mathcal{T}_2|_A)$ .*

## Theorem

*Let  $(X, \mathcal{T}_1) \simeq_s (X, \mathcal{T}_2)$ .*

*If  $A \in \mathcal{SO}(\mathcal{T}_1) \cap \mathcal{SO}(\mathcal{T}_2)$  then  $(A, \mathcal{T}_1|_A) \simeq_s (A, \mathcal{T}_2|_A)$ .*



# When the pair $(\mathcal{A}, \mathcal{I})$ is topological?

Let  $\mathcal{A} \subset 2^X$  be an arbitrary algebra and  $\mathcal{I} \subset \mathcal{A}$  - an arbitrary ideal of sets.

## Definition

We say that the pair  $(\mathcal{A}, \mathcal{I})$  is:

- HULL if for every  $A \subset X$  there exists  $H \in \mathcal{A}$  such that  $A \subset H$  and for any  $P \subset H \setminus A$ ,  $P \in \mathcal{A}$  we have  $P \in \mathcal{I}$ .

# When the pair $(\mathcal{A}, \mathcal{I})$ is topological?

Let  $\mathcal{A} \subset 2^X$  be an arbitrary algebra and  $\mathcal{I} \subset \mathcal{A}$  - an arbitrary ideal of sets.

## Definition

We say that the pair  $(\mathcal{A}, \mathcal{I})$  is:

- HULL if for every  $A \subset X$  there exists  $H \in \mathcal{A}$  such that  $A \subset H$  and for any  $P \subset H \setminus A$ ,  $P \in \mathcal{A}$  we have  $P \in \mathcal{I}$ .
- LDO if there exists lower density operator  $\Phi$  on  $(\mathcal{A}, \mathcal{I})$ .

# When the pair $(\mathcal{A}, \mathcal{I})$ is topological?

Let  $\mathcal{A} \subset 2^X$  be an arbitrary algebra and  $\mathcal{I} \subset \mathcal{A}$  - an arbitrary ideal of sets.

## Definition

We say that the pair  $(\mathcal{A}, \mathcal{I})$  is:

- HULL if for every  $A \subset X$  there exists  $H \in \mathcal{A}$  such that  $A \subset H$  and for any  $P \subset H \setminus A$ ,  $P \in \mathcal{A}$  we have  $P \in \mathcal{I}$ .
- LDO if there exists lower density operator  $\Phi$  on  $(\mathcal{A}, \mathcal{I})$ .
- LDO+ if there exists l.d.o.  $\Phi$  such that

$$\forall B \subset \mathcal{A} \quad \bigcup_{B \in \mathcal{B}} (B \cap \Phi(B)) \in \mathcal{A}.$$

# When the pair $(\mathcal{A}, \mathcal{I})$ is topological?

Let  $\mathcal{A} \subset 2^X$  be an arbitrary algebra and  $\mathcal{I} \subset \mathcal{A}$  - an arbitrary ideal of sets.

## Definition

We say that the pair  $(\mathcal{A}, \mathcal{I})$  is:

- HULL if for every  $A \subset X$  there exists  $H \in \mathcal{A}$  such that  $A \subset H$  and for any  $P \subset H \setminus A$ ,  $P \in \mathcal{A}$  we have  $P \in \mathcal{I}$ .
- LDO if there exists lower density operator  $\Phi$  on  $(\mathcal{A}, \mathcal{I})$ .
- LDO+ if there exists l.d.o.  $\Phi$  such that

$$\forall B \subset \mathcal{A} \quad \bigcup_{B \in \mathcal{B}} (B \cap \Phi(B)) \in \mathcal{A}.$$

- TOP if there exists a topology  $\mathcal{T}$  on  $X$  such that  $(\mathcal{A}, \mathcal{I}) = (\mathcal{NB}, \mathcal{ND})$ .

When the pair  $(\mathcal{A}, \mathcal{I})$  is topological?

Theorem, [Hejduk]

$$LDO \wedge HULL \implies TOP \implies HULL$$

# When the pair $(\mathcal{A}, \mathcal{I})$ is topological?

Theorem, [Hejduk]

$$LDO \wedge HULL \implies TOP \implies HULL$$

Theorem, [Bartoszewicz, Filipczak, Kowalski, Terepeta 2014]

*Every topology is similar to an abstract density topology.*

# When the pair $(\mathcal{A}, \mathcal{I})$ is topological?

Theorem, [Hejduk]

$$LDO \wedge HULL \implies TOP \implies HULL$$

Theorem, [Bartoszewicz, Filipczak, Kowalski, Terepeta 2014]

*Every topology is similar to an abstract density topology.*

*Since many properties connected with resolvability are  $\simeq_s$ -proof, it's enough to investigate these properties for ADT's.*

# When the pair $(\mathcal{A}, \mathcal{I})$ is topological?

Theorem, [Hejduk]

$$LDO \wedge HULL \implies TOP \implies HULL$$

Theorem, [Bartoszewicz, Filipczak, Kowalski, Terepeta 2014]

*Every topology is similar to an abstract density topology.*

*Since many properties connected with resolvability are  $\simeq_s$ -proof, it's enough to investigate these properties for ADT's.*

Corollary

$$TOP \iff LDO \wedge HULL$$



# When the pair $(\mathcal{A}, \mathcal{I})$ is topological?

## Theorem, [Hejduk]

$$LDO \wedge HULL \implies TOP \implies HULL$$

## Theorem, [Bartoszewicz, Filipczak, Kowalski, Terepeta 2014]

*Every topology is similar to an abstract density topology.*

*Since many properties connected with resolvability are  $\simeq_s$ -proof, it's enough to investigate these properties for ADT's.*

## Corollary

$$TOP \iff LDO \wedge HULL$$

## Theorem

$$TOP \iff LDO+$$

# When the pair $(\mathcal{A}, \mathcal{I})$ is topological?

## Corollary

*Let  $\mathcal{I} \subset 2^X$  is an arbitrary ideal. The following statements are equivalent:*

- *$(2^X, \mathcal{I})$  is TOP.*
- *there exists an submaximal topology  $\mathcal{T}$  on  $X$  such that  $(\mathcal{A}, \mathcal{I}) = (\mathcal{NB}, \mathcal{ND})$ .*
- *$(2^X, \mathcal{I})$  is LDO.*

# When the pair $(\mathcal{A}, \mathcal{I})$ is topological?

## Corollary

*Let  $(\mathcal{A}, \mathcal{I})$  be such that  $(\mathcal{H}(\mathcal{A}) \setminus \mathcal{I})$  is coinitial to  $(\mathcal{A} \setminus \mathcal{I})$ . Then the following holds*

$$TOP \implies \mathcal{A} = 2^X.$$

# When the pair $(\mathcal{A}, \mathcal{I})$ is topological?

## Corollary

*Let  $(\mathcal{A}, \mathcal{I})$  be such that  $(\mathcal{H}(\mathcal{A}) \setminus \mathcal{I})$  is coinitial to  $(\mathcal{A} \setminus \mathcal{I})$ . Then the following holds*

$$TOP \implies \mathcal{A} = 2^X.$$

## Theorem, [Zindulka]

*Every analytic set  $B \subset \mathbb{R}$  contains a universal measure zero set  $E$  such that  $\dim_H E = \dim_H B$ .*

# When the pair $(\mathcal{A}, \mathcal{I})$ is topological?

## Corollary

*Let  $(\mathcal{A}, \mathcal{I})$  be such that  $(\mathcal{H}(\mathcal{A}) \setminus \mathcal{I})$  is coinital to  $(\mathcal{A} \setminus \mathcal{I})$ . Then the following holds*

$$TOP \implies \mathcal{A} = 2^X.$$

## Theorem, [Zindulka]

*Every analytic set  $B \subset \mathbb{R}$  contains a universal measure zero set  $E$  such that  $\dim_H E = \dim_H B$ .*

Let  $\mathcal{L}$  -  $\sigma$ -ideal of Lebesgue measurable sets. Then  $\mathcal{H}(\mathcal{L}) = \mathcal{N}$ . For  $\alpha \in [0, 1)$  let  $\mathcal{I}_\alpha = \{E \in \mathcal{L} : \dim_H(E) \leq \alpha\}$ .

## Corollary

*For  $\alpha \in [0, 1)$  the pair  $(\mathcal{L}, \mathcal{I}_\alpha)$  is not TOP.*

# Bibliography



[A] A. Arhangel'ski, P. J. Collins, *On submaximal spaces*, Topology Appl. 64 (1995), no. 3, 219-241



[BC] A. Bartoszewicz, K. Ciesielski, *MB-representations and topological algebras* RAE, 27(2) 2001, pp. 749 - 756.



[BFKT] A. Bartoszewicz, M. Filipczak, A. Kowalski, M. Terepeta, *Similarity between topologies*, CEJM 12(4), 2014, 603-610.



[BK] A. Bartoszewicz, P. Koszmider, *When an atomic and complete algebra of sets is a field of sets with nowhere dense boundary*, J. App. Anal., 15 (2009), 119-127



[C] S. Gene Crossley, *A note on semitopological classes*, Proc. of AMS, 43 (2) 1974.



[He] J. Hejduk, *One more difference between measure and category*, Tatra Mt. Math. Publ. 49 (2011), 915.



[Hw] E. Hewitt, *A problem of set-theoretic topology*, Duke Mathematical Journal 10 (1943), no. 2, 309 - 333.



[RST] D. Rose, K. Sizemore, B. Thurston, *Strongly irresolvable spaces*, Intern. Journal of Mathematics and Mathematical Sciences, vol. 2006.



[Z] Ondrej Zindulka, *Universal measure zero sets with full Hausdorff dimension*