

Cantor's intersection theorem for cone metric spaces and its applications to fixed point theory

Jakub Klima

Institute of Mathematics, Łódź University of Technology, Poland

Stará Lesná, September 2, 2014

Cone metric spaces

Definition.

Let E be a real Banach space and $K \subset E$. K is called a cone iff:

- (i) K is closed, nonempty and $K \neq \{0\}$;
- (ii) if $a, b \in \mathbb{R}$, $a, b \geq 0$, $x, y \in K$, then $ax + by \in K$;
- (iii) if $x \in K$ and $-x \in K$, then $x = 0$.

Cone metric spaces

Definition.

Let E be a real Banach space and $K \subset E$. K is called a cone iff:

- (i) K is closed, nonempty and $K \neq \{0\}$;
- (ii) if $a, b \in \mathbb{R}$, $a, b \geq 0$, $x, y \in K$, then $ax + by \in K$;
- (iii) if $x \in K$ and $-x \in K$, then $x = 0$.

Definition.

A cone metric space is a pair (X, d) , where X is a nonempty set and $d : X \times X \rightarrow E$ satisfies three well-known axioms of a metric with respect to the following partial ordering \leq in E : $x \leq y$ iff $y - x \in K$.

Cone metric spaces

Definition.

Let E be a real Banach space and $K \subset E$. K is called a cone iff:

- (i) K is closed, nonempty and $K \neq \{0\}$;
- (ii) if $a, b \in \mathbb{R}$, $a, b \geq 0$, $x, y \in K$, then $ax + by \in K$;
- (iii) if $x \in K$ and $-x \in K$, then $x = 0$.

Definition.

A cone metric space is a pair (X, d) , where X is a nonempty set and $d : X \times X \rightarrow E$ satisfies three well-known axioms of a metric with respect to the following partial ordering \leq in E : $x \leq y$ iff $y - x \in K$.

Lemma.

Let $E = \mathbb{R}^n$ and $K = \{x \in \mathbb{R}^n : x_i \geq 0\}$. A function $d : X \times X \rightarrow \mathbb{R}^n$ ($d = (d_1, \dots, d_n)$) is a cone metric iff $\{d_1, \dots, d_n\}$ is a separating family of pseudometrics.

Convergence

Convergence

In the sequel we consider cones with a nonempty interior and $a \ll b$ means $b - a \in \text{int } K$.

Convergence

In the sequel we consider cones with a nonempty interior and $a \ll b$ means $b - a \in \text{int } K$.

Definition.

We say that a sequence (x_n) of elements of X is convergent to $x \in X$ if for any $0 \ll c$, there exists $k \in \mathbb{N}$ such that $d(x_n, x) \ll c$ for all $n \geq k$, and x is called the limit of (x_n) . Similarly, we define Cauchy sequence. Then the completeness of (X, d) is understood in an analogous way as in the case of real-valued metric.

Convergence

In the sequel we consider cones with a nonempty interior and $a \ll b$ means $b - a \in \text{int } K$.

Definition.

We say that a sequence (x_n) of elements of X is convergent to $x \in X$ if for any $0 \ll c$, there exists $k \in \mathbb{N}$ such that $d(x_n, x) \ll c$ for all $n \geq k$, and x is called the limit of (x_n) . Similarly, we define Cauchy sequence. Then the completeness of (X, d) is understood in an analogous way as in the case of real-valued metric.

Definition.

We say that a set $A \subset X$ is closed if it contains all of its limit points.

Convergence

In the sequel we consider cones with a nonempty interior and $a \ll b$ means $b - a \in \text{int } K$.

Definition.

We say that a sequence (x_n) of elements of X is convergent to $x \in X$ if for any $0 \ll c$, there exists $k \in \mathbb{N}$ such that $d(x_n, x) \ll c$ for all $n \geq k$, and x is called the limit of (x_n) . Similarly, we define Cauchy sequence. Then the completeness of (X, d) is understood in an analogous way as in the case of real-valued metric.

Definition.

We say that a set $A \subset X$ is closed if it contains all of its limit points.

Definition.

$B(x_0, r) := \{y \in X : d(x_0, y) \leq r\}$, $x_0 \in X, r \in K$.

Convergence

In the sequel we consider cones with a nonempty interior and $a \ll b$ means $b - a \in \text{int } K$.

Definition.


We say that a sequence (x_n) of elements of X is convergent to $x \in X$ if for any $0 \ll c$, there exists $k \in \mathbb{N}$ such that $d(x_n, x) \ll c$ for all $n \geq k$, and x is called the limit of (x_n) . Similarly, we define Cauchy sequence. Then the completeness of (X, d) is understood in an analogous way as in the case of real-valued metric.

Definition.

We say that a set $A \subset X$ is closed if it contains all of its limit points.

Definition.

$B(x_0, r) := \{y \in X : d(x_0, y) \leq r\}$, $x_0 \in X, r \in K$.

It is easy to show that $B(x_0, r)$ is a closed set. 

Boundedness

Definition.

A nonempty set $A \subset E$ is called bounded from above if there exists $c \in E$ such that $x \leq c$, for all $x \in A$.

Boundedness

Definition.

A nonempty set $A \subset E$ is called bounded from above if there exists $c \in E$ such that $x \leq c$, for all $x \in A$.

Definition.

A cone K is called strongly minihedral if every set which is bounded from above has a supremum.

Boundedness

Definition.

A nonempty set $A \subset E$ is called bounded from above if there exists $c \in E$ such that $x \leq c$, for all $x \in A$.

Definition.

A cone K is called strongly minihedral if every set which is bounded from above has a supremum.

Definition.

A nonempty set $B \subset X$ is called bounded if there exists $c \in K$ such that $d(x, y) \leq c$, for all $x, y \in B$. A vector c is called a bound of B .

Boundedness

Definition.

A nonempty set $A \subset E$ is called bounded from above if there exists $c \in E$ such that $x \leq c$, for all $x \in A$.

Definition.

A cone K is called strongly minihedral if every set which is bounded from above has a supremum.

Definition.

A nonempty set $B \subset X$ is called bounded if there exists $c \in K$ such that $d(x, y) \leq c$, for all $x, y \in B$. A vector c is called a bound of B .

Definition.

Let B be a subset of X . The diameter of B is the vector $\delta(B) := \sup \{d(x, y) : x, y \in B\}$ (if it exists).

Cantor's intersection theorem

Cantor's intersection theorem

Theorem (S. H. Alnafei, S. Radenović, N. Shahzad; 2011).

A cone metric space (X, d) over a strongly minihedral cone K is complete iff every decreasing sequence (A_n) of nonempty closed sets with $\delta(A_n) \xrightarrow{\|\cdot\|} 0$, has nonempty intersection.

Cantor's intersection theorem

Theorem (S. H. Alnafei, S. Radenović, N. Shahzad; 2011).

A cone metric space (X, d) over a strongly minihedral cone K is complete iff every decreasing sequence (A_n) of nonempty closed sets with $\delta(A_n) \xrightarrow{\|\cdot\|} 0$, has nonempty intersection.

Theorem.

Let (X, d) be a cone metric space. The following statements are equivalent:

- (i) X is complete;
- (ii) every decreasing sequence (A_n) of nonempty closed sets such that every A_n has a bound $c_n \in K$ and $c_n \xrightarrow{\|\cdot\|} 0$, has a nonempty intersection;
- (iii) every decreasing sequence $((B(x_n, r_n)))$ of closed balls with $r_n \xrightarrow{\|\cdot\|} 0$, has a nonempty intersection.

Banach's fixed point theorem

Banach's fixed point theorem

Theorem.

Let K be a cone in a Banach space E and $\Lambda : E \rightarrow E$ be a linear bounded operator, which is monotone, i.e., $a \leq b$ implies $\Lambda a \leq \Lambda b$ for $a, b \in K$. Let (X, d) be a complete cone metric space and $T : X \rightarrow X$ be such that

$$d(Tx, Ty) \leq \Lambda(d(x, y)) \text{ for all } x, y \in X.$$

If $\lim_{n \rightarrow \infty} \|\Lambda^n\|^{\frac{1}{n}} < 1$, then T has a unique fixed point x_* and for any $x_0 \in X$, $x_* = \lim_{n \rightarrow \infty} T^n x_0$.

Remark 1.

We can prove Banach's fixed point theorem using Cantor's intersection theorem in a similar way as in real-valued metric spaces:

- (i) We consider sets $A_n := \{x \in X : d(x, Tx) \leq \frac{1}{n}c\}$.
- (ii) The sequence (A_n) satisfies all the assumptions of Cantor's intersection theorem (the proof that the sets A_n are closed is different from the one in real-valued metric case).
- (iii) There exists unique $x_* \in \bigcap_{n \in \mathbb{N}} A_n$ and $Tx_* = x_*$.

Remark 2.

It can happen that $x_n \rightarrow x$ and $y_n \rightarrow y$ but

$d(x_n, y_n) \not\xrightarrow{\|\cdot\|} d(x, y)$, so a cone metric d need not be a continuous function.

Remark 2.

It can happen that $x_n \rightarrow x$ and $y_n \rightarrow y$ but

$d(x_n, y_n) \not\rightarrow d(x, y)$, so a cone metric d need not be a continuous function.

Example. Let $E = C^{(1)}[0, 1]$ with the norm $\|x\| = \|x\|_\infty + \|x'\|_\infty$, $K = \{x \in E : x(t) \geq 0, \text{ for all } t \in [0, 1]\}$. We define a cone metric $d : K \times K \rightarrow K$,

$$d(x, y) := \begin{cases} 0, & x = y \\ x + y, & x \neq y \end{cases}.$$

Consider $x_n(t) := \frac{1 + \sin(nt)}{n}$, $x(t) = y_n(t) = y(t) = 0$, then

$x_n \rightarrow x, y_n \rightarrow y$, but $d(x_n, y_n) \not\rightarrow d(x, y)$.

- ① L.-G. Huang and X. Zhang, *Cone metric spaces and fixed point theorems of contractive mappings*, J. Math. Anal. Appl. **332** (2007), 1468–1476.
- ② S. H. Alnafei, S. Radenović, N. Shahzad, *Fixed point theorems for mappings with convex diminishing diameters on cone metric spaces*, Applied Mathematics Letters **24** (2011), 2162–2166.
- ③ S. Janković, Z. Kadelburg and S. Radenović, *On cone metric spaces: a survey*, Nonlinear Anal. **74** (2011), 2591–2601.