

Faculty of Mathematics and Computer Science
Łódź University

On some properties of functions connected with focal points of entropy

Ewa Korczak-Kubiak, Anna Loranty, Ryszard J. Pawlak

Stará Lesná, September 2014

Notations and assumptions

Notations and assumptions

(X, ρ) - compact metric space

Notations and assumptions

(X, ρ) - compact metric space

$O(x_0)$ -the family of all neighbourhoods of x_0

Notations and assumptions

(X, ρ) - compact metric space

$O(x_0)$ -the family of all neighbourhoods of x_0

$$B_\rho(x_0, \varepsilon) = \{x \in X : \rho(x_0, x) < \varepsilon\}$$

Notations and assumptions

(X, ρ) - compact metric space

$O(x_0)$ -the family of all neighbourhoods of x_0

$$B_\rho(x_0, \varepsilon) = \{x \in X : \rho(x_0, x) < \varepsilon\}$$

$$f : X \rightarrow X, f^0(x) = x, f^{n+1}(x) = f(f^n(x))$$

Notations and assumptions

(X, ρ) - compact metric space

$O(x_0)$ -the family of all neighbourhoods of x_0

$$B_\rho(x_0, \varepsilon) = \{x \in X : \rho(x_0, x) < \varepsilon\}$$

$$f : X \rightarrow X, f^0(x) = x, f^{n+1}(x) = f(f^n(x))$$

$$\text{Fix}(f) = \{x \in X : f(x) = x\}$$

Notations and assumptions

(X, ρ) - compact metric space

$O(x_0)$ -the family of all neighbourhoods of x_0

$$B_\rho(x_0, \varepsilon) = \{x \in X : \rho(x_0, x) < \varepsilon\}$$

$$f : X \rightarrow X, f^0(x) = x, f^{n+1}(x) = f(f^n(x))$$

$$\text{Fix}(f) = \{x \in X : f(x) = x\}$$

$$A \xrightarrow[f]{} B \Leftrightarrow B \subset f(A)$$

Notations and assumptions

(X, ρ) - compact metric space

$O(x_0)$ -the family of all neighbourhoods of x_0

$$B_\rho(x_0, \varepsilon) = \{x \in X : \rho(x_0, x) < \varepsilon\}$$

$$f : X \rightarrow X, f^0(x) = x, f^{n+1}(x) = f(f^n(x))$$

$$\text{Fix}(f) = \{x \in X : f(x) = x\}$$

$$A \xrightarrow[f]{} B \Leftrightarrow B \subset f(A)$$

$h(f)$ - entropy of a function f

Notations and assumptions

(X, ρ) - compact metric space

$O(x_0)$ - the family of all neighbourhoods of x_0

$$B_\rho(x_0, \varepsilon) = \{x \in X : \rho(x_0, x) < \varepsilon\}$$

$$f : X \rightarrow X, f^0(x) = x, f^{n+1}(x) = f(f^n(x))$$

$$\text{Fix}(f) = \{x \in X : f(x) = x\}$$

$$A \xrightarrow{f} B \Leftrightarrow B \subset f(A)$$

$h(f)$ - entropy of a function f

$\mathcal{F} \subset \exp(X) \setminus \emptyset$ - a family such that each nonempty open set contains some element of \mathcal{F} .

Y - open subset of X .

$\vartheta_{\mathcal{F}}^Y = \{(A_1, \dots, A_m) : m \in \mathbb{N}, A_1 \dots A_m \in \mathcal{F}, A_1, \dots, A_m \subset Y, \overline{A_i} \cap \overline{A_j} = \emptyset\}$.

For a sequence $A = (A_1, \dots, A_m) \in \vartheta_{\mathcal{F}}^X$ we define the matrix

$\mathcal{M}_{A,f} = [m_{ij}]_{i,j=1}^m$ in the following way:

$$m_{ij} = \begin{cases} 1 & \text{if } A_i \xrightarrow{f} A_j \\ 0 & \text{otherwise.} \end{cases}$$

The entropy of a function f with respect to a sequence

$A = (A_1, \dots, A_m) \in \vartheta_{\mathcal{F}}^X$:

$$H_f(F) = \begin{cases} \log \sigma(\mathcal{M}_{F,f}) & \text{if } \sigma(\mathcal{M}_{F,f}) > 0, \\ 0 & \text{if } \sigma(\mathcal{M}_{F,f}) = 0. \end{cases}$$

where $\sigma(\mathcal{M}_{F,f}) = \limsup_{n \rightarrow \infty} \sqrt[n]{\text{tr}(\mathcal{M}_{F,f}^n)}$.

$$H_{\mathcal{F},f}(Y) = \sup \left\{ \frac{1}{n} H_{f^n}(F) : F \in \vartheta_{\mathcal{F}}^Y \wedge n \in \mathbb{N} \right\}.$$

$$d(\mathcal{F}, f, Y) = \begin{cases} \frac{H_{\mathcal{F},f}(Y)}{h(f)} & \text{if } h(f) \in (0, \infty), \\ 1 & \text{if } H_{\mathcal{F},f}(Y) = \infty \text{ or } h(f) = 0, \\ 0 & \text{if } H_{\mathcal{F},f}(Y) \in [0, \infty) \text{ and } h(f) = \infty. \end{cases}$$

A density of entropy of f with respect to \mathcal{F} at the point x_0 :

$$E_{\mathcal{F},f}(x_0) = \inf \{ d(\mathcal{F}, f, V) : V \in O(x_0) \}$$

We say that a point $x_0 \in X$ is an **\mathcal{F} -focal point of entropy of f** if:

$$E_{\mathcal{F},f}(x_0) = 1.$$

$E_{\mathcal{F}}(f)$ - the set of all \mathcal{F} -focal points of entropy of f .

Nonwandering and recurrent points

Nonwandering and recurrent points

Let $f : X \rightarrow X$. We say that a point $x_0 \in X$ is a **wandering point** for f if there exists a set $U \in \mathcal{O}(x_0)$ such that $U \cap f^i(U) = \emptyset$ for any $i \in \mathbb{N}$.

Nonwandering and recurrent points

Let $f : X \rightarrow X$. We say that a point $x_0 \in X$ is a **wandering point** for f if there exists a set $U \in \mathcal{O}(x_0)$ such that $U \cap f^i(U) = \emptyset$ for any $i \in \mathbb{N}$.

A point which is not a wandering point for f is called **nonwandering point** for f .

Nonwandering and recurrent points

Let $f : X \rightarrow X$. We say that a point $x_0 \in X$ is a **wandering point** for f if there exists a set $U \in \mathcal{O}(x_0)$ such that $U \cap f^i(U) = \emptyset$ for any $i \in \mathbb{N}$.

A point which is not a wandering point for f is called **nonwandering point** for f .

$\Omega(f)$ - the set of all nonwandering points for f .

Nonwandering and recurrent points

Let $f : X \rightarrow X$. We say that a point $x_0 \in X$ is a **wandering point** for f if there exists a set $U \in \mathcal{O}(x_0)$ such that $U \cap f^i(U) = \emptyset$ for any $i \in \mathbb{N}$.

A point which is not a wandering point for f is called **nonwandering point** for f .

$\Omega(f)$ - the set of all nonwandering points for f .

A point $x_0 \in X$ is called a **recurrent point** for f if there exists a sequence $\{n_k\}_{k \in \mathbb{N}}$ of positive integers such that $x_0 = \lim_{k \rightarrow \infty} f^{n_k}(x_0)$.

Nonwandering and recurrent points

Let $f : X \rightarrow X$. We say that a point $x_0 \in X$ is a **wandering point** for f if there exists a set $U \in \mathcal{O}(x_0)$ such that $U \cap f^i(U) = \emptyset$ for any $i \in \mathbb{N}$.

A point which is not a wandering point for f is called **nonwandering point** for f .

$\Omega(f)$ - the set of all nonwandering points for f .

A point $x_0 \in X$ is called a **recurrent point** for f if there exists a sequence $\{n_k\}_{k \in \mathbb{N}}$ of positive integers such that $x_0 = \lim_{k \rightarrow \infty} f^{n_k}(x_0)$.

$R(f)$ - the set of all recurrent points for f .

Nonwandering and recurrent points

Let $f : X \rightarrow X$. We say that a point $x_0 \in X$ is a **wandering point** for f if there exists a set $U \in \mathcal{O}(x_0)$ such that $U \cap f^i(U) = \emptyset$ for any $i \in \mathbb{N}$.

A point which is not a wandering point for f is called **nonwandering point** for f .

$\Omega(f)$ - the set of all nonwandering points for f .

A point $x_0 \in X$ is called a **recurrent point** for f if there exists a sequence $\{n_k\}_{k \in \mathbb{N}}$ of positive integers such that $x_0 = \lim_{k \rightarrow \infty} f^{n_k}(x_0)$.

$R(f)$ - the set of all recurrent points for f .

$$R(f) \subset \Omega(f)$$

Theorem 1.

Let $f : X \rightarrow X$ be a function such that $h(f) > 0$ and $\mathcal{F} \subset \exp(X) \setminus \{\emptyset\}$ be a family such that each open set contains some element of \mathcal{F} . If $x \in X$ is an \mathcal{F} -focal point of entropy of f then x is a nonwandering point for f .

Theorem 1.

Let $f : X \rightarrow X$ be a function such that $h(f) > 0$ and $\mathcal{F} \subset \exp(X) \setminus \{\emptyset\}$ be a family such that each open set contains some element of \mathcal{F} . If $x \in X$ is an \mathcal{F} -focal point of entropy of f then x is a nonwandering point for f .

Theorem 1.

Let $f : X \rightarrow X$ be a function such that $h(f) > 0$ and $\mathcal{F} \subset \exp(X) \setminus \{\emptyset\}$ be a family such that each open set contains some element of \mathcal{F} . If $x \in X$ is an \mathcal{F} -focal point of entropy of f then x is a nonwandering point for f .

Example

$X = [0, 1]$, $f(x) = 1$. Obviously $h(f) = 0$.

Theorem 1.

Let $f : X \rightarrow X$ be a function such that $h(f) > 0$ and $\mathcal{F} \subset \exp(X) \setminus \{\emptyset\}$ be a family such that each open set contains some element of \mathcal{F} . If $x \in X$ is an \mathcal{F} -focal point of entropy of f then x is a nonwandering point for f .

Example

$X = [0, 1]$, $f(x) = 1$. Obviously $h(f) = 0$.

Each point $x_0 \in [0, 1)$ is \mathcal{F} -focal point of entropy of f , but it is a wandering point for f .

$$h(f) = 0 \Rightarrow E_{\mathcal{F}}(f) = X.$$

Theorem 1.

Let $f : X \rightarrow X$ be a function such that $h(f) > 0$ and $\mathcal{F} \subset \exp(X) \setminus \{\emptyset\}$ be a family such that each open set contains some element of \mathcal{F} . If $x \in X$ is an \mathcal{F} -focal point of entropy of f then x is a nonwandering point for f .

Example

$X = [0, 1]$, $f(x) = 1$. Obviously $h(f) = 0$.


Each point $x_0 \in [0, 1)$ is \mathcal{F} -focal point of entropy of f , but it is a wandering point for f .

Let $U \in \mathcal{O}(x_0)$ be such that $1 \notin U$. For any $i \in \mathbb{N}$ we have $U \cap f^i(U) = U \cap \{1\} = \emptyset$.

We will say that $f \in \mathfrak{J}(\mathcal{F})$ if for any $n \in \mathbb{N}$ and for any sequence $(A_1, \dots, A_{m-1}, A_1) \in \vartheta_{\mathcal{F}}^X$ such that

$$A_1 \xrightarrow{f^n} A_2 \xrightarrow{f^n} \dots \xrightarrow{f^n} A_{m-1} \xrightarrow{f^n} A_m = A_1$$


there exists $x_0 \in A_1$ such that $f^{n+m-1}(x_0) = x_0$ and $f^{n+i-1}(x_0) \in A_{i+1}$, for $i \in \{1, 2, \dots, m-1\}$.



We will say that $f \in \mathfrak{J}(\mathcal{F})$ if for any $n \in \mathbb{N}$ and for any sequence $(A_1, \dots, A_{m-1}, A_1) \in \vartheta_{\mathcal{F}}^X$ such that

$$A_1 \xrightarrow{f^n} A_2 \xrightarrow{f^n} \dots \xrightarrow{f^n} A_{m-1} \xrightarrow{f^n} A_m = A_1$$

there exists $x_0 \in A_1$ such that $f^{n+m-1}(x_0) = x_0$ and $f^{n+i-1}(x_0) \in A_{i+1}$, for $i \in \{1, 2, \dots, m-1\}$.



Theorem 2.


Let $f : X \rightarrow X$ be a function such that $h(f) > 0$ and $f \in J(\mathcal{F})$. If x_0 is \mathcal{F} -focal point of entropy of f , then $h(f \upharpoonright (U \cap R(f))) = h(f)$ for any $U \in O(x_0)$.

Theorem 2.

Let $f : X \rightarrow X$ be a function such that $h(f) > 0$ and $f \in J(\mathcal{F})$. If x_0 is \mathcal{F} -focal point of entropy of f , then $h(f \upharpoonright (U \cap R(f))) = h(f)$ for any $U \in O(x_0)$.


Theorem 2.

Let $f : X \rightarrow X$ be a function such that $h(f) > 0$ and $f \in J(\mathcal{F})$. If x_0 is \mathcal{F} -focal point of entropy of f , then $h(f \upharpoonright (U \cap R(f))) = h(f)$ for any $U \in O(x_0)$.



Theorem 2.

Let $f : X \rightarrow X$ be a function such that $h(f) > 0$ and $f \in J(\mathcal{F})$. If x_0 is \mathcal{F} -focal point of entropy of f , then $h(f \upharpoonright (U \cap R(f))) = h(f)$ for any $U \in O(x_0)$.



$$E_{\mathcal{F}}(f) = X \text{ but } R(f) = \{0, 1\}.$$

\mathcal{F} -modifiable function

\mathcal{F} -modifiable function

(X, ρ) - compact metric space being an m -dimensional manifold with boundary

\mathcal{F} -modifiable function

(X, ρ) - compact metric space being an m -dimensional manifold with boundary

A topological space X is called an m -dimensional manifold with boundary if it is a second countable Hausdorff space and every point $x \in X$ has a neighbourhood that is homeomorphic to an open subset of $\mathbb{H}^m = \{(x_1, \dots, x_m) : x_m \geq 0\}$.

\mathcal{F} -modifiable function

(X, ρ) - compact metric space being an m -dimensional manifold with boundary

$$f, g : X \rightarrow X$$

\mathcal{F} -modifiable function

(X, ρ) - compact metric space being an m -dimensional manifold with boundary

$$f, g : X \rightarrow X$$

$$\nabla(f, g) = \{x \in X : f(x) \neq g(x)\}$$

\mathcal{F} -modifiable function

(X, ρ) - compact metric space being an m -dimensional manifold with boundary

$$f, g : X \rightarrow X$$

$$\nabla(f, g) = \{x \in X : f(x) \neq g(x)\}$$

ρ_u - metric of uniform convergence

\mathcal{F} -modifiable function

(X, ρ) - compact metric space being an m -dimensional manifold with boundary

$$f, g : X \rightarrow X$$

$$\nabla(f, g) = \{x \in X : f(x) \neq g(x)\}$$

ρ_u - metric of uniform convergence

We say that a continuous function $f : X \rightarrow X$ is **\mathcal{F} -modifiable at a point x_0** if

for any $\varepsilon > 0$ there exists a continuous function g such that x_0 is an \mathcal{F} -focal entropy point of g , $\rho_u(f, g) < \varepsilon$ and $\nabla(f, g) \subset B_\rho(x_0, \varepsilon)$.

\mathcal{F} -modifiable function

(X, ρ) - compact metric space being an m -dimensional manifold with boundary

$$f, g : X \rightarrow X$$

$$\nabla(f, g) = \{x \in X : f(x) \neq g(x)\}$$

ρ_u - metric of uniform convergence

We say that a continuous function $f : X \rightarrow X$ is **\mathcal{F} -modifiable at a point x_0** if

for any $\varepsilon > 0$ there exists a continuous function g such that x_0 is an \mathcal{F} -focal entropy point of g , $\rho_u(f, g) < \varepsilon$ and $\nabla(f, g) \subset B_\rho(x_0, \varepsilon)$.

Theorem 4.

Let X be a compact metric space being an m -dimensional manifold, $f : X \rightarrow X$ be a continuous function, $x_0 \in \text{Fix}(f)$ and \mathcal{F} be a family of all closed subsets of X . Then f is \mathcal{F} -modifiable at the point x_0 .

- [1] L. Alsedá, J. Llibre, M. Misiurewicz, *Combinatorial Dynamics and Entropy in Dimension One*, World Sci., 1993.
- [2] L. S. Block, W. A. Coppel *Dynamics in one dimension*, Springer-Verlag, Berlin Heidelberg, 1992.
- [3] M. Čiklová *Dynamical systems generated by functions with connected \mathcal{G}_δ graphs*, Real Anal. Exchange, vol. 30(2), 2004/2005, 617–638.
- [4] M. Misiurewicz *Horseshoes for mappings of the interval*, Bull.Acad. Polon. Sci. Ser. Sci. Math. 27 (1979), 167–169.
- [5] A. Loranty, R. J. Pawlak *The generalized entropy in the generalized topological spaces*, Topol. Appl. 159, 2012, 1734–1742.
- [6] R. J. Pawlak *On the entropy of Darboux functions*, Colloq. Math., 116(2), 2009, 227–241.