

# Rearranging series of vectors on a small set

Paweł Klinga

University of Gdańsk

September 4, 2014

## Theorem (Riemann)

*For any conditionally convergent series of reals  $\sum_{n=1}^{\infty} a_n$  and any  $a \in \mathbb{R}$  there exists a permutation  $\sigma : \mathbb{N} \rightarrow \mathbb{N}$  such that  $\sum_{n=1}^{\infty} a_{\sigma(n)} = a$ .*

## Theorem (Riemann)

*For any conditionally convergent series of reals  $\sum_{n=1}^{\infty} a_n$  and any  $a \in \mathbb{R}$  there exists a permutation  $\sigma : \mathbb{N} \rightarrow \mathbb{N}$  such that  $\sum_{n=1}^{\infty} a_{\sigma(n)} = a$ .*

## Theorem (Wilczyński)

*For any conditionally convergent series of reals  $\sum_{n=1}^{\infty} a_n$  and any  $a \in \mathbb{R}$  there exists a permutation  $\sigma : \mathbb{N} \rightarrow \mathbb{N}$  such that  $\sum_{n=1}^{\infty} a_{\sigma(n)} = a$  and  $\text{supp}(\sigma) = \{n \in \mathbb{N} : \sigma(n) \neq n\} \in \mathcal{I}_d$ , where*

$$\mathcal{I}_d = \{A \subseteq \mathbb{N} : \limsup_{n \rightarrow \infty} \frac{|A \cap \{0, 1, \dots, n-1\}|}{n} = 0\}.$$

## Definition

*We say that an ideal  $\mathcal{I} \subseteq \mathcal{P}(\mathbb{N})$  has the (R) property if for any conditionally convergent series of reals  $\sum_{n=1}^{\infty} a_n$  and any  $a \in \mathbb{R}$  there exists a permutation  $\sigma : \mathbb{N} \rightarrow \mathbb{N}$  such that  $\sum_{n=1}^{\infty} a_{\sigma(n)} = a$  and  $\text{supp}(\sigma) \in \mathcal{I}$ .*

## Definition

*We say that an ideal  $\mathcal{I} \subseteq \mathcal{P}(\mathbb{N})$  has the (R) property if for any conditionally convergent series of reals  $\sum_{n=1}^{\infty} a_n$  and any  $a \in \mathbb{R}$  there exists a permutation  $\sigma : \mathbb{N} \rightarrow \mathbb{N}$  such that  $\sum_{n=1}^{\infty} a_{\sigma(n)} = a$  and  $\text{supp}(\sigma) \in \mathcal{I}$ .*

## Theorem (Filipów, Szuca)

*Let  $\mathcal{I} \subseteq \mathcal{P}(\mathbb{N})$  be an ideal. The following are equivalent.*

- (i)  $\mathcal{I}$  has the (R) property.*
- (ii)  $\mathcal{I}$  cannot be extended to a summable ideal.*

# The Lévy-Steinitz Theorem

## Definition

Let  $(v_n)_n$  be a sequence of vectors in  $\mathbb{R}^m$ .

$$S\left(\sum_{n=1}^{\infty} v_n\right) = \left\{v \in \mathbb{R}^m : \exists \sigma : \mathbb{N} \rightarrow \mathbb{N} \text{ - permutation} \right. \\ \left. \sum_{n=1}^{\infty} v_{\sigma(n)} = v \right\}.$$

# The Lévy-Steinitz Theorem

## Definition

Let  $(v_n)_n$  be a sequence of vectors in  $\mathbb{R}^m$ .

$$S(\sum_{n=1}^{\infty} v_n) = \{v \in \mathbb{R}^m : \exists \sigma : \mathbb{N} \rightarrow \mathbb{N} \text{ - permutation} \\ \sum_{n=1}^{\infty} v_{\sigma(n)} = v\}.$$

## Definition

Let  $(v_n)_n$  be a sequence of vectors in  $\mathbb{R}^m$ .

$$S_{\mathcal{I}}(\sum_{n=1}^{\infty} v_n) = \{v \in \mathbb{R}^m : \exists \sigma : \mathbb{N} \rightarrow \mathbb{N} \text{ - permutation} \\ \sum_{n=1}^{\infty} v_{\sigma(n)} = v \text{ and } \text{supp}(\sigma) \in \mathcal{I}\}.$$

# The Lévy-Steinitz Theorem

## Theorem (Lévy, Steinitz)

*Let  $(v_n)_n$  be a sequence of vectors in  $R^m$ . The set  $S(\sum_{n=1}^{\infty} v_n)$  is either empty or is of the form  $s_0 + L$  for some vector  $s_0$  and some linear subspace  $L$ .*



# The Lévy-Steinitz Theorem

## Theorem (Lévy, Steinitz)

*Let  $(v_n)_n$  be a sequence of vectors in  $R^m$ . The set  $S(\sum_{n=1}^{\infty} v_n)$  is either empty or is of the form  $s_0 + L$  for some vector  $s_0$  and some linear subspace  $L$ .*

## Examples

$$S\left(\sum_{n=1}^{\infty} \left(\frac{(-1)^n}{n}, \frac{(-1)^n}{n}\right)\right) = \{(x, y) : x = y\},$$

# The Lévy-Steinitz Theorem

## Theorem (Lévy, Steinitz)

*Let  $(v_n)_n$  be a sequence of vectors in  $\mathbb{R}^m$ . The set  $S(\sum_{n=1}^{\infty} v_n)$  is either empty or is of the form  $s_0 + L$  for some vector  $s_0$  and some linear subspace  $L$ .*

## Examples

$$S\left(\sum_{n=1}^{\infty} \left(\frac{(-1)^n}{n}, \frac{(-1)^n}{n}\right)\right) = \{(x, y) : x = y\},$$

$$S\left(\sum_{n=1}^{\infty} \left(\frac{(-1)^n}{n}, \frac{(-1)^n}{\sqrt{n}}\right)\right) = \mathbb{R}^2.$$

# The Lévy-Steinitz Theorem

The form of the space  $s_0 + L$  in the Lévy-Steinitz theorem can be expressed in a more precise way.

# The Lévy-Steinitz Theorem

The form of the space  $s_0 + L$  in the Lévy-Steinitz theorem can be expressed in a more precise way.

Let  $F = \{w \in \mathbb{R}^m : \sum_{n=1}^{\infty} (w \circ v_n)^+ < \infty\}$ , where  $\circ$  denotes the real inner product and  $a^+ = \max\{a, 0\}$ .

# The Lévy-Steinitz Theorem

The form of the space  $s_0 + L$  in the Lévy-Steinitz theorem can be expressed in a more precise way.

Let  $F = \{w \in \mathbb{R}^m : \sum_{n=1}^{\infty} (w \circ v_n)^+ < \infty\}$ , where  $\circ$  denotes the real inner product and  $a^+ = \max\{a, 0\}$ .

Let  $F^\perp = \{v \in \mathbb{R}^m : \forall w \in F \ v \circ w = 0\}$ .

# The Lévy-Steinitz Theorem

The form of the space  $s_0 + L$  in the Lévy-Steinitz theorem can be expressed in a more precise way.

Let  $F = \{w \in \mathbb{R}^m : \sum_{n=1}^{\infty} (w \circ v_n)^+ < \infty\}$ , where  $\circ$  denotes the real inner product and  $a^+ = \max\{a, 0\}$ .

Let  $F^\perp = \{v \in \mathbb{R}^m : \forall w \in F \ v \circ w = 0\}$ .

Also let  $s_0$  be any sum (rearranged or not) of the series.

# The Lévy-Steinitz Theorem

The form of the space  $s_0 + L$  in the Lévy-Steinitz theorem can be expressed in a more precise way.

Let  $F = \{w \in \mathbb{R}^m : \sum_{n=1}^{\infty} (w \circ v_n)^+ < \infty\}$ , where  $\circ$  denotes the real inner product and  $a^+ = \max\{a, 0\}$ .

Let  $F^\perp = \{v \in \mathbb{R}^m : \forall w \in F \ v \circ w = 0\}$ .

Also let  $s_0$  be any sum (rearranged or not) of the series.

Finally, if  $S(\sum_{n=1}^{\infty} v_n)$  is not empty, then

$$S\left(\sum_{n=1}^{\infty} v_n\right) = s_0 + F^\perp.$$

# The two-dimensional case, ideal version

## Theorem

*Let  $l \subseteq \mathbb{R}^2$  be such line on the plane and  $\sum_{n=1}^{\infty} v_n$  such series in  $\mathbb{R}^2$  that*

$$S \left( \sum_{n=1}^{\infty} v_n \right) = l.$$

*Then*

$$S_{\mathcal{I}_d} \left( \sum_{n=1}^{\infty} v_n \right) = l.$$



# The two-dimensional case, ideal version

## Theorem

Let  $l \subseteq \mathbb{R}^2$  be such line on the plane and  $\sum_{n=1}^{\infty} v_n$  such series in  $\mathbb{R}^2$  that

$$S \left( \sum_{n=1}^{\infty} v_n \right) = l.$$

Then

$$S_{\mathcal{I}_d} \left( \sum_{n=1}^{\infty} v_n \right) = l.$$

Alternatively, instead of  $\mathcal{I}_d$  you can put any ideal that has the (R) property.

# The two-dimensional case, ideal version

## Definition

*We say that an ideal  $\mathcal{I} \subseteq \mathcal{P}(\mathbb{N})$  has the  $(R_2)$  property if for any conditionally convergent series of vectors on the plane  $\sum_{n=1}^{\infty} v_n$  such that  $S(\sum_{n=1}^{\infty} v_n) = \mathbb{R}^2$  and any  $v \in \mathbb{R}^2$  there exists a permutation  $\sigma : \mathbb{N} \rightarrow \mathbb{N}$  such that  $\sum_{n=1}^{\infty} v_{\sigma(n)} = v$  and  $\text{supp}(\sigma) \in \mathcal{I}$ .*

# The two-dimensional case, ideal version

## Definition

We say that an ideal  $\mathcal{I} \subseteq \mathcal{P}(\mathbb{N})$  has the  $(R_2)$  property if for any conditionally convergent series of vectors on the plane  $\sum_{n=1}^{\infty} v_n$  such that  $S(\sum_{n=1}^{\infty} v_n) = \mathbb{R}^2$  and any  $v \in \mathbb{R}^2$  there exists a permutation  $\sigma : \mathbb{N} \rightarrow \mathbb{N}$  such that  $\sum_{n=1}^{\infty} v_{\sigma(n)} = v$  and  $\text{supp}(\sigma) \in \mathcal{I}$ .

## Theorem (Folklore)

Let  $(v_n)_n \subseteq \mathbb{R}^2$ ,  $v_n \rightarrow 0$ ,  $\forall w \neq 0 \sum_{n=1}^{\infty} (w \circ v_n)^+ = \infty$ . Then

$$S\left(\sum_{n=1}^{\infty} v_n\right) = \mathbb{R}^2.$$

# The two-dimensional case, ideal version

## Theorem

*Let  $(v_n)_n \subseteq \mathbb{R}^2$ ,  $v_n \rightarrow 0$ . The following are equivalent:*

- $S(\sum_{n=1}^{\infty} v_n) = \mathbb{R}^2$ .
- *The set  $\{\sum_{n \in F} v_n : F \subseteq \mathbb{N}, |F| < \aleph_0\}$  is dense in  $\mathbb{R}^2$ .*

# The two-dimensional case, ideal version

## Theorem

*Let  $(v_n)_n \subseteq \mathbb{R}^2$ ,  $v_n \rightarrow 0$ . The following are equivalent:*

- $S(\sum_{n=1}^{\infty} v_n) = \mathbb{R}^2$ .
- *The set  $\{\sum_{n \in F} v_n : F \subseteq \mathbb{N}, |F| < \aleph_0\}$  is dense in  $\mathbb{R}^2$ .*

## Theorem

*Let  $S(\sum_{n=1}^{\infty} v_n) = \mathbb{R}^2$ . There exists a set  $A \subseteq \mathbb{N}$  such that both series  $\sum_{n \in A} v_n$  and  $\sum_{n \in \mathbb{N} \setminus A} v_n$  are conditionally convergent and  $S(\sum_{n \in A} v_n) = \mathbb{R}^2$ ,  $S(\sum_{n \in \mathbb{N} \setminus A} v_n) = \mathbb{R}^2$ .*

# The two-dimensional case, ideal version

## Theorem

*Let  $(v_n)_n \subseteq \mathbb{R}^2$ ,  $v_n \rightarrow 0$ . The following are equivalent:*

- $S(\sum_{n=1}^{\infty} v_n) = \mathbb{R}^2$ .*
- The set  $\{\sum_{n \in F} v_n : F \subseteq \mathbb{N}, |F| < \aleph_0\}$  is dense in  $\mathbb{R}^2$ .*

## Theorem

*Let  $S(\sum_{n=1}^{\infty} v_n) = \mathbb{R}^2$ . There exists a set  $A \subseteq \mathbb{N}$  such that both series  $\sum_{n \in A} v_n$  and  $\sum_{n \in \mathbb{N} \setminus A} v_n$  are conditionally convergent and  $S(\sum_{n \in A} v_n) = \mathbb{R}^2$ ,  $S(\sum_{n \in \mathbb{N} \setminus A} v_n) = \mathbb{R}^2$ .*

## Corollary

*If  $\mathcal{I} \subseteq \mathcal{P}(\mathbb{N})$  is a maximal ideal, then it has the  $(R_2)$  property.*

# The two-dimensional case, ideal version

## Theorem

*Let  $\mathcal{I} \subseteq \mathcal{P}(\mathbb{N})$  be an ideal. The following are equivalent.*

- (i) If  $(v_n)_n \subseteq \mathbb{R}^2$ ,  $v_n \rightarrow 0$  is such that  
 $\forall w \neq 0 \sum_{n=1}^{\infty} (w \circ v_n)^+ = \infty$  then  
 $\exists A \in \mathcal{I} \forall w \neq 0 \sum_{n \in A} (w \circ v_n)^+ = \infty$ .*
- (ii)  $\mathcal{I}$  cannot be extended to a summable ideal.*

Thank you.