

# On microperiodic multifunctions

Jolanta Olko

Institute of Mathematics, Pedagogical University of Cracow

28th ISCRF 2014 Stara Lesna

### Definition 1

Let  $(X, \cdot)$  be a topological group. A function  $f : X \rightarrow Y$  is called *microperiodic* if there exists a dense subset  $P$  of  $X$  such that

$$f(px) = f(x), \quad x \in X, p \in P.$$

### Definition 1

Let  $(X, \cdot)$  be a topological group. A function  $f : X \rightarrow Y$  is called microperiodic if there exists a dense subset  $P$  of  $X$  such that

$$f(px) = f(x), \quad x \in X, p \in P.$$

### Proposition 1

Let  $(X, \cdot)$  be a topological group,  $P$  dense subset of  $X$ . If  $f : X \rightarrow \overline{\mathbb{R}}$  satisfying

$$f(px) \leq f(x), \quad x \in X, p \in P, \tag{m}$$

is continuous at some  $x_0 \in X$ , then it is constant.

### Definition 1

Let  $(X, \cdot)$  be a topological group. A function  $f : X \rightarrow Y$  is called *microperiodic* if there exists a dense subset  $P$  of  $X$  such that

$$f(px) = f(x), \quad x \in \mathbb{R}, p \in P.$$

### Proposition 1

Let  $(X, \cdot)$  be a topological group,  $P$  dense subset of  $X$ . If  $f : X \rightarrow \overline{\mathbb{R}}$  satisfying

$$f(px) \leq f(x), \quad x \in X, p \in P, \tag{m}$$

is continuous at some  $x_0 \in X$ , then it is constant.

### Definition 2

Let  $(X, \cdot)$  be a topological group. We say that a multifunction  $F : X \rightarrow 2^Y$  is *microperiodic* if there exists a dense subset  $P$  of  $X$  such that

$$F(px) \subset F(x), \quad x \in X, p \in P. \tag{M}$$

### Definition 1

Let  $(X, \cdot)$  be a topological group. A function  $f : X \rightarrow Y$  is called *microperiodic* if there exists a dense subset  $P$  of  $X$  such that

$$f(px) = f(x), \quad x \in X, p \in P.$$

### Proposition 1

Let  $(X, \cdot)$  be a topological group,  $P$  dense subset of  $X$ . If  $f : X \rightarrow \overline{\mathbb{R}}$  satisfying

$$f(px) \leq f(x), \quad x \in X, p \in P, \tag{m}$$

is continuous at some  $x_0 \in X$ , then it is constant.

### Definition 2

Let  $(X, \cdot)$  be a topological group. We say that a multifunction  $F : X \rightarrow 2^Y$  is *microperiodic* if there exists a dense subset  $P$  of  $X$  such that

$$F(px) \subset F(x), \quad x \in X, p \in P. \tag{M}$$

Let  $A, B \subset Y$ . Multifunction  $F : X \rightarrow 2^Y$  of the form

$$F(x) = \begin{cases} A & \text{for } x \in \mathbb{Q}, \\ B & \text{for } x \notin \mathbb{Q} \end{cases}$$

is microperiodic.

Let  $X, Y$  be topological spaces,  $F : X \rightarrow 2^Y$ .

Let  $X, Y$  be topological spaces,  $F : X \rightarrow 2^Y$ .

- $F$  is upper semicontinuous (usc) at  $x_0 \in X$  if for every open set  $V \subset Y$  such that  $F(x_0) \subset V$  there exists  $U \in \mathcal{N}(x_0)$  such that  $F(x) \subset V$  for every  $x \in U$ ,

Let  $X, Y$  be topological spaces,  $F : X \rightarrow 2^Y$ .

- $F$  is upper semicontinuous (usc) at  $x_0 \in X$  if for every open set  $V \subset Y$  such that  $F(x_0) \subset V$  there exists  $U \in \mathcal{N}(x_0)$  such that  $F(x) \subset V$  for every  $x \in U$ ,
- $F$  is lower semicontinuous (lsc) at  $x_0 \in X$  if for every open set  $V \subset Y$  such that  $F(x_0) \cap V \neq \emptyset$  there exists  $U \in \mathcal{N}(x_0)$  such that  $F(x) \cap V \neq \emptyset$  for every  $x \in U$ ,



Let  $X, Y$  be topological spaces,  $F : X \rightarrow 2^Y$ .

- $F$  is upper semicontinuous (usc) at  $x_0 \in X$  if for every open set  $V \subset Y$  such that  $F(x_0) \subset V$  there exists  $U \in \mathcal{N}(x_0)$  such that  $F(x) \subset V$  for every  $x \in U$ ,
- $F$  is lower semicontinuous (lsc) at  $x_0 \in X$  if for every open set  $V \subset Y$  such that  $F(x_0) \cap V \neq \emptyset$  there exists  $U \in \mathcal{N}(x_0)$  such that  $F(x) \cap V \neq \emptyset$  for every  $x \in U$ ,
- $F$  is continuous at  $x_0$  if it is upper and lower semicontinuous at this point.

Let  $X, Y$  be topological spaces,  $F : X \rightarrow 2^Y$ .

- $F$  is upper semicontinuous (usc) at  $x_0 \in X$  if for every open set  $V \subset Y$  such that  $F(x_0) \subset V$  there exists  $U \in \mathcal{N}(x_0)$  such that  $F(x) \subset V$  for every  $x \in U$ ,
- $F$  is lower semicontinuous (lsc) at  $x_0 \in X$  if for every open set  $V \subset Y$  such that  $F(x_0) \cap V \neq \emptyset$  there exists  $U \in \mathcal{N}(x_0)$  such that  $F(x) \cap V \neq \emptyset$  for every  $x \in U$ ,
- $F$  is continuous at  $x_0$  if it is upper and lower semicontinuous at this point.

If  $(Y, d)$  is a metric space,  $A, B \subset Y$ , define

$$e(A, B) := \sup\{d(a, B) : a \in A\},$$
$$h(A, B) := \max\{e(A, B), e(B, A)\},$$

where  $d(a, B) := \inf\{d(a, b), b \in B\}$ .

Let  $X, Y$  be topological spaces,  $F : X \rightarrow 2^Y$ .

- $F$  is upper semicontinuous (usc) at  $x_0 \in X$  if for every open set  $V \subset Y$  such that  $F(x_0) \subset V$  there exists  $U \in \mathcal{N}(x_0)$  such that  $F(x) \subset V$  for every  $x \in U$ ,
- $F$  is lower semicontinuous (lsc) at  $x_0 \in X$  if for every open set  $V \subset Y$  such that  $F(x_0) \cap V \neq \emptyset$  there exists  $U \in \mathcal{N}(x_0)$  such that  $F(x) \cap V \neq \emptyset$  for every  $x \in U$ ,
- $F$  is continuous at  $x_0$  if it is upper and lower semicontinuous at this point.

If  $(Y, d)$  is a metric space,  $A, B \subset Y$ , define

$$e(A, B) := \sup\{d(a, B) : a \in A\},$$

$$h(A, B) := \max\{e(A, B), e(B, A)\},$$

where  $d(a, B) := \inf\{d(a, b), b \in B\}$ .

Let  $X$  be a topological space,  $(Y, d)$  is a metric space,  $F : X \rightarrow P(Y)$ ,

- $F$  is  $h$ -upper semicontinuous ( $h$ -usc) at  $x_0 \in X$  if  $x \mapsto e(F(x), F(x_0))$  is continuous at  $x_0$ ,

Let  $X, Y$  be topological spaces,  $F : X \rightarrow 2^Y$ .

- $F$  is upper semicontinuous (usc) at  $x_0 \in X$  if for every open set  $V \subset Y$  such that  $F(x_0) \subset V$  there exists  $U \in \mathcal{N}(x_0)$  such that  $F(x) \subset V$  for every  $x \in U$ ,
- $F$  is lower semicontinuous (lsc) at  $x_0 \in X$  if for every open set  $V \subset Y$  such that  $F(x_0) \cap V \neq \emptyset$  there exists  $U \in \mathcal{N}(x_0)$  such that  $F(x) \cap V \neq \emptyset$  for every  $x \in U$ ,
- $F$  is continuous at  $x_0$  if it is upper and lower semicontinuous at this point.

If  $(Y, d)$  is a metric space,  $A, B \subset Y$ , define

$$e(A, B) := \sup\{d(a, B) : a \in A\},$$

$$h(A, B) := \max\{e(A, B), e(B, A)\},$$

where  $d(a, B) := \inf\{d(a, b), b \in B\}$ .

Let  $X$  be a topological space,  $(Y, d)$  is a metric space,  $F : X \rightarrow P(Y)$ ,

- $F$  is  $h$ -upper semicontinuous ( $h$ -usc) at  $x_0 \in X$  if  $x \mapsto e(F(x), F(x_0))$  is continuous at  $x_0$ ,
- $F$  is  $h$ -lower semicontinuous ( $h$ -lsc) at  $x_0$  if  $x \mapsto e(F(x_0), F(x))$  is continuous at  $x_0$ ,

Let  $X, Y$  be topological spaces,  $F : X \rightarrow 2^Y$ .

- $F$  is upper semicontinuous (usc) at  $x_0 \in X$  if for every open set  $V \subset Y$  such that  $F(x_0) \subset V$  there exists  $U \in \mathcal{N}(x_0)$  such that  $F(x) \subset V$  for every  $x \in U$ ,
- $F$  is lower semicontinuous (lsc) at  $x_0 \in X$  if for every open set  $V \subset Y$  such that  $F(x_0) \cap V \neq \emptyset$  there exists  $U \in \mathcal{N}(x_0)$  such that  $F(x) \cap V \neq \emptyset$  for every  $x \in U$ ,
- $F$  is continuous at  $x_0$  if it is upper and lower semicontinuous at this point.

If  $(Y, d)$  is a metric space,  $A, B \subset Y$ , define

$$e(A, B) := \sup\{d(a, B) : a \in A\},$$

$$h(A, B) := \max\{e(A, B), e(B, A)\},$$

where  $d(a, B) := \inf\{d(a, b), b \in B\}$ .

Let  $X$  be a topological space,  $(Y, d)$  is a metric space,  $F : X \rightarrow P(Y)$ ,

- $F$  is  $h$ -upper semicontinuous ( $h$ -usc) at  $x_0 \in X$  if  $x \mapsto e(F(x), F(x_0))$  is continuous at  $x_0$ ,
- $F$  is  $h$ -lower semicontinuous ( $h$ -lsc) at  $x_0$  if  $x \mapsto e(F(x_0), F(x))$  is continuous at  $x_0$ ,
- $F$  is  $h$ -continuous at  $x_0$  if it is  $h$ -usc and  $h$ -lsc at  $x_0$ .

### Example 1

$$1 \quad F_1(x) = \begin{cases} [0, 1] & \text{for } x \in \mathbb{Q}, \\ (0, 1) & \text{for } x \notin \mathbb{Q} \end{cases}$$

$$2 \quad F_2(x) = \begin{cases} (0, 1) & \text{for } x \in \mathbb{Q}, \\ (0, 1) \setminus \{\frac{1}{2}\} & \text{for } x \notin \mathbb{Q} \end{cases}$$

$$3 \quad F_3(x) = \begin{cases} [0, 1] & \text{for } x \in \mathbb{Q}, \\ \mathbb{Q} \cap [0, 1] & \text{for } x \notin \mathbb{Q} \end{cases}$$

### Example 1

$$1 \quad F_1(x) = \begin{cases} [0, 1] & \text{for } x \in \mathbb{Q}, \\ (0, 1) & \text{for } x \notin \mathbb{Q} \end{cases}$$

$$2 \quad F_2(x) = \begin{cases} (0, 1) & \text{for } x \in \mathbb{Q}, \\ (0, 1) \setminus \{\frac{1}{2}\} & \text{for } x \notin \mathbb{Q} \end{cases}$$

$$3 \quad F_3(x) = \begin{cases} [0, 1] & \text{for } x \in \mathbb{Q}, \\ \mathbb{Q} \cap [0, 1] & \text{for } x \notin \mathbb{Q} \end{cases}$$

$$h(F_i(x), F_i(y)) = 0, \quad x, y \in \mathbb{R}$$

### Example 1

$$1 \quad F_1(x) = \begin{cases} [0, 1] & \text{for } x \in \mathbb{Q}, \\ (0, 1) & \text{for } x \notin \mathbb{Q} \end{cases}$$

$$2 \quad F_2(x) = \begin{cases} (0, 1) & \text{for } x \in \mathbb{Q}, \\ (0, 1) \setminus \{\frac{1}{2}\} & \text{for } x \notin \mathbb{Q} \end{cases}$$

$$3 \quad F_3(x) = \begin{cases} [0, 1] & \text{for } x \in \mathbb{Q}, \\ \mathbb{Q} \cap [0, 1] & \text{for } x \notin \mathbb{Q} \end{cases}$$

$$h(F_i(x), F_i(y)) = 0, \quad x, y \in \mathbb{R}$$

- each  $F_i$  is  $h$ -continuous on  $\mathbb{R}$



### Example 1

$$1 \quad F_1(x) = \begin{cases} [0, 1] & \text{for } x \in \mathbb{Q}, \\ (0, 1) & \text{for } x \notin \mathbb{Q} \end{cases}$$

$$2 \quad F_2(x) = \begin{cases} (0, 1) & \text{for } x \in \mathbb{Q}, \\ (0, 1) \setminus \{\frac{1}{2}\} & \text{for } x \notin \mathbb{Q} \end{cases}$$

$$3 \quad F_3(x) = \begin{cases} [0, 1] & \text{for } x \in \mathbb{Q}, \\ \mathbb{Q} \cap [0, 1] & \text{for } x \notin \mathbb{Q} \end{cases}$$

$$h(F_i(x), F_i(y)) = 0, \quad x, y \in \mathbb{R}$$

- each  $F_i$  is  $h$ -continuous on  $\mathbb{R}$
- each  $F_i$  is lsc on  $\mathbb{R}$

### Example 1

$$1 \quad F_1(x) = \begin{cases} [0, 1] & \text{for } x \in \mathbb{Q}, \\ (0, 1) & \text{for } x \notin \mathbb{Q} \end{cases}$$

$$2 \quad F_2(x) = \begin{cases} (0, 1) & \text{for } x \in \mathbb{Q}, \\ (0, 1) \setminus \{\frac{1}{2}\} & \text{for } x \notin \mathbb{Q} \end{cases}$$

$$3 \quad F_3(x) = \begin{cases} [0, 1] & \text{for } x \in \mathbb{Q}, \\ \mathbb{Q} \cap [0, 1] & \text{for } x \notin \mathbb{Q} \end{cases}$$

$$h(F_i(x), F_i(y)) = 0, \quad x, y \in \mathbb{R}$$

- each  $F_i$  is  $h$ -continuous on  $\mathbb{R}$
- each  $F_i$  is lsc on  $\mathbb{R}$
- each  $F_i$  is usc only on  $\mathbb{Q}$

### Theorem 1

*Let  $X$  be a topological group,  $Y$  a metric space,  $F : X \rightarrow P(Y)$  a microperiodic multifunction. If  $F$  is  $h$ -continuous at some  $x_0$ , then the multifunction  $\overline{F}$  is constant.*

### Theorem 1

*Let  $X$  be a topological group,  $Y$  a metric space,  $F : X \rightarrow P(Y)$  a microperiodic multifunction. If  $F$  is  $h$ -continuous at some  $x_0$ , then the multifunction  $\overline{F}$  is constant.*

### Proof.

1 Define  $f(x) := e(F(x_0), F(x))$ ,  $x \in X$



### Theorem 1

*Let  $X$  be a topological group,  $Y$  a metric space,  $F : X \rightarrow P(Y)$  a microperiodic multifunction. If  $F$  is  $h$ -continuous at some  $x_0$ , then the multifunction  $\overline{F}$  is constant.*

### Proof.

- 1 Define  $f(x) := e(F(x_0), F(x))$ ,  $x \in X$
- 2  $f(x) \leq f(px)$ ,  $x \in X$ ,  $p \in P$



### Theorem 1

*Let  $X$  be a topological group,  $Y$  a metric space,  $F : X \rightarrow P(Y)$  a microperiodic multifunction. If  $F$  is  $h$ -continuous at some  $x_0$ , then the multifunction  $\overline{F}$  is constant.*

### Proof.

- 1 Define  $f(x) := e(F(x_0), F(x))$ ,  $x \in X$
- 2  $f(x) \leq f(px)$ ,  $x \in X$ ,  $p \in P$
- 3  $f$  is continuous at  $x_0$



### Theorem 1

*Let  $X$  be a topological group,  $Y$  a metric space,  $F : X \rightarrow P(Y)$  a microperiodic multifunction. If  $F$  is  $h$ -continuous at some  $x_0$ , then the multifunction  $\overline{F}$  is constant.*

### Proof.

- 1 Define  $f(x) := e(F(x_0), F(x))$ ,  $x \in X$
- 2  $f(x) \leq f(px)$ ,  $x \in X$ ,  $p \in P$
- 3  $f$  is continuous at  $x_0$
- 4  $f$  is constant, therefore  $e(F(x_0), F(x)) = f(x) = f(x_0) = e(F(x_0), F(x_0)) = 0$  for  $x \in X$



### Theorem 1

*Let  $X$  be a topological group,  $Y$  a metric space,  $F : X \rightarrow P(Y)$  a microperiodic multifunction. If  $F$  is  $h$ -continuous at some  $x_0$ , then the multifunction  $\overline{F}$  is constant.*

### Proof.

- 1 Define  $f(x) := e(F(x_0), F(x))$ ,  $x \in X$
- 2  $f(x) \leq f(px)$ ,  $x \in X$ ,  $p \in P$
- 3  $f$  is continuous at  $x_0$
- 4  $f$  is constant, therefore  $e(F(x_0), F(x)) = f(x) = f(x_0) = e(F(x_0), F(x_0)) = 0$  for  $x \in X$
- 5  $F(x_0) \subset \overline{F}(x)$ ,  $x \in X$





### Theorem 1

Let  $X$  be a topological group,  $Y$  a metric space,  $F : X \rightarrow P(Y)$  a microperiodic multifunction. If  $F$  is  $h$ -continuous at some  $x_0$ , then the multifunction  $\overline{F}$  is constant.

### Proof.

- 1 Define  $f(x) := e(F(x_0), F(x))$ ,  $x \in X$
- 2  $f(x) \leq f(px)$ ,  $x \in X$ ,  $p \in P$
- 3  $f$  is continuous at  $x_0$
- 4  $f$  is constant, therefore  $e(F(x_0), F(x)) = f(x) = f(x_0) = e(F(x_0), F(x_0)) = 0$  for  $x \in X$
- 5  $F(x_0) \subset \overline{F}(x)$ ,  $x \in X$
- 6  $g(x) := e(F(x), F(x_0))$ ,  $x \in X$



## Theorem 1

*Let  $X$  be a topological group,  $Y$  a metric space,  $F : X \rightarrow P(Y)$  a microperiodic multifunction. If  $F$  is  $h$ -continuous at some  $x_0$ , then the multifunction  $\overline{F}$  is constant.*

## Proof.

- 1 Define  $f(x) := e(F(x_0), F(x))$ ,  $x \in X$
- 2  $f(x) \leq f(px)$ ,  $x \in X$ ,  $p \in P$
- 3  $f$  is continuous at  $x_0$
- 4  $f$  is constant, therefore  $e(F(x_0), F(x)) = f(x) = f(x_0) = e(F(x_0), F(x_0)) = 0$  for  $x \in X$
- 5  $F(x_0) \subset \overline{F}(x)$ ,  $x \in X$
- 6  $g(x) := e(F(x), F(x_0))$ ,  $x \in X$
- 7  $F(x) \subset \overline{F}(x_0) \subset \overline{F}(x)$



## Theorem 2

*Assume that  $X$  is a topological group,  $Y$  a normed space,  $F : X \rightarrow P_{fc}(Y)$  is a microperiodic multifunction. If  $F$  is lsc at  $x_0$  and usc at  $x_0$  ( $Y$  endowed with the weak topology), then it is constant.*

## Theorem 2

*Assume that  $X$  is a topological group,  $Y$  a normed space,  $F : X \rightarrow P_{fc}(Y)$  is a microperiodic multifunction. If  $F$  is lsc at  $x_0$  and usc at  $x_0$  ( $Y$  endowed with the weak topology), then it is constant.*

## Proof.

1  $y^* \in Y^*$



## Theorem 2

*Assume that  $X$  is a topological group,  $Y$  a normed space,  $F : X \rightarrow P_{fc}(Y)$  is a microperiodic multifunction. If  $F$  is lsc at  $x_0$  and usc at  $x_0$  ( $Y$  endowed with the weak topology), then it is constant.*

## Proof.

1  $y^* \in Y^*$

2  $f_{y^*}(x) := \sigma(y^*, F(x)) = \sup\{(y^*, y) : y \in F(x)\}, x \in X$



## Theorem 2

*Assume that  $X$  is a topological group,  $Y$  a normed space,  $F : X \rightarrow P_{fc}(Y)$  is a microperiodic multifunction. If  $F$  is lsc at  $x_0$  and usc at  $x_0$  ( $Y$  endowed with the weak topology), then it is constant.*

## Proof.

- 1  $y^* \in Y^*$
- 2  $f_{y^*}(x) := \sigma(y^*, F(x)) = \sup\{(y^*, y) : y \in F(x)\}, x \in X$
- 3  $f_{y^*}$  is continuous at  $x_0$



## Theorem 2

*Assume that  $X$  is a topological group,  $Y$  a normed space,  $F : X \rightarrow P_{fc}(Y)$  is a microperiodic multifunction. If  $F$  is lsc at  $x_0$  and usc at  $x_0$  ( $Y$  endowed with the weak topology), then it is constant.*

## Proof.

- 1  $y^* \in Y^*$
- 2  $f_{y^*}(x) := \sigma(y^*, F(x)) = \sup\{(y^*, y) : y \in F(x)\}, x \in X$
- 3  $f_{y^*}$  is continuous at  $x_0$
- 4  $f_{y^*}(px) \leq f_{y^*}(x), x \in X, p \in P$



## Theorem 2

Assume that  $X$  is a topological group,  $Y$  a normed space,  $F : X \rightarrow P_{fc}(Y)$  is a microperiodic multifunction. If  $F$  is lsc at  $x_0$  and usc at  $x_0$  ( $Y$  endowed with the weak topology), then it is constant.

## Proof.

- 1  $y^* \in Y^*$
- 2  $f_{y^*}(x) := \sigma(y^*, F(x)) = \sup\{(y^*, y) : y \in F(x)\}, x \in X$
- 3  $f_{y^*}$  is continuous at  $x_0$
- 4  $f_{y^*}(px) \leq f_{y^*}(x), x \in X, p \in P$
- 5  $f_{y^*}$  is constant





## Theorem 2

Assume that  $X$  is a topological group,  $Y$  a normed space,  $F : X \rightarrow P_{fc}(Y)$  is a microperiodic multifunction. If  $F$  is lsc at  $x_0$  and usc at  $x_0$  ( $Y$  endowed with the weak topology), then it is constant.

## Proof.

- 1  $y^* \in Y^*$
- 2  $f_{y^*}(x) := \sigma(y^*, F(x)) = \sup\{(y^*, y) : y \in F(x)\}, x \in X$
- 3  $f_{y^*}$  is continuous at  $x_0$
- 4  $f_{y^*}(px) \leq f_{y^*}(x), x \in X, p \in P$
- 5  $f_{y^*}$  is constant
- 6  $\sigma(y^*, F(x)) = \sigma(y^*, F(x_0))$  for  $x \in X$



## Theorem 2

*Assume that  $X$  is a topological group,  $Y$  a normed space,  $F : X \rightarrow P_{fc}(Y)$  is a microperiodic multifunction. If  $F$  is lsc at  $x_0$  and usc at  $x_0$  ( $Y$  endowed with the weak topology), then it is constant.*

## Proof.

- 1  $y^* \in Y^*$
- 2  $f_{y^*}(x) := \sigma(y^*, F(x)) = \sup\{(y^*, y) : y \in F(x)\}, x \in X$
- 3  $f_{y^*}$  is continuous at  $x_0$
- 4  $f_{y^*}(px) \leq f_{y^*}(x), x \in X, p \in P$
- 5  $f_{y^*}$  is constant
- 6  $\sigma(y^*, F(x)) = \sigma(y^*, F(x_0))$  for  $x \in X$
- 7 by the separation theorem  $F(x) = F(x_0), x \in X$



### Theorem 3

*Assume that  $X$  is a topological group,  $Y$  a topological space,  $F : X \rightarrow P(Y)$  a microperiodic multifunction with open values. If  $F$  is usc on some open neighborhood of  $x_0$ , then it is constant.*



Let  $(X, \mathcal{M})$  be a measurable space,  $(Y, d)$  a separable metric space,  $F : X \rightarrow 2^Y$

- $F$  is weakly measurable (measurable for short) if the lower inverse image  $F^-(U) := \{x \in X : F(x) \cap U \neq \emptyset\}$  is measurable for every open set  $U \subset Y$
- $F$  is strongly measurable if the lower inverse image  $F^-(C) := \{x \in X : F(x) \cap C \neq \emptyset\}$  is measurable for every closed set  $C \subset Y$

Let  $(X, \mathcal{M})$  be a measurable space,  $(Y, d)$  a separable metric space,  $F : X \rightarrow 2^Y$

- $F$  is weakly measurable (measurable for short) if the lower inverse image  $F^-(U) := \{x \in X : F(x) \cap U \neq \emptyset\}$  is measurable for every open set  $U \subset Y$
- $F$  is strongly measurable if the lower inverse image  $F^-(C) := \{x \in X : F(x) \cap C \neq \emptyset\}$  is measurable for every closed set  $C \subset Y$

### Example 2

Let  $A, B \subset Y$ ,  $P \subset X$  is dense and for every  $p \in P$  the following condition holds

$$px \in P, \text{ for } x \in P \quad \text{and} \quad px \in X \setminus P, \text{ for } x \in X \setminus P.$$

Let  $(X, \mathcal{M})$  be a measurable space,  $(Y, d)$  a separable metric space,  $F : X \rightarrow 2^Y$

- $F$  is weakly measurable (measurable for short) if the lower inverse image  $F^-(U) := \{x \in X : F(x) \cap U \neq \emptyset\}$  is measurable for every open set  $U \subset Y$
- $F$  is strongly measurable if the lower inverse image  $F^-(C) := \{x \in X : F(x) \cap C \neq \emptyset\}$  is measurable for every closed set  $C \subset Y$

### Example 2

Let  $A, B \subset Y$ ,  $P \subset X$  is dense and for every  $p \in P$  the following condition holds

$$px \in P, \text{ for } x \in P \quad \text{and} \quad px \in X \setminus P, \text{ for } x \in X \setminus P.$$

$$F(x) = \begin{cases} A & \text{for } x \in P, \\ B & \text{for } x \in X \setminus P \end{cases}.$$

Let  $(X, \mathcal{M})$  be a measurable space,  $(Y, d)$  a separable metric space,  $F : X \rightarrow 2^Y$

- $F$  is weakly measurable (measurable for short) if the lower inverse image  $F^-(U) := \{x \in X : F(x) \cap U \neq \emptyset\}$  is measurable for every open set  $U \subset Y$
- $F$  is strongly measurable if the lower inverse image  $F^-(C) := \{x \in X : F(x) \cap C \neq \emptyset\}$  is measurable for every closed set  $C \subset Y$

### Example 2

Let  $A, B \subset Y$ ,  $P \subset X$  is dense and for every  $p \in P$  the following condition holds

$$px \in P, \text{ for } x \in P \quad \text{and} \quad px \in X \setminus P, \text{ for } x \in X \setminus P.$$

$$F(x) = \begin{cases} A & \text{for } x \in P, \\ B & \text{for } x \in X \setminus P \end{cases}.$$

- $F$  is microperiodic



Let  $(X, \mathcal{M})$  be a measurable space,  $(Y, d)$  a separable metric space,  $F : X \rightarrow 2^Y$

- $F$  is weakly measurable (measurable for short) if the lower inverse image  $F^-(U) := \{x \in X : F(x) \cap U \neq \emptyset\}$  is measurable for every open set  $U \subset Y$
- $F$  is strongly measurable if the lower inverse image  $F^-(C) := \{x \in X : F(x) \cap C \neq \emptyset\}$  is measurable for every closed set  $C \subset Y$

### Example 2

Let  $A, B \subset Y$ ,  $P \subset X$  is dense and for every  $p \in P$  the following condition holds

$$px \in P, \text{ for } x \in P \quad \text{and} \quad px \in X \setminus P, \text{ for } x \in X \setminus P.$$

$$F(x) = \begin{cases} A & \text{for } x \in P, \\ B & \text{for } x \in X \setminus P \end{cases}.$$

- $F$  is microperiodic
- $F$  strongly measurable iff  $P \in \mathcal{M}$

Let  $(X, \mathcal{M})$  be a measurable space,  $(Y, d)$  a separable metric space,  $F : X \rightarrow 2^Y$

- $F$  is weakly measurable (measurable for short) if the lower inverse image  $F^-(U) := \{x \in X : F(x) \cap U \neq \emptyset\}$  is measurable for every open set  $U \subset Y$
- $F$  is strongly measurable if the lower inverse image  $F^-(C) := \{x \in X : F(x) \cap C \neq \emptyset\}$  is measurable for every closed set  $C \subset Y$

### Example 2

Let  $A, B \subset Y$ ,  $P \subset X$  is dense and for every  $p \in P$  the following condition holds

$$px \in P, \text{ for } x \in P \quad \text{and} \quad px \in X \setminus P, \text{ for } x \in X \setminus P.$$

$$F(x) = \begin{cases} A & \text{for } x \in P, \\ B & \text{for } x \in X \setminus P \end{cases}.$$

- $F$  is microperiodic
- $F$  strongly measurable iff  $P \in \mathcal{M}$

### Example 3

Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be an additive function with small graph (i.e.  $\text{card } f(\mathbb{R}) = \aleph_0$ ).

Let  $(X, \mathcal{M})$  be a measurable space,  $(Y, d)$  a separable metric space,  $F : X \rightarrow 2^Y$

- $F$  is weakly measurable (measurable for short) if the lower inverse image  $F^-(U) := \{x \in X : F(x) \cap U \neq \emptyset\}$  is measurable for every open set  $U \subset Y$
- $F$  is strongly measurable if the lower inverse image  $F^-(C) := \{x \in X : F(x) \cap C \neq \emptyset\}$  is measurable for every closed set  $C \subset Y$

### Example 2

Let  $A, B \subset Y$ ,  $P \subset X$  is dense and for every  $p \in P$  the following condition holds

$$px \in P, \text{ for } x \in P \quad \text{and} \quad px \in X \setminus P, \text{ for } x \in X \setminus P.$$

$$F(x) = \begin{cases} A & \text{for } x \in P, \\ B & \text{for } x \in X \setminus P \end{cases}.$$

- $F$  is microperiodic
- $F$  strongly measurable iff  $P \in \mathcal{M}$

### Example 3

Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be an additive function with small graph (i.e.  $\text{card } f(\mathbb{R}) = \aleph_0$ ). Assume that  $C \in P(\mathbb{R})$  such that  $d(0, C) > 0$ .

Let  $(X, \mathcal{M})$  be a measurable space,  $(Y, d)$  a separable metric space,  $F : X \rightarrow 2^Y$

- $F$  is weakly measurable (measurable for short) if the lower inverse image  $F^-(U) := \{x \in X : F(x) \cap U \neq \emptyset\}$  is measurable for every open set  $U \subset Y$
- $F$  is strongly measurable if the lower inverse image  $F^-(C) := \{x \in X : F(x) \cap C \neq \emptyset\}$  is measurable for every closed set  $C \subset Y$

### Example 2

Let  $A, B \subset Y$ ,  $P \subset X$  is dense and for every  $p \in P$  the following condition holds

$$px \in P, \text{ for } x \in P \quad \text{and} \quad px \in X \setminus P, \text{ for } x \in X \setminus P.$$

$$F(x) = \begin{cases} A & \text{for } x \in P, \\ B & \text{for } x \in X \setminus P \end{cases}.$$

- $F$  is microperiodic
- $F$  strongly measurable iff  $P \in \mathcal{M}$

### Example 3

Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be an additive function with small graph (i.e.  $\text{card } f(\mathbb{R}) = \aleph_0$ ). Assume that  $C \in P(\mathbb{R})$  such that  $d(0, C) > 0$ .

$$F(x) = |f(x)| \cdot C = \{|f(x)| \cdot c : c \in C\}, \quad x \in \mathbb{R}$$

### Theorem 4

Let  $X$  be a topological group,  $(Y, d)$  a separable metric space,  $P$  a dense subset of  $X$ . Assume that  $X$  is locally compact (of the second category, respectively),  $F : X \rightarrow P(Y)$  is  $\mathcal{H}$ -measurable ( $\mathcal{B}$ -measurable) satisfying one of the conditions

- (i)  $F$  is microperiodic multifunction,
- (ii)  $P$  is countable and  $F(px) \subset F(x)$  a.e. for every  $p \in P$ .

Then  $\overline{F}$  is constant a.e..

### Theorem 4

Let  $X$  be a topological group,  $(Y, d)$  a separable metric space,  $P$  a dense subset of  $X$ . Assume that  $X$  is locally compact (of the second category, respectively),  $F : X \rightarrow P(Y)$  is  $\mathcal{H}$ -measurable ( $\mathcal{B}$ -measurable) satisfying one of the conditions

- (i)  $F$  is microperiodic multifunction,
- (ii)  $P$  is countable and  $F(px) \subset F(x)$  a.e. for every  $p \in P$ .

Then  $\overline{F}$  is constant a.e..

### Theorem 5

Let  $I \subset \mathbb{R}$  be a nontrivial interval,  $P$  a dense subset of  $\mathbb{R}$ . If  $F : I \rightarrow P(\mathbb{R})$  is  $\mathcal{L}$ -measurable ( $\mathcal{B}$ -measurable, respectively) and there exists  $E \in \mathcal{L}_0$  ( $E \in \mathcal{B}_0$ ) such that

$$F(x + p) \subset F(x) \text{ for every } x \in I \setminus E, p \in P \text{ such that } x + p \in I,$$

then  $\overline{F}$  is constant a.e..

**Proposition 2 (Theorem 2 in [5])**

*Assume that  $X$  is a topological group,  $P \subset X$  a dense set,  $(Y, d)$  is a metric space. Let  $\varepsilon \geq 0$  and let  $f : X \rightarrow Y$  satisfies*

$$d(f(px), f(x)) \leq \varepsilon, \quad x \in X, p \in P.$$

*If  $f$  is continuous at some  $x_0$ , then*

$$d(f(x), f(x_0)) \leq \varepsilon, \quad x \in X.$$

### Proposition 2 (Theorem 2 in [5])

Assume that  $X$  is a topological group,  $P \subset X$  a dense set,  $(Y, d)$  is a metric space. Let  $\varepsilon \geq 0$  and let  $f : X \rightarrow Y$  satisfies

$$d(f(px), f(x)) \leq \varepsilon, \quad x \in X, p \in P.$$

If  $f$  is continuous at some  $x_0$ , then

$$d(f(x), f(x_0)) \leq \varepsilon, \quad x \in X.$$

### Theorem 6

Assume that  $X$  is a topological group,  $P \subset X$  a dense set,  $Y$  is a metric space. Let  $\varepsilon \geq 0$  and let  $F : X \rightarrow P(Y)$  satisfies

$$e(F(x), F(px)) \leq \varepsilon, \quad x \in X, p \in P. \quad (1)$$

If  $F$  is  $h$ -continuous at some  $x_0$ , then

$$h(F(x), F(x_0)) \leq \varepsilon, \quad x \in X.$$



## Corollary 1

Assume that  $X$  is a topological group,  $P \subset X$  a dense set,  $Y$  is a normed space. Let  $C \subset Y$  and let  $F : X \rightarrow P(Y)$  satisfies

$$F(px) \subset F(x) + C, \quad x \in X, p \in P.$$

If  $F$  is  $h$ -continuous at some  $x_0$  and there exists  $y_0 \in Y$  and  $\varepsilon > 0$  such that  $C \subset y_0 + \varepsilon S$  ( $S$  is the closed unit ball in  $Y$ ), then

$$h(F(x), F(x_0)) \leq \|y_0\| + \varepsilon, \quad x \in X.$$

## Corollary 1

Assume that  $X$  is a topological group,  $P \subset X$  a dense set,  $Y$  is a normed space. Let  $C \subset Y$  and let  $F : X \rightarrow P(Y)$  satisfies

$$F(px) \subset F(x) + C, \quad x \in X, p \in P.$$

If  $F$  is  $h$ -continuous at some  $x_0$  and there exists  $y_0 \in Y$  and  $\varepsilon > 0$  such that  $C \subset y_0 + \varepsilon S$  ( $S$  is the closed unit ball in  $Y$ ), then

$$h(F(x), F(x_0)) \leq \|y_0\| + \varepsilon, \quad x \in X.$$

## Corollary 2

Assume that  $X$  is a topological group,  $P \subset X$  a dense set,  $Y$  is a normed space. Let  $C \subset Y$  and let  $f : X \rightarrow Y$  satisfies

$$f(px) \in f(x) + C, \quad x \in X, p \in P.$$

If  $f$  is continuous at some  $x_0$  and there exists  $y_0 \in Y$  and  $\varepsilon > 0$  such that  $C \subset y_0 + \varepsilon S$  ( $S$  is the closed unit ball in  $Y$ ), then

$$\|f(x) - f(x_0)\| \leq \|y_0\| + \varepsilon, \quad x \in X.$$

# References I

- [1] Aubin, J.P., Frankowska, H.: Set-Valued Analysis. Birkhauser, Boston, Basel, Berlin (1990)
- [2] Berge, C.: Topological Spaces: Including a Treatment of Multi-Valued Functions, Vector Spaces and Convexity. Dover Books (2010)
- [3] Brzdęk, J.: On Functions Satisfying some Inequalities. Abh. Math. Sem. Hamburg. 63, 277-281 (1993)
- [4] Brzdęk J.: Generalizations of some results concerning microperiodic mappings. Manuscripta Math. 265–276, (2006).
- [5] Brzdęk, J.: On approximately microperiodic mappings. Acta. Math. Hungar. 117(1-2), 179-186 (2007)
- [6] Castaing, C., Valadier, M.: Convex analysis and measurable multifunctions. Lecture Notes in Math. vol. 580, Springer, Berlin, New York (1977)
- [7] Ger R., Kominek Z., Sablik M.: Generalized Smital's Lemma and a Theorem of Steinhaus. Radovi Mat. 101–119, (1985).

## References II

- [8] Hu, S., Papageorgiu, N.S.: Handbook of Multivalued Analysis. Vol. I: Theory, Mathematics and Its Applications. Vol. 419, Kluwer Academic Publishers, Dordrecht, Boston, London (1977)
- [9] Kuczma M.: An introduction to the theory of functional equations and inequalities. Prace Naukowe Uniwersytetu Śląskiego, Vol. 489, PWN Warszawa, Kraków, Katowice (1985).
- [10] Kuczma M.: Note on microperiodic functions. Rad. Mat. 5, 127–140 (1989)
- [11] Laczovich M.: Operators commuting with translations, and systems of difference equations. Coll. Math. Vol. 80(1), 1–22 (1999)
- [12] Łomnicki A.: O wielookresowych funkcjach jednoznacznych zmiennej rzeczywistej. Sprawozdania Tow. Nauk. Warsz. 808–846, (1918).
- [13] Semadeni Z.: Periods of Measurable Functions and the Stone-Čech Compactification. Amer. Math. Monthly Vol. 71, 891–893, (1964).
- [14] Xenikakis P.J.: Results on Similarities in  $\mathbb{R}^N$ . J. Math. Anal. Appl. Vol. 131, 554–564 (1988).

- 1 Microperiodic multifunctions
- 2 Continuity
- 3 Measurability
- 4 Approximately microperiodic multifunctions