

On openness of bilinear mappings

Adam Majchrzycki

Łódź University of Technology, Poland

28th Summer Conference on Real Functions Theory, Stara Lesna, 2014

Definitions

Let X and Y be metric spaces. A mapping $T: X \rightarrow Y$ is called

- **open** if it maps every open set onto an open set;
- **open at a point** $x \in X$ if $T(x) \in \text{int } T[V]$ for each neighbourhood V of x .

Fact

A mapping $T: X \rightarrow Y$ is open if and only if it is open at every point.

Theorem (Banach openness principle)

Every continuous linear operator from a Banach space onto a Banach space is an open mapping.

Definitions

Let X and Y be metric spaces. A mapping $T: X \rightarrow Y$ is called

- **open** if it maps every open set onto an open set;
- **open at a point** $x \in X$ if $T(x) \in \text{int } T[V]$ for each neighbourhood V of x .

Fact

A mapping $T: X \rightarrow Y$ is open if and only if it is open at every point.

Theorem (Banach openness principle)

Every continuous linear operator from a Banach space onto a Banach space is an open mapping.

Definitions

Let X and Y be metric spaces. A mapping $T: X \rightarrow Y$ is called

- **open** if it maps every open set onto an open set;
- **open at a point** $x \in X$ if $T(x) \in \text{int } T[V]$ for each neighbourhood V of x .

Fact

A mapping $T: X \rightarrow Y$ is open if and only if it is open at every point.

Theorem (Banach openness principle)

Every continuous linear operator from a Banach space onto a Banach space is an open mapping.

Definitions

Let X and Y be metric spaces. A mapping $T: X \rightarrow Y$ is called

- **open** if it maps every open set onto an open set;
- **open at a point** $x \in X$ if $T(x) \in \text{int } T[V]$ for each neighbourhood V of x .

Fact

A mapping $T: X \rightarrow Y$ is open if and only if it is open at every point.

Theorem (Banach openness principle)

Every continuous linear operator from a Banach space onto a Banach space is an open mapping.

Definitions

Let X and Y be metric spaces. A mapping $T: X \rightarrow Y$ is called

- **open** if it maps every open set onto an open set;
- **open at a point** $x \in X$ if $T(x) \in \text{int } T[V]$ for each neighbourhood V of x .

Fact

A mapping $T: X \rightarrow Y$ is open if and only if it is open at every point.

Theorem (Banach openness principle)

Every continuous linear operator from a Banach space onto a Banach space is an open mapping.

Fact

The bilinear analogue of the Banach openness principle is false.

Indeed,

- **W. Rudin**: The mapping $T: \mathbb{R} \times \mathbb{R}^2 \rightarrow \mathbb{R}^2$, $T(t,x,y):=(tx,ty)$, is not open at $(0,1,1)$;
- **P.J. Cohen (1974)** and **C. Horowitz (1975)** found bilinear continuous surjections that are not open at 0 (finite dimensional, and infinite dimensional cases).
- Multiplication in $C[0,1]$ is not open, for instance at (f,f) where $f(x) = x - 1/2$; **D.H. Fremlin** (oral communication) (see also, **M. Balcerzak, A. Wachowicz, W. Wilczyński, 2005**).

Fact

The bilinear analogue of the Banach openness principle is false.

Indeed,

- **W. Rudin**: The mapping $T: \mathbb{R} \times \mathbb{R}^2 \rightarrow \mathbb{R}^2$, $T(t,x,y):=(tx,ty)$, is not open at $(0,1,1)$;
- **P.J. Cohen (1974)** and **C. Horowitz (1975)** found bilinear continuous surjections that are not open at 0 (finite dimensional, and infinite dimensional cases).
- Multiplication in $C[0,1]$ is not open, for instance at (f,f) where $f(x) = x - 1/2$; **D.H. Fremlin** (oral communication) (see also, **M. Balcerzak, A. Wachowicz, W. Wilczyński, 2005**).

Fact

The bilinear analogue of the Banach openness principle is false.

Indeed,

- **W. Rudin**: The mapping $T: \mathbb{R} \times \mathbb{R}^2 \rightarrow \mathbb{R}^2$, $T(t,x,y):=(tx,ty)$, is not open at $(0,1,1)$;
- **P.J. Cohen (1974)** and **C. Horowitz (1975)** found bilinear continuous surjections that are not open at 0 (finite dimensional, and infinite dimensional cases).
- Multiplication in $C[0,1]$ is not open, for instance at (f,f) where $f(x) = x - 1/2$; **D.H. Fremlin** (oral communication) (see also, **M. Balcerzak, A. Wachowicz, W. Wilczyński, 2005**).

Fact

The bilinear analogue of the Banach openness principle is false.

Indeed,

- **W. Rudin**: The mapping $T: \mathbb{R} \times \mathbb{R}^2 \rightarrow \mathbb{R}^2$, $T(t,x,y):=(tx,ty)$, is not open at $(0,1,1)$;
- **P.J. Cohen (1974)** and **C. Horowitz (1975)** found bilinear continuous surjections that are not open at 0 (finite dimensional, and infinite dimensional cases).
- Multiplication in $C[0,1]$ is not open, for instance at (f,f) where $f(x) = x - 1/2$; **D.H. Fremlin** (oral communication) (see also, **M. Balcerzak, A. Wachowicz, W. Wilczyński, 2005**).

Fact

The bilinear analogue of the Banach openness principle is false.

Indeed,

- **W. Rudin**: The mapping $T: \mathbb{R} \times \mathbb{R}^2 \rightarrow \mathbb{R}^2$, $T(t,x,y):=(tx,ty)$, is not open at $(0, 1, 1)$;
- **P.J. Cohen (1974) and C. Horowitz (1975)** found bilinear continuous surjections that are not open at 0 (finite dimensional, and infinite dimensional cases).
- Multiplication in $C[0, 1]$ is not open, for instance at (f, f) where $f(x) = x - 1/2$; **D.H. Fremlin** (oral communication) (see also, **M. Balcerzak, A. Wachowicz, W. Wilczyński, 2005**).

The points of openness of multiplication in $C[0, 1]$ were characterized by [E. Behrends, 2011](#).

On the other hand, in the space $C_{\mathbb{C}}[0, 1]$ of continuous complex-valued functions, multiplication is open; [E. Behrends, 2011](#).

We have obtained some other positive results.

Theorem (M. Balcerzak, A. M., A. Wachowicz, 2013)

Let (X, \mathcal{S}, μ) be a measure space with $\mu(X) > 0$. Then the multiplication from $L_p \times L_q$ to L_1 (for $p \in [1, \infty]$, $1/p + 1/q = 1$) is an open continuous bilinear surjection. In particular, this holds for multiplication from $\ell_p \times \ell_q$ to ℓ_1 . Also, the multiplication from $\ell_1 \times c_0$ to ℓ_1 is an open bilinear surjection.

The points of openness of multiplication in $C[0, 1]$ were characterized by [E. Behrends, 2011](#).

On the other hand, in the space $C_{\mathbb{C}}[0, 1]$ of continuous complex-valued functions, multiplication is open; [E. Behrends, 2011](#).

We have obtained some other positive results.

Theorem (M. Balcerzak, A. M., A. Wachowicz, 2013)

Let (X, \mathcal{S}, μ) be a measure space with $\mu(X) > 0$. Then the multiplication from $L_p \times L_q$ to L_1 (for $p \in [1, \infty]$, $1/p + 1/q = 1$) is an open continuous bilinear surjection. In particular, this holds for multiplication from $\ell_p \times \ell_q$ to ℓ_1 . Also, the multiplication from $\ell_1 \times c_0$ to ℓ_1 is an open bilinear surjection.

The points of openness of multiplication in $C[0, 1]$ were characterized by [E. Behrends, 2011](#).

On the other hand, in the space $C_{\mathbb{C}}[0, 1]$ of continuous complex-valued functions, multiplication is open; [E. Behrends, 2011](#).

We have obtained some other positive results.

Theorem (M. Balcerzak, A. M., A. Wachowicz, 2013)

Let (X, \mathcal{S}, μ) be a measure space with $\mu(X) > 0$. Then the multiplication from $L_p \times L_q$ to L_1 (for $p \in [1, \infty]$, $1/p + 1/q = 1$) is an open continuous bilinear surjection. In particular, this holds for multiplication from $\ell_p \times \ell_q$ to ℓ_1 . Also, the multiplication from $\ell_1 \times c_0$ to ℓ_1 is an open bilinear surjection.

The points of openness of multiplication in $C[0, 1]$ were characterized by [E. Behrends, 2011](#).

On the other hand, in the space $C_{\mathbb{C}}[0, 1]$ of continuous complex-valued functions, multiplication is open; [E. Behrends, 2011](#).

We have obtained some other positive results.

Theorem (M. Balcerzak, A. M., A. Wachowicz, 2013)

Let (X, \mathcal{S}, μ) be a measure space with $\mu(X) > 0$. Then the multiplication from $L_p \times L_q$ to L_1 (for $p \in [1, \infty]$, $1/p + 1/q = 1$) is an open continuous bilinear surjection. In particular, this holds for multiplication from $\ell_p \times \ell_q$ to ℓ_1 . Also, the multiplication from $\ell_1 \times c_0$ to ℓ_1 is an open bilinear surjection.

The points of openness of multiplication in $C[0, 1]$ were characterized by [E. Behrends, 2011](#).

On the other hand, in the space $C_{\mathbb{C}}[0, 1]$ of continuous complex-valued functions, multiplication is open; [E. Behrends, 2011](#).

We have obtained some other positive results.

Theorem (M. Balcerzak, A. M., A. Wachowicz, 2013)

Let (X, \mathcal{S}, μ) be a measure space with $\mu(X) > 0$. Then the multiplication from $L_p \times L_q$ to L_1 (for $p \in [1, \infty]$, $1/p + 1/q = 1$) is an open continuous bilinear surjection. In particular, this holds for multiplication from $\ell_p \times \ell_q$ to ℓ_1 . Also, the multiplication from $\ell_1 \times c_0$ to ℓ_1 is an open bilinear surjection.

The points of openness of multiplication in $C[0, 1]$ were characterized by [E. Behrends, 2011](#).

On the other hand, in the space $C_{\mathbb{C}}[0, 1]$ of continuous complex-valued functions, multiplication is open; [E. Behrends, 2011](#).

We have obtained some other positive results.

Theorem (M. Balcerzak, A. M., A. Wachowicz, 2013)

Let (X, \mathcal{S}, μ) be a measure space with $\mu(X) > 0$. Then the multiplication from $L_p \times L_q$ to L_1 (for $p \in [1, \infty]$, $1/p + 1/q = 1$) is an open continuous bilinear surjection. In particular, this holds for multiplication from $\ell_p \times \ell_q$ to ℓ_1 . Also, the multiplication from $\ell_1 \times c_0$ to ℓ_1 is an open bilinear surjection.

Let X and Y be metric spaces and let $f: X \rightarrow Y$.

Remark

The mapping f is open if and only if

$$\forall x \in X \forall \varepsilon > 0 \exists \delta > 0 B(f(x), \delta) \subset f[B(x, \varepsilon)].$$

We introduce a stronger notion of openness.

Definition

The mapping f is called *uniformly open* whenever

$$\forall \varepsilon > 0 \exists \delta > 0 \forall x \in X B(f(x), \delta) \subset f[B(x, \varepsilon)].$$

Let X and Y be metric spaces and let $f: X \rightarrow Y$.

Remark

The mapping f is open if and only if

$$\forall x \in X \forall \varepsilon > 0 \exists \delta > 0 B(f(x), \delta) \subset f[B(x, \varepsilon)].$$

We introduce a stronger notion of openness.

Definition

The mapping f is called *uniformly open* whenever

$$\forall \varepsilon > 0 \exists \delta > 0 \forall x \in X B(f(x), \delta) \subset f[B(x, \varepsilon)].$$

Let X and Y be metric spaces and let $f: X \rightarrow Y$.

Remark

The mapping f is open if and only if

$$\forall x \in X \forall \varepsilon > 0 \exists \delta > 0 B(f(x), \delta) \subset f[B(x, \varepsilon)].$$

We introduce a stronger notion of openness.

Definition

The mapping f is called *uniformly open* whenever

$$\forall \varepsilon > 0 \exists \delta > 0 \forall x \in X B(f(x), \delta) \subset f[B(x, \varepsilon)].$$

Note that

- The function $f: \mathbb{R} \rightarrow (-\pi/2, \pi/2)$, $f(x) = \arctan x$, is open but not uniformly open.
- Every continuous linear operator from a Banach space onto a Banach space is a uniformly open mapping.

Fact (folklore?)

Let X, Y be Banach spaces. Then every nonzero bilinear continuous functional $T: X \times Y \rightarrow \mathbb{R}$ is a surjective open mapping.

Proposition (M. Balcerzak, A. M., F. Strobin)

Every nonzero bilinear continuous functional is uniformly open.

Note that

- The function $f: \mathbb{R} \rightarrow (-\pi/2, \pi/2)$, $f(x) = \arctan x$, is open but not uniformly open.
- Every continuous linear operator from a Banach space onto a Banach space is a uniformly open mapping.

Fact (folklore?)

Let X, Y be Banach spaces. Then every nonzero bilinear continuous functional $T: X \times Y \rightarrow \mathbb{R}$ is a surjective open mapping.

Proposition (M. Balcerzak, A. M., F. Stobin)

Every nonzero bilinear continuous functional is uniformly open.

Note that

- The function $f: \mathbb{R} \rightarrow (-\pi/2, \pi/2)$, $f(x) = \arctan x$, is open but not uniformly open.
- Every continuous linear operator from a Banach space onto a Banach space is a uniformly open mapping.

Fact (folklore?)

Let X, Y be Banach spaces. Then every nonzero bilinear continuous functional $T: X \times Y \rightarrow \mathbb{R}$ is a surjective open mapping.

Proposition (M. Balcerzak, A. M., F. Strobin)

Every nonzero bilinear continuous functional is uniformly open.

Note that

- The function $f: \mathbb{R} \rightarrow (-\pi/2, \pi/2)$, $f(x) = \arctan x$, is open but not uniformly open.
- Every continuous linear operator from a Banach space onto a Banach space is a uniformly open mapping.

Fact (folklore?)

Let X, Y be Banach spaces. Then every nonzero bilinear continuous functional $T: X \times Y \rightarrow \mathbb{R}$ is a surjective open mapping.

Proposition (M.Balcerzak, A. M., F. Strobil)

Every nonzero bilinear continuous functional is uniformly open.

Note that

- The function $f: \mathbb{R} \rightarrow (-\pi/2, \pi/2)$, $f(x) = \arctan x$, is open but not uniformly open.
- Every continuous linear operator from a Banach space onto a Banach space is a uniformly open mapping.

Fact (folklore?)

Let X, Y be Banach spaces. Then every nonzero bilinear continuous functional $T: X \times Y \rightarrow \mathbb{R}$ is a surjective open mapping.

Proposition (M. Balcerzak, A. M., F. Strobil)

Every nonzero bilinear continuous functional is uniformly open.

Recently, we have improved our former result.

Theorem (M.Balcerzak, A. M., F. Strobin)

Let (X, \mathcal{S}, μ) be a measure space with $\mu(X) > 0$. Then the multiplication from $L_p \times L_q$ to L_1 (for $p \in [1, \infty]$, $1/p + 1/q = 1$) is a uniformly open mapping. In particular, this holds for multiplication from $\ell_p \times \ell_q$ to ℓ_1 .

Recently, we have improved our former result.

Theorem (M.Balcerzak, A. M., F. Strobin)

Let (X, \mathcal{S}, μ) be a measure space with $\mu(X) > 0$. Then the multiplication from $L_p \times L_q$ to L_1 (for $p \in [1, \infty]$, $1/p + 1/q = 1$) is a uniformly open mapping. In particular, this holds for multiplication from $\ell_p \times \ell_q$ to ℓ_1 .

Recently, we have improved our former result.

Theorem (M.Balcerzak, A. M., F. Strobin)

Let (X, \mathcal{S}, μ) be a measure space with $\mu(X) > 0$. Then the multiplication from $L_p \times L_q$ to L_1 (for $p \in [1, \infty]$, $1/p + 1/q = 1$) is a uniformly open mapping. In particular, this holds for multiplication from $\ell_p \times \ell_q$ to ℓ_1 .

Recently, we have improved our former result.

Theorem (M.Balcerzak, A. M., F. Strobin)

Let (X, \mathcal{S}, μ) be a measure space with $\mu(X) > 0$. Then the multiplication from $L_p \times L_q$ to L_1 (for $p \in [1, \infty]$, $1/p + 1/q = 1$) is a uniformly open mapping. In particular, this holds for multiplication from $\ell_p \times \ell_q$ to ℓ_1 .

Definition

We will say that (X, \mathcal{S}, μ) is a topological measure space if X is a topological space with all Borel sets belonging to \mathcal{S} .

We will assume that X is σ -compact.

Let L_∞^0 denote the space of all functions $f \in L_\infty$ which vanish at infinity, i.e.

for any $\varepsilon > 0$, there is a compact set $K \subset X$ such that $\mu(\{x \in X \setminus K : |f(x)| > \varepsilon\}) = 0$.

Then L_∞^0 is a closed linear subspace of L_∞ and so, L_∞^0 with the norm $\|\cdot\|_\infty$ is a Banach space.

Theorem (M. Balcerzak, A. M., F. Strobil)

Assume that (X, \mathcal{S}, μ) is a topological measure space where X is σ -compact. Then the multiplication from $L_1 \times L_\infty^0$ to L_1 is a continuous surjection which is a uniformly open mapping.

Definition

We will say that (X, \mathcal{S}, μ) is a topological measure space if X is a topological space with all Borel sets belonging to \mathcal{S} .

We will assume that X is σ -compact.

Let L_∞^0 denote the space of all functions $f \in L_\infty$ which vanish at infinity, i.e.

for any $\varepsilon > 0$, there is a compact set $K \subset X$ such that $\mu(\{x \in X \setminus K : |f(x)| > \varepsilon\}) = 0$.

Then L_∞^0 is a closed linear subspace of L_∞ and so, L_∞^0 with the norm $\|\cdot\|_\infty$ is a Banach space.

Theorem (M. Balcerzak, A. M., F. Strobil)

Assume that (X, \mathcal{S}, μ) is a topological measure space where X is σ -compact. Then the multiplication from $L_1 \times L_\infty^0$ to L_1 is a continuous surjection which is a uniformly open mapping.

Definition

We will say that (X, \mathcal{S}, μ) is a topological measure space if X is a topological space with all Borel sets belonging to \mathcal{S} .

We will assume that X is σ -compact.

Let L_∞^0 denote the space of all functions $f \in L_\infty$ which vanish at infinity, i.e.

for any $\varepsilon > 0$, there is a compact set $K \subset X$ such that $\mu(\{x \in X \setminus K : |f(x)| > \varepsilon\}) = 0$.

Then L_∞^0 is a closed linear subspace of L_∞ and so, L_∞^0 with the norm $\|\cdot\|_\infty$ is a Banach space.

Theorem (M. Balcerzak, A. M., F. Strobil)

Assume that (X, \mathcal{S}, μ) is a topological measure space where X is σ -compact. Then the multiplication from $L_1 \times L_\infty^0$ to L_1 is a continuous surjection which is a uniformly open mapping.

Definition

We will say that (X, \mathcal{S}, μ) is a topological measure space if X is a topological space with all Borel sets belonging to \mathcal{S} .

We will assume that X is σ -compact.

Let L_∞^0 denote the space of all functions $f \in L_\infty$ which vanish at infinity, i.e.

for any $\varepsilon > 0$, there is a compact set $K \subset X$ such that $\mu(\{x \in X \setminus K : |f(x)| > \varepsilon\}) = 0$.

Then L_∞^0 is a closed linear subspace of L_∞ and so, L_∞^0 with the norm $\|\cdot\|_\infty$ is a Banach space.

Theorem (M. Balcerzak, A. M., F. Strobil)

Assume that (X, \mathcal{S}, μ) is a topological measure space where X is σ -compact. Then the multiplication from $L_1 \times L_\infty^0$ to L_1 is a continuous surjection which is a uniformly open mapping.

Definition

We will say that (X, \mathcal{S}, μ) is a topological measure space if X is a topological space with all Borel sets belonging to \mathcal{S} .

We will assume that X is σ -compact.

Let L_∞^0 denote the space of all functions $f \in L_\infty$ which vanish at infinity, i.e.

for any $\varepsilon > 0$, there is a compact set $K \subset X$ such that $\mu(\{x \in X \setminus K : |f(x)| > \varepsilon\}) = 0$.

Then L_∞^0 is a closed linear subspace of L_∞ and so, L_∞^0 with the norm $\|\cdot\|_\infty$ is a Banach space.

Theorem (M. Balcerzak, A. M., F. Strobil)

Assume that (X, \mathcal{S}, μ) is a topological measure space where X is σ -compact. Then the multiplication from $L_1 \times L_\infty^0$ to L_1 is a continuous surjection which is a uniformly open mapping.

Definition

We will say that (X, \mathcal{S}, μ) is a topological measure space if X is a topological space with all Borel sets belonging to \mathcal{S} .

We will assume that X is σ -compact.

Let L_∞^0 denote the space of all functions $f \in L_\infty$ which vanish at infinity, i.e.

for any $\varepsilon > 0$, there is a compact set $K \subset X$ such that $\mu(\{x \in X \setminus K : |f(x)| > \varepsilon\}) = 0$.

Then L_∞^0 is a closed linear subspace of L_∞ and so, L_∞^0 with the norm $\|\cdot\|_\infty$ is a Banach space.

Theorem (M. Balcerzak, A. M., F. Strobil)

Assume that (X, \mathcal{S}, μ) is a topological measure space where X is σ -compact. Then the multiplication from $L_1 \times L_\infty^0$ to L_1 is a continuous surjection which is a uniformly open mapping.

For $X = \mathbb{N}$, $\mathcal{S} = \mathcal{P}(\mathbb{N})$ and the counting measure, we get

Corollary

The multiplication from $\ell_1 \times c_0$ to ℓ_1 is a uniformly open mapping.

Note that the statement of the Theorem remains true if we assume there is a σ -compact subset Y of X such that $\mu(X \setminus Y) = 0$.

We then call X *almost σ -compact*.

However, if we drop the assumption of the almost σ -compactness of X , the surjectivity of multiplication can be lost.

For $X = \mathbb{N}$, $\mathcal{S} = \mathcal{P}(\mathbb{N})$ and the counting measure, we get

Corollary

The multiplication from $\ell_1 \times c_0$ to ℓ_1 is a uniformly open mapping.

Note that the statement of the Theorem remains true if we assume there is a σ -compact subset Y of X such that $\mu(X \setminus Y) = 0$.

We then call X *almost σ -compact*.

However, if we drop the assumption of the almost σ -compactness of X , the surjectivity of multiplication can be lost.

For $X = \mathbb{N}$, $\mathcal{S} = \mathcal{P}(\mathbb{N})$ and the counting measure, we get

Corollary

The multiplication from $\ell_1 \times c_0$ to ℓ_1 is a uniformly open mapping.

Note that the statement of the Theorem remains true if we assume there is a σ -compact subset Y of X such that $\mu(X \setminus Y) = 0$.

We then call X *almost σ -compact*.

However, if we drop the assumption of the almost σ -compactness of X , the surjectivity of multiplication can be lost.

For $X = \mathbb{N}$, $\mathcal{S} = \mathcal{P}(\mathbb{N})$ and the counting measure, we get

Corollary

The multiplication from $\ell_1 \times c_0$ to ℓ_1 is a uniformly open mapping.

Note that the statement of the Theorem remains true if we assume there is a σ -compact subset Y of X such that $\mu(X \setminus Y) = 0$.

We then call X *almost σ -compact*.

However, if we drop the assumption of the almost σ -compactness of X , the surjectivity of multiplication can be lost.

For $X = \mathbb{N}$, $\mathcal{S} = \mathcal{P}(\mathbb{N})$ and the counting measure, we get

Corollary

The multiplication from $\ell_1 \times c_0$ to ℓ_1 is a uniformly open mapping.

Note that the statement of the Theorem remains true if we assume there is a σ -compact subset Y of X such that $\mu(X \setminus Y) = 0$.

We then call X *almost σ -compact*.

However, if we drop the assumption of the almost σ -compactness of X , the surjectivity of multiplication can be lost.

Example

Let $X = [0, 1)$ with the Sorgenfrey topology and the Lebesgue measure. Then the Borel sets are the same as in the natural topology, and all compact sets are countable. Hence this measure space is not almost σ -compact.

Let $h(x) := 1$ for every $x \in X$. Then $h \in L_1$, and if $h = fg$ then $g(x) \neq 0$ for every $x \in X$.

Suppose that $g \in L_\infty^0$. Then for each $n \in \mathbb{N}$ there is a compact set K_n such that $\mu(\{x \in X \setminus K_n : |g(x)| > 1/n\}) = 0$. Hence

$$X = \{x \in X : g(x) \neq 0\} = \bigcup_{n \in \mathbb{N}} K_n \cup \bigcup_{n \in \mathbb{N}} \{x \in X \setminus K_n : |g(x)| > 1/n\}.$$

It follows that X is of measure zero, a contradiction.

Example

Let $X = [0, 1)$ with the Sorgenfrey topology and the Lebesgue measure. Then the Borel sets are the same as in the natural topology, and all compact sets are countable. Hence this measure space is not almost σ -compact.

Let $h(x) := 1$ for every $x \in X$. Then $h \in L_1$, and if $h = fg$ then $g(x) \neq 0$ for every $x \in X$.

Suppose that $g \in L_\infty^0$. Then for each $n \in \mathbb{N}$ there is a compact set K_n such that $\mu(\{x \in X \setminus K_n : |g(x)| > 1/n\}) = 0$. Hence

$$X = \{x \in X : g(x) \neq 0\} = \bigcup_{n \in \mathbb{N}} K_n \cup \bigcup_{n \in \mathbb{N}} \{x \in X \setminus K_n : |g(x)| > 1/n\}.$$

It follows that X is of measure zero, a contradiction.

Example

Let $X = [0, 1)$ with the Sorgenfrey topology and the Lebesgue measure. Then the Borel sets are the same as in the natural topology, and all compact sets are countable. Hence this measure space is not almost σ -compact.

Let $h(x) := 1$ for every $x \in X$. Then $h \in L_1$, and if $h = fg$ then $g(x) \neq 0$ for every $x \in X$.

Suppose that $g \in L_\infty^0$. Then for each $n \in \mathbb{N}$ there is a compact set K_n such that $\mu(\{x \in X \setminus K_n : |g(x)| > 1/n\}) = 0$. Hence

$$X = \{x \in X : g(x) \neq 0\} = \bigcup_{n \in \mathbb{N}} K_n \cup \bigcup_{n \in \mathbb{N}} \{x \in X \setminus K_n : |g(x)| > 1/n\}.$$

It follows that X is of measure zero, a contradiction.

Example

Let $X = [0, 1)$ with the Sorgenfrey topology and the Lebesgue measure. Then the Borel sets are the same as in the natural topology, and all compact sets are countable. Hence this measure space is not almost σ -compact.

Let $h(x) := 1$ for every $x \in X$. Then $h \in L_1$, and if $h = fg$ then $g(x) \neq 0$ for every $x \in X$.

Suppose that $g \in L_\infty^0$. Then for each $n \in \mathbb{N}$ there is a compact set K_n such that $\mu(\{x \in X \setminus K_n : |g(x)| > 1/n\}) = 0$. Hence

$$X = \{x \in X : g(x) \neq 0\} = \bigcup_{n \in \mathbb{N}} K_n \cup \bigcup_{n \in \mathbb{N}} \{x \in X \setminus K_n : |g(x)| > 1/n\}.$$

It follows that X is of measure zero, a contradiction.

Example

Let $X = [0, 1)$ with the Sorgenfrey topology and the Lebesgue measure. Then the Borel sets are the same as in the natural topology, and all compact sets are countable. Hence this measure space is not almost σ -compact.

Let $h(x) := 1$ for every $x \in X$. Then $h \in L_1$, and if $h = fg$ then $g(x) \neq 0$ for every $x \in X$.

Suppose that $g \in L_\infty^0$. Then for each $n \in \mathbb{N}$ there is a compact set K_n such that $\mu(\{x \in X \setminus K_n : |g(x)| > 1/n\}) = 0$. Hence

$$X = \{x \in X : g(x) \neq 0\} = \bigcup_{n \in \mathbb{N}} K_n \cup \bigcup_{n \in \mathbb{N}} \{x \in X \setminus K_n : |g(x)| > 1/n\}.$$

It follows that X is of measure zero, a contradiction.

Thank you for your attention.