


Periodic points of some maps of Jordan domains

Peter Maličský
Matej Bel University, Banská Bystrica
Slovakia


28th Summer School on Real Function Theory
Stará Lesná
August 31 - September 5, 2014

Notation

Let D be a closed domain in the plane the boundary of which consists of three arcs a , b and c .




Notation




Consider a partition $D = D_0 \cup D_1$ given by an arc d . Let $G : D \rightarrow D$ be a continuous map for which

Notation




Consider a partition $D = D_0 \cup D_1$ given by an arc d . Let $G : D \rightarrow D$ be a continuous map for which

- $G(a_0) = a = G(a_1)$,




Consider a partition $D = D_0 \cup D_1$ given by an arc d . Let $G : D \rightarrow D$ be a continuous map for which

- $G(a_0) = a = G(a_1)$,
- $G(b) = \{P_0\}$,




Consider a partition $D = D_0 \cup D_1$ given by an arc d . Let $G : D \rightarrow D$ be a continuous map for which

- $G(a_0) = a = G(a_1)$,
- $G(b) = \{P_0\}$,
- $G(c_0) = b = G(c_1)$,




Consider a partition $D = D_0 \cup D_1$ given by an arc d . Let $G : D \rightarrow D$ be a continuous map for which

- $G(a_0) = a = G(a_1)$,
- $G(b) = \{P_0\}$,
- $G(c_0) = b = G(c_1)$,
- G is injective on $D_0 \setminus b$ and D_1 ,



Consider a partition $D = D_0 \cup D_1$ given by an arc d . Let $G : D \rightarrow D$ be a continuous map for which

- $G(a_0) = a = G(a_1)$,
- $G(b) = \{P_0\}$,
- $G(c_0) = b = G(c_1)$,
- G is injective on $D_0 \setminus b$ and D_1 ,
- $G(D_0 \setminus b) = D \setminus \{P_0\}$ and $G(D_1) = D$.



Consider a partition $D = D_0 \cup D_1$ given by an arc d . Let $G : D \rightarrow D$ be a continuous map for which

- $G(a_0) = a = G(a_1)$,
- $G(b) = \{P_0\}$,
- $G(c_0) = b = G(c_1)$,
- G is injective on $D_0 \setminus b$ and D_1 ,
- $G(D_0 \setminus b) = D \setminus \{P_0\}$ and $G(D_1) = D$.

History of the problem

In 1993 A. N. Sharkovskii formulated some problems concerning the properties of the plain map $(x, y) \mapsto ((y - 2)^2, xy)$. One of the questions was about interior periodic points.

History of the problem

In 1993 A. N. Sharkovskii formulated some problems concerning the properties of the plain map $(x, y) \mapsto ((y - 2)^2, xy)$. One of the questions was about interior periodic points. This map leaves the plane triangle $\Delta = \{ (x, y) : 0 \leq x, 0 \leq y, x + y \leq 4 \}$ invariant.

History of the problem

In 1993 A. N. Sharkovskii formulated some problems concerning the properties of the plain map $(x, y) \mapsto ((y - 2)^2, xy)$. One of the questions was about interior periodic points. This map leaves the plane triangle $\Delta = \{ (x, y) : 0 \leq x, 0 \leq y, x + y \leq 4 \}$ invariant. Balibrea, García Guirao, Lampart and Llibre studied the map defined by

$$F : \Delta \rightarrow \Delta, (x, y) \mapsto (x(4 - x - y), xy) ,$$

History of the problem

In 1993 A. N. Sharkovskii formulated some problems concerning the properties of the plain map $(x, y) \mapsto ((y - 2)^2, xy)$. One of the questions was about interior periodic points. This map leaves the plane triangle $\Delta = \{ (x, y) : 0 \leq x, 0 \leq y, x + y \leq 4 \}$ invariant. Balibrea, García Guirao, Lampart and Llibre studied the map defined by

$$F : \Delta \rightarrow \Delta, (x, y) \mapsto (x(4 - x - y), xy) ,$$

which is conjugate with the map of Sharkovskii via the conjugation $H : (x, y) \mapsto (4 - x - y, x)$.

History of the problem

In 1993 A. N. Sharkovskii formulated some problems concerning the properties of the plain map $(x, y) \mapsto ((y - 2)^2, xy)$. One of the questions was about interior periodic points. This map leaves the plane triangle $\Delta = \{ (x, y) : 0 \leq x, 0 \leq y, x + y \leq 4 \}$ invariant. Balibrea, García Guirao, Lampart and Llibre studied the map defined by

$$F : \Delta \rightarrow \Delta, (x, y) \mapsto (x(4 - x - y), xy) ,$$

which is conjugate with the map of Sharkovskii via the conjugation $H : (x, y) \mapsto (4 - x - y, x)$. In 2006 they published a paper in which an interior periodic point with period 4 was found

History of the problem

In 1993 A. N. Sharkovskii formulated some problems concerning the properties of the plain map $(x, y) \mapsto ((y - 2)^2, xy)$. One of the questions was about interior periodic points. This map leaves the plane triangle $\Delta = \{ (x, y) : 0 \leq x, 0 \leq y, x + y \leq 4 \}$ invariant. Balibrea, García Guirao, Lampart and Llibre studied the map defined by

$$F : \Delta \rightarrow \Delta, (x, y) \mapsto (x(4 - x - y), xy) ,$$

which is conjugate with the map of Sharkovskii via the conjugation $H : (x, y) \mapsto (4 - x - y, x)$. In 2006 they published a paper in which an interior periodic point with period 4 was found and it was proved that there are no such points with period 2 and 3.

History of the problem

In 1993 A. N. Sharkovskii formulated some problems concerning the properties of the plain map $(x, y) \mapsto ((y - 2)^2, xy)$. One of the questions was about interior periodic points. This map leaves the plane triangle $\Delta = \{ (x, y) : 0 \leq x, 0 \leq y, x + y \leq 4 \}$ invariant. Balibrea, García Guirao, Lampart and Llibre studied the map defined by


$$F : \Delta \rightarrow \Delta, (x, y) \mapsto (x(4 - x - y), xy) ,$$

which is conjugate with the map of Sharkovskii via the conjugation $H : (x, y) \mapsto (4 - x - y, x)$. In 2006 they published a paper in which an interior periodic point with period 4 was found and it was proved that there are no such points with period 2 and 3.

Relationship between lower and interior periodic points

Theorem (Maličký 2012)


Let P be a lower *saddle* fixed point of the map F^n . Then there is an interior fixed point Q of F^n with *the same period and itinerary*, where the itinerary is considered with respect to the sets Δ_L and Δ_R .




Relationship between lower and interior periodic points

Theorem (Maličký 2012)

Let P be a lower *saddle* fixed point of the map F^n . Then there is an interior fixed point Q of F^n with *the same period and itinerary*, where the itinerary is considered with respect to the sets Δ_L and Δ_R .




Problem to be solved



Let $G : D \rightarrow D$ be a continuous map for which

- $G(a_0) = a = G(a_1)$,
- $G(b) = \{P_0\}$,


Problem to be solved



Let $G : D \rightarrow D$ be a continuous map for which

- $G(a_0) = a = G(a_1)$,
- $G(b) = \{P_0\}$,
- $G(c_0) = b = G(c_1)$,


Problem to be solved



Let $G : D \rightarrow D$ be a continuous map for which

- $G(a_0) = a = G(a_1)$,
- $G(b) = \{P_0\}$,
- $G(c_0) = b = G(c_1)$,
- G is injective on $D_0 \setminus b$ and D_1 ,


Problem to be solved



Let $G : D \rightarrow D$ be a continuous map for which

- $G(a_0) = a = G(a_1)$,
- $G(b) = \{P_0\}$,
- $G(c_0) = b = G(c_1)$,
- G is injective on $D_0 \setminus b$ and D_1 ,
- $G(D_0 \setminus b) = D \setminus \{P_0\}$ and $G(D_1) = D$.


Problem to be solved



Let $G : D \rightarrow D$ be a continuous map for which

- $G(a_0) = a = G(a_1)$,
- $G(b) = \{P_0\}$,
- $G(c_0) = b = G(c_1)$,
- G is injective on $D_0 \setminus b$ and D_1 ,
- $G(D_0 \setminus b) = D \setminus \{P_0\}$ and $G(D_1) = D$.

Problem to be solved




Let $G : D \rightarrow D$ be a continuous map for which

- $G(a_0) = a = G(a_1)$,
- $G(b) = \{P_0\}$,
- $G(c_0) = b = G(c_1)$,
- G is injective on $D_0 \setminus b$ and D_1 ,
- $G(D_0 \setminus b) = D \setminus \{P_0\}$ and $G(D_1) = D$.

We are interested in periodic points of the map G .

Problem to be solved




Let $G : D \rightarrow D$ be a continuous map for which

- $G(a_0) = a = G(a_1)$,
- $G(b) = \{P_0\}$,
- $G(c_0) = b = G(c_1)$,
- G is injective on $D_0 \setminus b$ and D_1 ,
- $G(D_0 \setminus b) = D \setminus \{P_0\}$ and $G(D_1) = D$.

We are interested in periodic points of the map G . It is easy to see that periodic points lying on the arc a exist.

Problem to be solved



Let $G : D \rightarrow D$ be a continuous map for which

- $G(a_0) = a = G(a_1)$,
- $G(b) = \{P_0\}$,
- $G(c_0) = b = G(c_1)$,
- G is injective on $D_0 \setminus b$ and D_1 ,
- $G(D_0 \setminus b) = D \setminus \{P_0\}$ and $G(D_1) = D$.

We are interested in periodic points of the map G . It is easy to see that periodic points lying on the arc a exist.

Itinerary

For a periodic point P of the map G we consider its itinerary W as a sequence $(w_i)_{i=0}^{\infty}$ defined by

$$w_i = \begin{cases} 0 & \text{if } G^i(P) \in D_0, \\ 1 & \text{if } G^i(P) \in D_1. \end{cases}$$


For any periodic itinerary W there is a periodic point $P \in a$ of the map G with the itinerary W .

Itinerary

For a periodic point P of the map G we consider its itinerary W as a sequence $(w_i)_{i=0}^{\infty}$ defined by

$$w_i = \begin{cases} 0 & \text{if } G^i(P) \in D_0, \\ 1 & \text{if } G^i(P) \in D_1. \end{cases}$$

For any periodic itinerary W there is a periodic point $P \in a$ of the map G with the itinerary W . Such a point need not be unique.




Itinerary

For a periodic point P of the map G we consider its itinerary W as a sequence $(w_i)_{i=0}^{\infty}$ defined by

$$w_i = \begin{cases} 0 & \text{if } G^i(P) \in D_0, \\ 1 & \text{if } G^i(P) \in D_1. \end{cases}$$

For any periodic itinerary W there is a periodic point $P \in a$ of the map G with the itinerary W . Such a point need not be unique.



Saddle point

Let $U \subset \mathbb{R}^2$ be open and $P = (x_0, y_0) \in U$ be a fixed point of the map $G : U \rightarrow \mathbb{R}^2$, $(x, y) \mapsto (g_1(x, y), g_2(x, y))$. P is said to be a saddle fixed point of the map F if the Jacobi matrix

$$\begin{pmatrix} \frac{\partial g_1}{\partial x}(x_0, y_0) & \frac{\partial g_1}{\partial y}(x_0, y_0) \\ \frac{\partial g_2}{\partial x}(x_0, y_0) & \frac{\partial g_2}{\partial y}(x_0, y_0) \end{pmatrix}$$

Saddle point

Let $U \subset \mathbb{R}^2$ be open and $P = (x_0, y_0) \in U$ be a fixed point of the map $G : U \rightarrow \mathbb{R}^2$, $(x, y) \mapsto (g_1(x, y), g_2(x, y))$. P is said to be a saddle fixed point of the map F if the Jacobi matrix

$$\begin{pmatrix} \frac{\partial g_1}{\partial x}(x_0, y_0) & \frac{\partial g_1}{\partial y}(x_0, y_0) \\ \frac{\partial g_2}{\partial x}(x_0, y_0) & \frac{\partial g_2}{\partial y}(x_0, y_0) \end{pmatrix}$$

has eigenvalues $\lambda_{1,2}$

Saddle point

Let $U \subset \mathbb{R}^2$ be open and $P = (x_0, y_0) \in U$ be a fixed point of the map $G : U \rightarrow \mathbb{R}^2$, $(x, y) \mapsto (g_1(x, y), g_2(x, y))$. P is said to be a saddle fixed point of the map F if the Jacobi matrix

$$\begin{pmatrix} \frac{\partial g_1}{\partial x}(x_0, y_0) & \frac{\partial g_1}{\partial y}(x_0, y_0) \\ \frac{\partial g_2}{\partial x}(x_0, y_0) & \frac{\partial g_2}{\partial y}(x_0, y_0) \end{pmatrix}$$

has eigenvalues $\lambda_{1,2}$ with

$$|\lambda_1| < 1 < |\lambda_2|.$$

Saddle point


Let $U \subset \mathbb{R}^2$ be open and $P = (x_0, y_0) \in U$ be a fixed point of the map $G : U \rightarrow \mathbb{R}^2$, $(x, y) \mapsto (g_1(x, y), g_2(x, y))$. P is said to be a saddle fixed point of the map F if the Jacobi matrix

$$\begin{pmatrix} \frac{\partial g_1}{\partial x}(x_0, y_0) & \frac{\partial g_1}{\partial y}(x_0, y_0) \\ \frac{\partial g_2}{\partial x}(x_0, y_0) & \frac{\partial g_2}{\partial y}(x_0, y_0) \end{pmatrix}$$

has eigenvalues $\lambda_{1,2}$ with


$$|\lambda_1| < 1 < |\lambda_2| .$$

Saddle point




Now assume that $A = U \cup a$, where U is an open subset and $a \subset \text{Bd}(U)$ is an arc, $G : A \rightarrow \mathbb{R}^2$ is a continuous map, $P = (x_0, y_0) \in a$ is a fixed point of G ,

Saddle point




Now assume that $A = U \cup a$, where U is an open subset and $a \subset \text{Bd}(U)$ is an arc, $G : A \rightarrow \mathbb{R}^2$ is a continuous map, $P = (x_0, y_0) \in a$ is a fixed point of G , $G(U \cap V) \subset U$,

Saddle point




Now assume that $A = U \cup a$, where U is an open subset and $a \subset \text{Bd}(U)$ is an arc, $G : A \rightarrow \mathbb{R}^2$ is a continuous map, $P = (x_0, y_0) \in a$ is a fixed point of G , $G(U \cap V) \subset U$, $G(a \cap V) \subset a$ and for some neighbourhood V of P .

Saddle point



Now assume that $A = U \cup a$, where U is an open subset and $a \subset \text{Bd}(U)$ is an arc, $G : A \rightarrow \mathbb{R}^2$ is a continuous map, $P = (x_0, y_0) \in a$ is a fixed point of G , $G(U \cap V) \subset U$, $G(a \cap V) \subset a$ and for some neighbourhood V of P . We want to define the notion of saddle fixed point.

Saddle point



Now assume that $A = U \cup a$, where U is an open subset and $a \subset \text{Bd}(U)$ is an arc, $G : A \rightarrow \mathbb{R}^2$ is a continuous map, $P = (x_0, y_0) \in a$ is a fixed point of G , $G(U \cap V) \subset U$, $G(a \cap V) \subset a$ and for some neighbourhood V of P . We want to define the notion of saddle fixed point.

Saddle point

P is said to be a saddle fixed point of G if there exist

- a neighbourhood W of P ,

Saddle point

P is said to be a saddle fixed point of G if there exist

- a neighbourhood W of P ,
- $\delta > 0$,

Saddle point

P is said to be a saddle fixed point of G if there exist

- a neighbourhood W of P ,
- $\delta > 0$,
- a homeomorphism $H : A \cap W \rightarrow (-\delta, \delta) \times \langle 0, \delta \rangle$ and

Saddle point

P is said to be a saddle fixed point of G if there exist

- a neighbourhood W of P ,
- $\delta > 0$,
- a homeomorphism $H : A \cap W \rightarrow (-\delta, \delta) \times \langle 0, \delta \rangle$ and
- a map $\tilde{G} : (-\delta, \delta) \times \langle 0, \delta \rangle \rightarrow \mathbb{R}^2$, $(x, y) \mapsto (\tilde{g}_1(x, y), \tilde{g}_2(x, y))$

Saddle point

P is said to be a saddle fixed point of G if there exist

- a neighbourhood W of P ,
- $\delta > 0$,
- a homeomorphism $H : A \cap W \rightarrow (-\delta, \delta) \times \langle 0, \delta \rangle$ and
- a map $\tilde{G} : (-\delta, \delta) \times \langle 0, \delta \rangle \rightarrow \mathbb{R}^2$, $(x, y) \mapsto (\tilde{g}_1(x, y), \tilde{g}_2(x, y))$

Saddle point

P is said to be a saddle fixed point of G if there exist

- a neighbourhood W of P ,
- $\delta > 0$,
- a homeomorphism $H : A \cap W \rightarrow (-\delta, \delta) \times \langle 0, \delta \rangle$ and
- a map $\tilde{G} : (-\delta, \delta) \times \langle 0, \delta \rangle \rightarrow \mathbb{R}^2$, $(x, y) \mapsto (\tilde{g}_1(x, y), \tilde{g}_2(x, y))$

such that

- $H(P) = (0, 0)$,

Saddle point

P is said to be a saddle fixed point of G if there exist

- a neighbourhood W of P ,
- $\delta > 0$,
- a homeomorphism $H : A \cap W \rightarrow (-\delta, \delta) \times \langle 0, \delta \rangle$ and
- a map $\tilde{G} : (-\delta, \delta) \times \langle 0, \delta \rangle \rightarrow \mathbb{R}^2$, $(x, y) \mapsto (\tilde{g}_1(x, y), \tilde{g}_2(x, y))$

such that

- $H(P) = (0, 0)$,
- $\tilde{G}(H(Q)) = H(G(Q))$ for all $Q \in A \cap W \cap G^{-1}(A \cap W)$,

Saddle point

P is said to be a saddle fixed point of G if there exist

- a neighbourhood W of P ,
- $\delta > 0$,
- a homeomorphism $H : A \cap W \rightarrow (-\delta, \delta) \times \langle 0, \delta \rangle$ and
- a map $\tilde{G} : (-\delta, \delta) \times \langle 0, \delta \rangle \rightarrow \mathbb{R}^2$, $(x, y) \mapsto (\tilde{g}_1(x, y), \tilde{g}_2(x, y))$

such that

- $H(P) = (0, 0)$,
- $\tilde{G}(H(Q)) = H(G(Q))$ for all $Q \in A \cap W \cap G^{-1}(A \cap W)$,
- $\tilde{g}_2(x, 0) = 0$ for all $x \in (-\delta, \delta)$,

Saddle point

P is said to be a saddle fixed point of G if there exist

- a neighbourhood W of P ,
- $\delta > 0$,
- a homeomorphism $H : A \cap W \rightarrow (-\delta, \delta) \times \langle 0, \delta \rangle$ and
- a map $\tilde{G} : (-\delta, \delta) \times \langle 0, \delta \rangle \rightarrow \mathbb{R}^2$, $(x, y) \mapsto (\tilde{g}_1(x, y), \tilde{g}_2(x, y))$

such that

- $H(P) = (0, 0)$,
- $\tilde{G}(H(Q)) = H(G(Q))$ for all $Q \in A \cap W \cap G^{-1}(A \cap W)$,
- $\tilde{g}_2(x, 0) = 0$ for all $x \in (-\delta, \delta)$,
- $|\tilde{g}_1(x, 0)| > |x|$ for all $x \in (-\delta, \delta)$, $x \neq 0$,

Saddle point

P is said to be a saddle fixed point of G if there exist

- a neighbourhood W of P ,
- $\delta > 0$,
- a homeomorphism $H : A \cap W \rightarrow (-\delta, \delta) \times \langle 0, \delta \rangle$ and
- a map $\tilde{G} : (-\delta, \delta) \times \langle 0, \delta \rangle \rightarrow \mathbb{R}^2$, $(x, y) \mapsto (\tilde{g}_1(x, y), \tilde{g}_2(x, y))$

such that

- $H(P) = (0, 0)$,
- $\tilde{G}(H(Q)) = H(G(Q))$ for all $Q \in A \cap W \cap G^{-1}(A \cap W)$,
- $\tilde{g}_2(x, 0) = 0$ for all $x \in (-\delta, \delta)$,
- $|\tilde{g}_1(x, 0)| > |x|$ for all $x \in (-\delta, \delta)$, $x \neq 0$,
- $0 < \tilde{g}_2(x, y) < y$ for all $x \in (-\delta, \delta)$, $y \in (0, \delta)$.


Saddle point

P is said to be a saddle fixed point of G if there exist

- a neighbourhood W of P ,
- $\delta > 0$,
- a homeomorphism $H : A \cap W \rightarrow (-\delta, \delta) \times \langle 0, \delta \rangle$ and
- a map $\tilde{G} : (-\delta, \delta) \times \langle 0, \delta \rangle \rightarrow \mathbb{R}^2$, $(x, y) \mapsto (\tilde{g}_1(x, y), \tilde{g}_2(x, y))$


such that

- $H(P) = (0, 0)$,
- $\tilde{G}(H(Q)) = H(G(Q))$ for all $Q \in A \cap W \cap G^{-1}(A \cap W)$,
- $\tilde{g}_2(x, 0) = 0$ for all $x \in (-\delta, \delta)$,
- $|\tilde{g}_1(x, 0)| > |x|$ for all $x \in (-\delta, \delta)$, $x \neq 0$,
- $0 < \tilde{g}_2(x, y) < y$ for all $x \in (-\delta, \delta)$, $y \in (0, \delta)$.




Consider a partition $D = D_0 \cup D_1$ given by an arc d . Let $G : D \rightarrow D$ be a continuous map for which

- $G(a_0) = a = G(a_1)$,
- $G(b) = \{P_0\}$,




Consider a partition $D = D_0 \cup D_1$ given by an arc d . Let $G : D \rightarrow D$ be a continuous map for which

- $G(a_0) = a = G(a_1)$,
- $G(b) = \{P_0\}$,
- $G(c_0) = b = G(c_1)$,




Consider a partition $D = D_0 \cup D_1$ given by an arc d . Let $G : D \rightarrow D$ be a continuous map for which

- $G(a_0) = a = G(a_1)$,
- $G(b) = \{P_0\}$,
- $G(c_0) = b = G(c_1)$,
- G is injective on $D_0 \setminus b$ and D_1 ,



Consider a partition $D = D_0 \cup D_1$ given by an arc d . Let $G : D \rightarrow D$ be a continuous map for which

- $G(a_0) = a = G(a_1)$,
- $G(b) = \{P_0\}$,
- $G(c_0) = b = G(c_1)$,
- G is injective on $D_0 \setminus b$ and D_1 ,
- $G(D_0 \setminus b) = D \setminus \{P_0\}$ and $G(D_1) = D$.



Consider a partition $D = D_0 \cup D_1$ given by an arc d . Let $G : D \rightarrow D$ be a continuous map for which

- $G(a_0) = a = G(a_1)$,
- $G(b) = \{P_0\}$,
- $G(c_0) = b = G(c_1)$,
- G is injective on $D_0 \setminus b$ and D_1 ,
- $G(D_0 \setminus b) = D \setminus \{P_0\}$ and $G(D_1) = D$.

Main result

Theorem

Let $P \in a$, $P \neq P_0$ be a periodic point of the map G with period n such that

- P is a saddle fixed point of the map G^n ,*
- P is a unique periodic point on the arc a with its itinerary.*

Main result

Theorem

Let $P \in a$, $P \neq P_0$ be a periodic point of the map G with period n such that

- P is a saddle fixed point of the map G^n ,*
- P is a unique periodic point on the arc a with its itinerary.*

Main result

Theorem

Let $P \in a$, $P \neq P_0$ be a periodic point of the map G with period n such that

- P is a saddle fixed point of the map G^n ,*
- P is a unique periodic point on the arc a with its itinerary.*

Then there is a periodic point $Q \in \text{Int } D$ with the same period and itinerary.

Main result

Theorem

Let $P \in a$, $P \neq P_0$ be a periodic point of the map G with period n such that

- P is a saddle fixed point of the map G^n ,*
- P is a unique periodic point on the arc a with its itinerary.*

Then there is a periodic point $Q \in \text{Int } D$ with the same period and itinerary.

Jacobi matrix

Let $P = (x_0, 0) \in \Delta$ be a fixed point of the map F^n . In this case $P = \left(4 \sin^2 \frac{k\pi}{2^{n\pm 1}}, 0\right)$. Then the Jacobi matrix of the map F^n at the point P has a form

$$\begin{pmatrix} \lambda_1 & \mu \\ 0 & \lambda_2 \end{pmatrix} = \begin{pmatrix} \mp 2^n & \mu \\ 0 & \prod_{i=0}^{n-1} 4 \sin^2 \frac{2^i k\pi}{2^{n\pm 1}} \end{pmatrix}.$$

Jacobi matrix

Let $P = (x_0, 0) \in \Delta$ be a fixed point of the map F^n . In this case $P = \left(4 \sin^2 \frac{k\pi}{2^n \pm 1}, 0\right)$. Then the Jacobi matrix of the map F^n at the point P has a form

$$\begin{pmatrix} \lambda_1 & \mu \\ 0 & \lambda_2 \end{pmatrix} = \begin{pmatrix} \mp 2^n & \mu \\ 0 & \prod_{i=0}^{n-1} 4 \sin^2 \frac{2^i k\pi}{2^n \pm 1} \end{pmatrix}.$$

Classification

For λ_2 we have the possibilities

Saddle point

$$0 \leq \lambda_2 < 1, \text{ e.g. } x_0 = 4 \sin^2 \frac{\pi}{17}$$

Nonhyperbolic point

$$\lambda_2 = 1, \text{ e.g. } x_0 = 4 \sin^2 \frac{\pi}{15}$$

Repulsive point

$$1 < \lambda_2, \text{ e.g. } x_0 = 4 \sin^2 \frac{3\pi}{17}$$

Remark

All above points $(x_0, 0)$ have period 4.

Classification

Saddle point

Lower periodic points with period n and $0 < \lambda_2 < 1$ appear for all $n \geq 4$.

Nonhyperbolic point

Lower periodic points with period n and $\lambda_2 = 1$ appear for infinitely many n , e.g. $n = 4 \cdot 3^i \cdot 5^j$, where $i \geq 0, j \geq 0$.

Repulsive point

Lower periodic points with period n and $1 < \lambda_2$ appear for all $n \geq 1$.

Modifications

Assume that for any $x \in (0, 4)$ we have an increasing homeomorphism φ_x of the interval $\langle 0, 4 - x \rangle$ onto itself. Moreover let the function $\varphi(x, y) = \varphi_x(y)$ be continuous in the domain

$$\hat{\Delta} = \{ (x, y) : 0 < x < 4, 0 \leq y \leq 4 - x \} .$$

Modifications

Assume that for any $x \in (0, 4)$ we have an increasing homeomorphism φ_x of the interval $\langle 0, 4 - x \rangle$ onto itself. Moreover let the function $\varphi(x, y) = \varphi_x(y)$ be continuous in the domain

$$\hat{\Delta} = \{ (x, y) : 0 < x < 4, 0 \leq y \leq 4 - x \} .$$

Modifications

To obtain such above family φ_x , choose for $0 < x < 4$ a family of increasing homeomorphisms ψ_x of the interval $\langle 0, 1 \rangle$ such that the function $\psi(x, y) = \psi_x(y)$ is continuous in $(0, 4) \times \langle 0, 1 \rangle$ and put

$$\varphi_x(y) = (4 - x)\psi_x\left(\frac{y}{4 - x}\right).$$

Modifications

To obtain such above family φ_x , choose for $0 < x < 4$ a family of increasing homeomorphisms ψ_x of the interval $\langle 0, 1 \rangle$ such that the function $\psi(x, y) = \psi_x(y)$ is continuous in $(0, 4) \times \langle 0, 1 \rangle$ and put

$$\varphi_x(y) = (4 - x)\psi_x\left(\frac{y}{4 - x}\right).$$

Modifications

It is natural to put $\varphi_4(0)=0$. On the other hand we assume nothing about existence and properties of the limit

$$\lim_{x \rightarrow 0} \varphi_x(y).$$

Modifications

It is natural to put $\varphi_4(0)=0$. On the other hand we assume nothing about existence and properties of the limit

$$\lim_{x \rightarrow 0} \varphi_x(y).$$

Modifications

Let $G : \Delta \rightarrow \Delta$ be defined by

$$G(x, y) = \begin{cases} (0, 0) & \text{if } x = 0, \\ (x(4 - x - \varphi_x(y)), x\varphi_x(y)) & \text{otherwise} \end{cases} .$$

Then G is called a modified Lotka–Volterra map.

Modifications

Let $G : \Delta \rightarrow \Delta$ be defined by

$$G(x, y) = \begin{cases} (0, 0) & \text{if } x = 0, \\ (x(4 - x - \varphi_x(y)), x\varphi_x(y)) & \text{otherwise} \end{cases} .$$

Then G is called a modified Lotka–Volterra map.

Modifications

All such modifications have properties

- G is continuous on Δ

Modifications

All such modifications have properties

- G is continuous on Δ
- $G(\Delta_L) = \Delta = G(\Delta_R)$

Modifications

All such modifications have properties

- G is continuous on Δ
- $G(\Delta_L) = \Delta = G(\Delta_R)$
- G restricted to $\tilde{\Delta}_L$ and Δ_R is invertible

Modifications

All such modifications have properties

- G is continuous on Δ
- $G(\Delta_L) = \Delta = G(\Delta_R)$
- G restricted to $\tilde{\Delta}_L$ and Δ_R is invertible
- G restricted to the lower side is a logistic map.

Modifications

All such modifications have properties

- G is continuous on Δ
- $G(\Delta_L) = \Delta = G(\Delta_R)$
- G restricted to $\tilde{\Delta}_L$ and Δ_R is invertible
- G restricted to the lower side is a logistic map.

Formula for λ_2

We have

$$\tilde{\lambda}_2 = \prod_{i=0}^{n-1} x_i \varphi'_{x_i}(0) = \prod_{i=0}^{n-1} x_i \psi'_{x_i}(0) ,$$

or equivalently

$$\tilde{\lambda}_2 = \prod_{i=0}^{n-1} x_i \frac{\partial \varphi}{\partial y}(x_i, 0) = \prod_{i=0}^{n-1} x_i \frac{\partial \psi}{\partial y}(x_i, 0) ,$$

where $\varphi(x, y) = \varphi_x(y)$ and

Formula for λ_2

We have

$$\tilde{\lambda}_2 = \prod_{i=0}^{n-1} x_i \varphi'_{x_i}(0) = \prod_{i=0}^{n-1} x_i \psi'_{x_i}(0) ,$$

or equivalently

$$\tilde{\lambda}_2 = \prod_{i=0}^{n-1} x_i \frac{\partial \varphi}{\partial y}(x_i, 0) = \prod_{i=0}^{n-1} x_i \frac{\partial \psi}{\partial y}(x_i, 0) ,$$

where $\varphi(x, y) = \varphi_x(y)$ and $\psi(x, y) = \psi_x(y)$.

Formula for λ_2

We have

$$\tilde{\lambda}_2 = \prod_{i=0}^{n-1} x_i \varphi'_{x_i}(0) = \prod_{i=0}^{n-1} x_i \psi'_{x_i}(0) ,$$

or equivalently

$$\tilde{\lambda}_2 = \prod_{i=0}^{n-1} x_i \frac{\partial \varphi}{\partial y}(x_i, 0) = \prod_{i=0}^{n-1} x_i \frac{\partial \psi}{\partial y}(x_i, 0) ,$$

where $\varphi(x, y) = \varphi_x(y)$ and $\psi(x, y) = \psi_x(y)$.

(i) Let $0 \leq \alpha \leq 2$ and $\psi_x(y) = \alpha y + (1 - \alpha)y^2$. Then we obtain

$$\varphi_x(y) = \alpha y + \frac{(1-\alpha)y^2}{4-x}.$$

(ii) Let $\psi_x(y) = \frac{\sqrt{2y+x^2}-\sqrt{y+x^2}}{\sqrt{2+x^2}-\sqrt{1+x^2}}$. Then we obtain

$$\varphi_x(y) = \frac{\sqrt{2y(4-x) + x^2(4-x)^2} - \sqrt{y(4-x) + x^2(4-x)^2}}{\sqrt{2+x^2} - \sqrt{1+x^2}}.$$

(i) Let $0 \leq \alpha \leq 2$ and $\psi_x(y) = \alpha y + (1 - \alpha)y^2$. Then we obtain $\varphi_x(y) = \alpha y + \frac{(1-\alpha)y^2}{4-x}$.

(ii) Let $\psi_x(y) = \frac{\sqrt{2y+x^2}-\sqrt{y+x^2}}{\sqrt{2+x^2}-\sqrt{1+x^2}}$. Then we obtain

$$\varphi_x(y) = \frac{\sqrt{2y(4-x) + x^2(4-x)^2} - \sqrt{y(4-x) + x^2(4-x)^2}}{\sqrt{2+x^2} - \sqrt{1+x^2}}.$$

(iii) Let $\psi_x(y) = \sqrt{y}$. Then $\varphi_x(y) = \sqrt{(4-x)y}$.

(i) Let $0 \leq \alpha \leq 2$ and $\psi_x(y) = \alpha y + (1 - \alpha)y^2$. Then we obtain $\varphi_x(y) = \alpha y + \frac{(1-\alpha)y^2}{4-x}$.

(ii) Let $\psi_x(y) = \frac{\sqrt{2y+x^2}-\sqrt{y+x^2}}{\sqrt{2+x^2}-\sqrt{1+x^2}}$. Then we obtain

$$\varphi_x(y) = \frac{\sqrt{2y(4-x) + x^2(4-x)^2} - \sqrt{y(4-x) + x^2(4-x)^2}}{\sqrt{2+x^2} - \sqrt{1+x^2}}.$$

(iii) Let $\psi_x(y) = \sqrt{y}$. Then $\varphi_x(y) = \sqrt{(4-x)y}$.

(iv) Let $\psi_x(y) = y^x$. Then $\varphi_x(y) = (4-x) \left(\frac{y}{4-x} \right)^x$.

(i) Let $0 \leq \alpha \leq 2$ and $\psi_x(y) = \alpha y + (1 - \alpha)y^2$. Then we obtain $\varphi_x(y) = \alpha y + \frac{(1-\alpha)y^2}{4-x}$.

(ii) Let $\psi_x(y) = \frac{\sqrt{2y+x^2}-\sqrt{y+x^2}}{\sqrt{2+x^2}-\sqrt{1+x^2}}$. Then we obtain

$$\varphi_x(y) = \frac{\sqrt{2y(4-x) + x^2(4-x)^2} - \sqrt{y(4-x) + x^2(4-x)^2}}{\sqrt{2+x^2} - \sqrt{1+x^2}}.$$

(iii) Let $\psi_x(y) = \sqrt{y}$. Then $\varphi_x(y) = \sqrt{(4-x)y}$.

(iv) Let $\psi_x(y) = y^x$. Then $\varphi_x(y) = (4-x) \left(\frac{y}{4-x} \right)^x$.

Modification (i)

Let $0 \leq \alpha \leq 2$ and $G : \Delta \rightarrow \Delta$ be defined by

$$G(x, y) = \begin{cases} (0, 0) & \text{if } x = 4, \\ \left(x \left(4 - x - \alpha y - \frac{(1-\alpha)y^2}{4-x} \right), x \left(\alpha y + \frac{(1-\alpha)y^2}{4-x} \right) \right) & \text{otherwise} \end{cases}$$

In this case

$$\tilde{\lambda}_2 = \alpha^n \lambda_2 .$$

Modification (i)

Let $0 \leq \alpha \leq 2$ and $G : \Delta \rightarrow \Delta$ be defined by

$$G(x, y) = \begin{cases} (0, 0) & \text{if } x = 4, \\ \left(x \left(4 - x - \alpha y - \frac{(1-\alpha)y^2}{4-x} \right), x \left(\alpha y + \frac{(1-\alpha)y^2}{4-x} \right) \right) & \text{otherwise} \end{cases}$$

In this case

$$\tilde{\lambda}_2 = \alpha^n \lambda_2 .$$

- If $0 \leq \alpha < 1/3$ then G has interior periodic points for all periodic itineraries, because all lower fixed points of G^n are saddle fixed points.

Modification (i)

Let $0 \leq \alpha \leq 2$ and $G : \Delta \rightarrow \Delta$ be defined by

$$G(x, y) = \begin{cases} (0, 0) & \text{if } x = 4, \\ \left(x \left(4 - x - \alpha y - \frac{(1-\alpha)y^2}{4-x} \right), x \left(\alpha y + \frac{(1-\alpha)y^2}{4-x} \right) \right) & \text{otherwise} \end{cases}$$

In this case

$$\tilde{\lambda}_2 = \alpha^n \lambda_2 .$$

- If $0 \leq \alpha < 1/3$ then G has interior periodic points for all periodic itineraries, because all lower fixed points of G^n are saddle fixed points.
- If $\alpha = 1/3$ then $(3, 0)$ is not lower saddle fixed point of G and there is no interior fixed point of G lying in Δ_R . The other lower fixed points of G^n are saddle points for any $n > 1$.

Modification (i)

Let $0 \leq \alpha \leq 2$ and $G : \Delta \rightarrow \Delta$ be defined by

$$G(x, y) = \begin{cases} (0, 0) & \text{if } x = 4, \\ \left(x \left(4 - x - \alpha y - \frac{(1-\alpha)y^2}{4-x} \right), x \left(\alpha y + \frac{(1-\alpha)y^2}{4-x} \right) \right) & \text{otherwise} \end{cases}$$

In this case

$$\tilde{\lambda}_2 = \alpha^n \lambda_2 .$$

- If $0 \leq \alpha < 1/3$ then G has interior periodic points for all periodic itineraries, because all lower fixed points of G^n are saddle fixed points.
- If $\alpha = 1/3$ then $(3, 0)$ is not lower saddle fixed point of G and there is no interior fixed point of G lying in Δ_R . The other lower fixed points of G^n are saddle points for any $n > 1$.

Modification (i)

- If $0 \leq \alpha < \sqrt[4]{1 + 4/\sqrt{17}} \doteq 1.1847437 \dots$ then $P = (4 \sin^2 \frac{\pi}{17}, 0)$ is a saddle fixed point of G^4 .
- If $\sqrt[4]{1 + 4/\sqrt{17}} \leq \alpha \leq 2$ then $P = (4 \sin^2 \frac{\pi}{17}, 0)$ is not a saddle fixed point of G^4 .

Modification (i)

- If $0 \leq \alpha < \sqrt[4]{1 + 4/\sqrt{17}} \doteq 1.1847437 \dots$ then $P = (4 \sin^2 \frac{\pi}{17}, 0)$ is a saddle fixed point of G^4 .
- If $\sqrt[4]{1 + 4/\sqrt{17}} \leq \alpha \leq 2$ then $P = (4 \sin^2 \frac{\pi}{17}, 0)$ is not a saddle fixed point of G^4 .

Modifications (ii) and (iii)

Let

$$G(x, y) = (x(4 - x - \varphi_x(y)), x\varphi_x(y)) ,$$

where

$$\varphi_x(y) = \frac{\sqrt{2y(4-x) + x^2(4-x)^2} - \sqrt{y(4-x) + x^2(4-x)^2}}{\sqrt{2+x^2} - \sqrt{1+x^2}}$$

Modifications (ii) and (iii)

Let

$$G(x, y) = (x(4 - x - \varphi_x(y)), x\varphi_x(y)) ,$$

where

$$\varphi_x(y) = \frac{\sqrt{2y(4-x) + x^2(4-x)^2} - \sqrt{y(4-x) + x^2(4-x)^2}}{\sqrt{2+x^2} - \sqrt{1+x^2}}$$

or

$$\varphi_x(y) = \sqrt{(4-x)y} .$$

Modifications (ii) and (iii)

Let

$$G(x, y) = (x(4 - x - \varphi_x(y)), x\varphi_x(y)) ,$$

where

$$\varphi_x(y) = \frac{\sqrt{2y(4-x) + x^2(4-x)^2} - \sqrt{y(4-x) + x^2(4-x)^2}}{\sqrt{2+x^2} - \sqrt{1+x^2}}$$

or

$$\varphi_x(y) = \sqrt{(4-x)y} .$$

Then all lower fixed points different from $(0,0)$ of the map G^n are repulsive.

Modifications (ii) and (iii)

Let

$$G(x, y) = (x(4 - x - \varphi_x(y)), x\varphi_x(y)) ,$$

where

$$\varphi_x(y) = \frac{\sqrt{2y(4-x) + x^2(4-x)^2} - \sqrt{y(4-x) + x^2(4-x)^2}}{\sqrt{2+x^2} - \sqrt{1+x^2}}$$

or

$$\varphi_x(y) = \sqrt{(4-x)y} .$$

Then all lower fixed points different from $(0,0)$ of the map G^n are repulsive. In the case (ii)

$$\tilde{\lambda}_2 > \left(\frac{\sqrt{2}+1}{2} \right)^n > 1 .$$

Modifications (ii) and (iii)

Let

$$G(x, y) = (x(4 - x - \varphi_x(y)), x\varphi_x(y)) ,$$

where

$$\varphi_x(y) = \frac{\sqrt{2y(4-x) + x^2(4-x)^2} - \sqrt{y(4-x) + x^2(4-x)^2}}{\sqrt{2+x^2} - \sqrt{1+x^2}}$$

or

$$\varphi_x(y) = \sqrt{(4-x)y} .$$

Then all lower fixed points different from $(0,0)$ of the map G^n are repulsive. In the case (ii)

$$\tilde{\lambda}_2 > \left(\frac{\sqrt{2}+1}{2} \right)^n > 1 .$$

In the case (iii) the map G is not differentiable on the lower side.

Modifications (ii) and (iii)

Let

$$G(x, y) = (x(4 - x - \varphi_x(y)), x\varphi_x(y)) ,$$

where

$$\varphi_x(y) = \frac{\sqrt{2y(4-x) + x^2(4-x)^2} - \sqrt{y(4-x) + x^2(4-x)^2}}{\sqrt{2+x^2} - \sqrt{1+x^2}}$$

or

$$\varphi_x(y) = \sqrt{(4-x)y} .$$

Then all lower fixed points different from $(0,0)$ of the map G^n are repulsive. In the case (ii)

$$\tilde{\lambda}_2 > \left(\frac{\sqrt{2}+1}{2} \right)^n > 1 .$$

In the case (iii) the map G is not differentiable on the lower side.

Modification (iv)

Let

$$G(x, y) = \begin{cases} (0, 0) & \text{if } x \in \{0, 4\} \\ (x(4 - x - \varphi_x(y)), x\varphi_x(y)) & \text{otherwise,} \end{cases}$$

where

$$\varphi_x(y) = (4 - x) \left(\frac{y}{4 - x} \right)^x.$$

Let $P = (x_0, 0) \neq (0, 0)$ be a fixed point of the map F^n (and G^n as well). In this case $P = \left(4 \sin^2 \frac{k\pi}{2^{n\pm 1}}, 0 \right)$

Modification (iv)

Let

$$G(x, y) = \begin{cases} (0, 0) & \text{if } x \in \{0, 4\} \\ (x(4 - x - \varphi_x(y)), x\varphi_x(y)) & \text{otherwise,} \end{cases}$$

where

$$\varphi_x(y) = (4 - x) \left(\frac{y}{4 - x} \right)^x.$$

Let $P = (x_0, 0) \neq (0, 0)$ be a fixed point of the map F^n (and G^n as well). In this case $P = \left(4 \sin^2 \frac{k\pi}{2^{n\pm 1}}, 0 \right)$ and $\lambda_2 = \prod_{i=0}^{n-1} 4 \sin^2 \frac{2^i k\pi}{2^{n\pm 1}}$.

Modification (iv)

Let

$$G(x, y) = \begin{cases} (0, 0) & \text{if } x \in \{0, 4\} \\ (x(4 - x - \varphi_x(y)), x\varphi_x(y)) & \text{otherwise,} \end{cases}$$

where

$$\varphi_x(y) = (4 - x) \left(\frac{y}{4 - x} \right)^x.$$

Let $P = (x_0, 0) \neq (0, 0)$ be a fixed point of the map F^n (and G^n as well). In this case $P = \left(4 \sin^2 \frac{k\pi}{2^{n\pm 1}}, 0 \right)$ and $\lambda_2 = \prod_{i=0}^{n-1} 4 \sin^2 \frac{2^i k\pi}{2^{n\pm 1}}$. If $\lambda_2 > 1$ then P is a repulsive fixed point of F^n

Modification (iv)

Let

$$G(x, y) = \begin{cases} (0, 0) & \text{if } x \in \{0, 4\} \\ (x(4 - x - \varphi_x(y)), x\varphi_x(y)) & \text{otherwise,} \end{cases}$$

where

$$\varphi_x(y) = (4 - x) \left(\frac{y}{4 - x} \right)^x.$$

Let $P = (x_0, 0) \neq (0, 0)$ be a fixed point of the map F^n (and G^n as well). In this case $P = \left(4 \sin^2 \frac{k\pi}{2^{n\pm 1}}, 0 \right)$ and $\lambda_2 = \prod_{i=0}^{n-1} 4 \sin^2 \frac{2^i k\pi}{2^{n\pm 1}}$. If $\lambda_2 > 1$ then P is a repulsive fixed point of F^n and a saddle fixed point of G^n

Modification (iv)

Let

$$G(x, y) = \begin{cases} (0, 0) & \text{if } x \in \{0, 4\} \\ (x(4 - x - \varphi_x(y)), x\varphi_x(y)) & \text{otherwise,} \end{cases}$$

where

$$\varphi_x(y) = (4 - x) \left(\frac{y}{4 - x} \right)^x.$$

Let $P = (x_0, 0) \neq (0, 0)$ be a fixed point of the map F^n (and G^n as well). In this case $P = \left(4 \sin^2 \frac{k\pi}{2^{n\pm 1}}, 0 \right)$ and $\lambda_2 = \prod_{i=0}^{n-1} 4 \sin^2 \frac{2^i k\pi}{2^{n\pm 1}}$. If $\lambda_2 > 1$ then P is a repulsive fixed point of F^n and a saddle fixed point of G^n and there exists an interior fixed point of G^n with the same period and itinerary.

Modification (iv)

Let

$$G(x, y) = \begin{cases} (0, 0) & \text{if } x \in \{0, 4\} \\ (x(4 - x - \varphi_x(y)), x\varphi_x(y)) & \text{otherwise,} \end{cases}$$

where

$$\varphi_x(y) = (4 - x) \left(\frac{y}{4 - x} \right)^x.$$

Let $P = (x_0, 0) \neq (0, 0)$ be a fixed point of the map F^n (and G^n as well). In this case $P = \left(4 \sin^2 \frac{k\pi}{2^{n\pm 1}}, 0 \right)$ and $\lambda_2 = \prod_{i=0}^{n-1} 4 \sin^2 \frac{2^i k\pi}{2^{n\pm 1}}$. If $\lambda_2 > 1$ then P is a repulsive fixed point of F^n and a saddle fixed point of G^n and there exists an interior fixed point of G^n with the same period and itinerary. If $\lambda_2 < 1$ then P is a saddle fixed point of F^n

Modification (iv)

Let

$$G(x, y) = \begin{cases} (0, 0) & \text{if } x \in \{0, 4\} \\ (x(4 - x - \varphi_x(y)), x\varphi_x(y)) & \text{otherwise,} \end{cases}$$

where

$$\varphi_x(y) = (4 - x) \left(\frac{y}{4 - x} \right)^x.$$

Let $P = (x_0, 0) \neq (0, 0)$ be a fixed point of the map F^n (and G^n as well). In this case $P = \left(4 \sin^2 \frac{k\pi}{2^{n\pm 1}}, 0 \right)$ and $\lambda_2 = \prod_{i=0}^{n-1} 4 \sin^2 \frac{2^i k\pi}{2^{n\pm 1}}$. If $\lambda_2 > 1$ then P is a repulsive fixed point of F^n and a saddle fixed point of G^n and there exists an interior fixed point of G^n with the same period and itinerary. If $\lambda_2 < 1$ then P is a saddle fixed point of F^n and a repulsive fixed point of G^n .

Modification (iv)

Let

$$G(x, y) = \begin{cases} (0, 0) & \text{if } x \in \{0, 4\} \\ (x(4 - x - \varphi_x(y)), x\varphi_x(y)) & \text{otherwise,} \end{cases}$$

where

$$\varphi_x(y) = (4 - x) \left(\frac{y}{4 - x} \right)^x.$$

Let $P = (x_0, 0) \neq (0, 0)$ be a fixed point of the map F^n (and G^n as well). In this case $P = \left(4 \sin^2 \frac{k\pi}{2^{n\pm 1}}, 0 \right)$ and $\lambda_2 = \prod_{i=0}^{n-1} 4 \sin^2 \frac{2^i k \pi}{2^{n\pm 1}}$. If $\lambda_2 > 1$ then P is a repulsive fixed point of F^n and a saddle fixed point of G^n and there exists an interior fixed point of G^n with the same period and itinerary. If $\lambda_2 < 1$ then P is a saddle fixed point of F^n and a repulsive fixed point of G^n . What about $\lambda_2 = 1$?

Modification (iv)






Let

$$G(x, y) = \begin{cases} (0, 0) & \text{if } x \in \{0, 4\} \\ (x(4 - x - \varphi_x(y)), x\varphi_x(y)) & \text{otherwise,} \end{cases}$$

where

$$\varphi_x(y) = (4 - x) \left(\frac{y}{4 - x} \right)^x.$$

Let $P = (x_0, 0) \neq (0, 0)$ be a fixed point of the map F^n (and G^n as well). In this case $P = \left(4 \sin^2 \frac{k\pi}{2^{n\pm 1}}, 0 \right)$ and $\lambda_2 = \prod_{i=0}^{n-1} 4 \sin^2 \frac{2^i k\pi}{2^{n\pm 1}}$. If $\lambda_2 > 1$ then P is a repulsive fixed point of F^n and a saddle fixed point of G^n and there exists an interior fixed point of G^n with the same period and itinerary. If $\lambda_2 < 1$ then P is a saddle fixed point of F^n and a repulsive fixed point of G^n . What about $\lambda_2 = 1$?

-  F. Balibrea, J. L. G. Guirao, M. Lampart and J. Llibre. Dynamics of a Lotka–Volterra map. *Fund. Math.* **191** (2006), 265–279.
-  J. L. G. Guirao and M. Lampart. Transitivity of a Lotka–Volterra map. *Discrete Contin. Dyn. Syst. Ser. B* **9** (2008), no. 1, 75–82 (electronic).
-  P. Maličský. Interior periodic points of a Lotka–Volterra map. *J. Difference Equ. Appl.* **18** (2012), no. 4, 553–567.
-  A. N. Sharkovskii. Low dimensional dynamics. *Tagungsbericht 20/1993, Proceedings of Mathematisches Forschungsinstitut Oberwolfach* 1993, 17.
-  G. Swirszcz. On a certain map of the triangle. *Fund. Math.* **155** (1998), 45–57.