



Faculty of Mathematics
and Computer Science
University of Łódź



On \mathcal{I} -continuous functions

Jacek Hejduk & Anna Loranty & Renata Wiertelak

The 28th International Summer Conference
on Real Functions Theory
Stara Lesna, August 31-September 5, 2014

Table of contents

1 \mathcal{J} -density topologies

2 Continuous functions

Definition 1 (H. Lebesgue).

A point $x_0 \in \mathbb{R}$ is a **density point** of a Lebesgue measurable set A if

$$\lim_{h \rightarrow 0^+} \frac{\lambda(A \cap [x_0 - h, x_0 + h])}{2h} = 1.$$

Definition 1 (H. Lebesgue).

A point $x_0 \in \mathbb{R}$ is a **density point** of a Lebesgue measurable set A if

$$\lim_{h \rightarrow 0^+} \frac{\lambda(A \cap [x_0 - h, x_0 + h])}{2h} = 1.$$

$$\begin{aligned} \lim_{h \rightarrow 0^+} \frac{\lambda(A \cap [x_0 - h, x_0 + h])}{2h} = 1 &\Leftrightarrow \\ \lim_{n \rightarrow \infty} \frac{\lambda(A \cap [x_0 - \frac{1}{n}, x_0 + \frac{1}{n}])}{\frac{2}{n}} &= 1. \end{aligned}$$

A – a Lebesgue measurable set,

$$\Phi_d(A) = \{x \in \mathbb{R} : x \text{ is a density point of } A\}.$$

A – a Lebesgue measurable set,

$$\Phi_d(A) = \{x \in \mathbb{R} : x \text{ is a density point of } A\}.$$

Theorem 1 (Lebesgue, 1910).

$$\lambda(A \triangle \Phi_d(A)) = 0.$$

A – a Lebesgue measurable set,

$$\Phi_d(A) = \{x \in \mathbb{R} : x \text{ is a density point of } A\}.$$

Theorem 1 (Lebesgue, 1910).

$$\lambda(A \triangle \Phi_d(A)) = 0.$$

Theorem 2.

For any sets $A, B \in \mathcal{L}$ we have :

- ❶ $\Phi_d(\emptyset) = \emptyset, \quad \Phi_d(\mathbb{R}) = \mathbb{R};$
- ❷ $\lambda(A \triangle B) = 0 \Rightarrow \Phi_d(A) = \Phi_d(B);$
- ❸ $\lambda(A \triangle \Phi_d(A)) = 0;$
- ❹ $\Phi_d(A \cap B) = \Phi_d(A) \cap \Phi_d(B).$

Theorem 3 (O. Haupt, C. Pauc, 1952).

The family

$$\mathcal{T}_d = \{A \in \mathcal{L}: A \subset \Phi_d(A)\}$$

forms a topology on \mathbb{R} called the density topology.

Moreover, we have $\mathcal{T}_{nat} \subset \mathcal{T}_d$.

$\mathcal{J} = \{J_n\}_{n \in \mathbb{N}}$ is a sequence of intervals tending to zero if

$$\text{diam}\{J_n \cup \{0\}\} \xrightarrow[n \rightarrow \infty]{} 0.$$

$\mathcal{J} = \{J_n\}_{n \in \mathbb{N}}$ is a sequence of intervals tending to zero if

$$\text{diam}\{J_n \cup \{0\}\} \xrightarrow{n \rightarrow \infty} 0.$$

Definition 2.

We shall say that a point $x_0 \in \mathbb{R}$ is a **\mathcal{J} -density point** of a measurable set A , if

$$\lim_{n \rightarrow \infty} \frac{\lambda(A \cap (J_n + x_0))}{|J_n|} = 1.$$

A – a Lebesgue measurable set,

$$\Phi_{\mathcal{J}}(A) = \{x \in \mathbb{R} : x \text{ is a } \mathcal{J}\text{-density point of } A\}.$$

A – a Lebesgue measurable set,

$$\Phi_{\mathcal{J}}(A) = \{x \in \mathbb{R} : x \text{ is a } \mathcal{J}\text{-density point of } A\}.$$

Theorem 4.

For any sets $A, B \in \mathcal{L}$ we have:

- ❶ $\Phi_{\mathcal{J}}(\emptyset) = \emptyset, \quad \Phi_{\mathcal{J}}(\mathbb{R}) = \mathbb{R};$
- ❷ $\lambda(A \triangle B) = 0 \Rightarrow \Phi_{\mathcal{J}}(A) = \Phi_{\mathcal{J}}(B);$
- ❸ $\Phi_{\mathcal{J}}(A \cap B) = \Phi_{\mathcal{J}}(A) \cap \Phi_{\mathcal{J}}(B).$

The equivalent of Lebesgue Density Theorem does not hold. It was proved by M. Csörnyei in *Density theorems revisited*.

The equivalent of Lebesgue Density Theorem does not hold. It was proved by M. Csörnyei in *Density theorems revisited*.

$$\exists_{J \rightarrow 0} \exists_{A \in \mathcal{L}} \quad \lambda(A) > 0 \wedge \lambda(A \cap \Phi_{\mathcal{J}}(A)) = 0.$$

The equivalent of Lebesgue Density Theorem does not hold. It was proved by M. Csörnyei in *Density theorems revisited*.

$$\exists_{J \rightarrow 0} \exists_{A \in \mathcal{L}} \lambda(A) > 0 \wedge \lambda(A \cap \Phi_{\mathcal{J}}(A)) = 0.$$

Theorem 5.

If $A \in \mathcal{L}$ then $\Phi_{\mathcal{J}}(A) \in \mathcal{L}$.

The equivalent of Lebesgue Density Theorem does not hold. It was proved by M. Csörnyei in *Density theorems revisited*.

$$\exists_{J \rightarrow 0} \exists_{A \in \mathcal{L}} \lambda(A) > 0 \wedge \lambda(A \cap \Phi_{\mathcal{J}}(A)) = 0.$$

Theorem 5.

If $A \in \mathcal{L}$ then $\Phi_{\mathcal{J}}(A) \in \mathcal{L}$.

Theorem 6.

For $A \in \mathcal{L}$ we have

$$\lambda(\Phi_{\mathcal{J}}(A) \setminus A) = 0.$$

Theorem 7.

Let \mathcal{J} be a sequence of intervals tending to zero. Then

$$\mathcal{T}_{\mathcal{J}} = \{A \in \mathcal{L} : A \subset \Phi_{\mathcal{J}}(A)\}.$$

*is a topology on \mathbb{R} , which will be called \mathcal{J} -density topology.
Moreover, we have $\mathcal{T}_{nat} \subset \mathcal{T}_{\mathcal{J}}$.*

Theorem 8.

For every sequence \mathcal{J} of intervals tending to zero

Theorem 8.

For every sequence \mathcal{J} of intervals tending to zero

- (i) $\mathcal{T}_{nat} \subset \mathcal{T}_{\mathcal{J}}$ and the inclusion is proper. In particular $\mathcal{T}_{\mathcal{J}}$ is Hausdorff;*

Theorem 8.

For every sequence \mathcal{J} of intervals tending to zero

- (i) $\mathcal{T}_{nat} \subset \mathcal{T}_{\mathcal{J}}$ and the inclusion is proper. In particular $\mathcal{T}_{\mathcal{J}}$ is Hausdorff;*
- (ii) a subset C of \mathbb{R} is closed and discrete with respect to $\mathcal{T}_{\mathcal{J}}$ if, and only if, $\lambda(C) = 0$;*

Theorem 8.

For every sequence \mathcal{J} of intervals tending to zero

- (i) $\mathcal{T}_{nat} \subset \mathcal{T}_{\mathcal{J}}$ and the inclusion is proper. In particular $\mathcal{T}_{\mathcal{J}}$ is Hausdorff;*
- (ii) a subset C of \mathbb{R} is closed and discrete with respect to $\mathcal{T}_{\mathcal{J}}$ if, and only if, $\lambda(C) = 0$;*
- (iii) $(\mathbb{R}, \mathcal{T}_{\mathcal{J}})$ is neither separable nor has the Lindelöf property;*

Theorem 8.

For every sequence \mathcal{J} of intervals tending to zero

- (i) $\mathcal{T}_{nat} \subset \mathcal{T}_{\mathcal{J}}$ and the inclusion is proper. In particular $\mathcal{T}_{\mathcal{J}}$ is Hausdorff;*
- (ii) a subset C of \mathbb{R} is closed and discrete with respect to $\mathcal{T}_{\mathcal{J}}$ if, and only if, $\lambda(C) = 0$;*
- (iii) $(\mathbb{R}, \mathcal{T}_{\mathcal{J}})$ is neither separable nor has the Lindelöf property;*
- (iv) a set A is compact with respect to $\mathcal{T}_{\mathcal{J}}$ if, and only if, it is finite.*

Table of contents

1 \mathcal{J} -density topologies

2 Continuous functions

For $\mathcal{J} \in \mathfrak{S}$ we consider four families of continuous functions defined as follows:

$$\mathcal{C}_{nat,nat} = \{f: (\mathbb{R}, \mathcal{T}_{nat}) \rightarrow (\mathbb{R}, \mathcal{T}_{nat})\},$$

$$\mathcal{C}_{nat,\mathcal{J}} = \{f: (\mathbb{R}, \mathcal{T}_{nat}) \rightarrow (\mathbb{R}, \mathcal{T}_{\mathcal{J}})\},$$

$$\mathcal{C}_{\mathcal{J},nat} = \{f: (\mathbb{R}, \mathcal{T}_{\mathcal{J}}) \rightarrow (\mathbb{R}, \mathcal{T}_{nat})\},$$

$$\mathcal{C}_{\mathcal{J},\mathcal{J}} = \{f: (\mathbb{R}, \mathcal{T}_{\mathcal{J}}) \rightarrow (\mathbb{R}, \mathcal{T}_{\mathcal{J}})\}.$$

For $\mathcal{J} \in \mathfrak{S}$ we consider four families of continuous functions defined as follows:

$$\mathcal{C}_{nat,nat} = \{f: (\mathbb{R}, \mathcal{T}_{nat}) \rightarrow (\mathbb{R}, \mathcal{T}_{nat})\},$$

$$\mathcal{C}_{nat,\mathcal{J}} = \{f: (\mathbb{R}, \mathcal{T}_{nat}) \rightarrow (\mathbb{R}, \mathcal{T}_{\mathcal{J}})\},$$

$$\mathcal{C}_{\mathcal{J},nat} = \{f: (\mathbb{R}, \mathcal{T}_{\mathcal{J}}) \rightarrow (\mathbb{R}, \mathcal{T}_{nat})\},$$

$$\mathcal{C}_{\mathcal{J},\mathcal{J}} = \{f: (\mathbb{R}, \mathcal{T}_{\mathcal{J}}) \rightarrow (\mathbb{R}, \mathcal{T}_{\mathcal{J}})\}.$$

Property 1.

For $\mathcal{J} \in \mathfrak{S}$ the family $\mathcal{C}_{nat,\mathcal{J}}$ consists of constant functions.

For $\mathcal{J} \in \mathfrak{S}$ we consider four families of continuous functions defined as follows:

$$\mathcal{C}_{nat,nat} = \{f: (\mathbb{R}, \mathcal{T}_{nat}) \rightarrow (\mathbb{R}, \mathcal{T}_{nat})\},$$

$$\mathcal{C}_{nat,\mathcal{J}} = \{f: (\mathbb{R}, \mathcal{T}_{nat}) \rightarrow (\mathbb{R}, \mathcal{T}_{\mathcal{J}})\},$$

$$\mathcal{C}_{\mathcal{J},nat} = \{f: (\mathbb{R}, \mathcal{T}_{\mathcal{J}}) \rightarrow (\mathbb{R}, \mathcal{T}_{nat})\},$$

$$\mathcal{C}_{\mathcal{J},\mathcal{J}} = \{f: (\mathbb{R}, \mathcal{T}_{\mathcal{J}}) \rightarrow (\mathbb{R}, \mathcal{T}_{\mathcal{J}})\}.$$

Property 1.

For $\mathcal{J} \in \mathfrak{S}$ the family $\mathcal{C}_{nat,\mathcal{J}}$ consists of constant functions.

Let $f \in \mathcal{C}_{nat,\mathcal{J}}$ and $a, b \in \mathbb{R}$, $a < b$.

For $\mathcal{J} \in \mathfrak{S}$ we consider four families of continuous functions defined as follows:

$$\mathcal{C}_{nat,nat} = \{f: (\mathbb{R}, \mathcal{T}_{nat}) \rightarrow (\mathbb{R}, \mathcal{T}_{nat})\},$$

$$\mathcal{C}_{nat,\mathcal{J}} = \{f: (\mathbb{R}, \mathcal{T}_{nat}) \rightarrow (\mathbb{R}, \mathcal{T}_{\mathcal{J}})\},$$

$$\mathcal{C}_{\mathcal{J},nat} = \{f: (\mathbb{R}, \mathcal{T}_{\mathcal{J}}) \rightarrow (\mathbb{R}, \mathcal{T}_{nat})\},$$

$$\mathcal{C}_{\mathcal{J},\mathcal{J}} = \{f: (\mathbb{R}, \mathcal{T}_{\mathcal{J}}) \rightarrow (\mathbb{R}, \mathcal{T}_{\mathcal{J}})\}.$$

Property 1.

For $\mathcal{J} \in \mathfrak{S}$ the family $\mathcal{C}_{nat,\mathcal{J}}$ consists of constant functions.

Let $f \in \mathcal{C}_{nat,\mathcal{J}}$ and $a, b \in \mathbb{R}$, $a < b$.

$f([a, b])$ is $\mathcal{T}_{\mathcal{J}}$ -compact and $\mathcal{T}_{\mathcal{J}}$ -connected.

For $\mathcal{T} \in \mathfrak{S}$ we consider four families of continuous functions defined as follows:

$$\mathcal{C}_{nat,nat} = \{f: (\mathbb{R}, \mathcal{T}_{nat}) \rightarrow (\mathbb{R}, \mathcal{T}_{nat})\},$$

$$\mathcal{C}_{nat,\mathcal{T}} = \{f: (\mathbb{R}, \mathcal{T}_{nat}) \rightarrow (\mathbb{R}, \mathcal{T})\},$$

$$\mathcal{C}_{\mathcal{T},nat} = \{f: (\mathbb{R}, \mathcal{T}) \rightarrow (\mathbb{R}, \mathcal{T}_{nat})\},$$

$$\mathcal{C}_{\mathcal{T},\mathcal{T}} = \{f: (\mathbb{R}, \mathcal{T}) \rightarrow (\mathbb{R}, \mathcal{T})\}.$$

Property 1.

For $\mathcal{T} \in \mathfrak{S}$ the family $\mathcal{C}_{nat,\mathcal{T}}$ consists of constant functions.

Let $f \in \mathcal{C}_{nat,\mathcal{T}}$ and $a, b \in \mathbb{R}$, $a < b$.

$f([a, b])$ is $\mathcal{T}_{\mathcal{T}}$ -compact and $\mathcal{T}_{\mathcal{T}}$ -connected.

$f([a, b])$ is a singleton,

For $\mathcal{J} \in \mathfrak{S}$ we consider four families of continuous functions defined as follows:

$$\mathcal{C}_{nat,nat} = \{f: (\mathbb{R}, \mathcal{T}_{nat}) \rightarrow (\mathbb{R}, \mathcal{T}_{nat})\},$$

$$\mathcal{C}_{nat,\mathcal{J}} = \{f: (\mathbb{R}, \mathcal{T}_{nat}) \rightarrow (\mathbb{R}, \mathcal{T}_{\mathcal{J}})\},$$

$$\mathcal{C}_{\mathcal{J},nat} = \{f: (\mathbb{R}, \mathcal{T}_{\mathcal{J}}) \rightarrow (\mathbb{R}, \mathcal{T}_{nat})\},$$

$$\mathcal{C}_{\mathcal{J},\mathcal{J}} = \{f: (\mathbb{R}, \mathcal{T}_{\mathcal{J}}) \rightarrow (\mathbb{R}, \mathcal{T}_{\mathcal{J}})\}.$$

Property 1.

For $\mathcal{J} \in \mathfrak{S}$ the family $\mathcal{C}_{nat,\mathcal{J}}$ consists of constant functions.

Let $f \in \mathcal{C}_{nat,\mathcal{J}}$ and $a, b \in \mathbb{R}$, $a < b$.

$f([a, b])$ is $\mathcal{T}_{\mathcal{J}}$ -compact and $\mathcal{T}_{\mathcal{J}}$ -connected.

$f([a, b])$ is a singleton, $f(a) = f(b)$ and f is constant.

Property 2.

For $\mathcal{J} \in \mathfrak{S}$ the following inclusions holds:

$$\begin{aligned} \mathcal{C}_{nat, \mathcal{J}} \subsetneq \mathcal{C}_{nat, nat} &\subset \mathcal{C}_{\mathcal{J}, nat} \\ \mathcal{C}_{nat, \mathcal{J}} \subsetneq \mathcal{C}_{\mathcal{J}, \mathcal{J}} &\subset \mathcal{C}_{\mathcal{J}, nat} \end{aligned}$$

Property 2.

For $\mathcal{J} \in \mathfrak{S}$ the following inclusions holds:

$$\begin{aligned}\mathcal{C}_{nat,\mathcal{J}} &\subsetneq \mathcal{C}_{nat,nat} \subset \mathcal{C}_{\mathcal{J},nat} \\ \mathcal{C}_{nat,\mathcal{J}} &\subsetneq \mathcal{C}_{\mathcal{J},\mathcal{J}} \subset \mathcal{C}_{\mathcal{J},nat}\end{aligned}$$

$f(x) = x$ is the member of $\mathcal{C}_{nat,nat}$ and $\mathcal{C}_{\mathcal{J},\mathcal{J}}$ but not $\mathcal{C}_{nat,\mathcal{J}}$.

We say that a sequence of intervals $\mathcal{J} = \{[a_n, b_n]\}_{n \in \mathbb{N}} \in \mathfrak{S}$, is **right-side (left-side) tending to zero** if there exists $n_0 \in \mathbb{N}$ such that $b_n > 0$ ($a_n < 0$) for $n \geq n_0$ and

$$\lim_{n \rightarrow \infty} \frac{\min\{0, a_n\}}{b_n} = 0 \quad \left(\lim_{n \rightarrow \infty} \frac{\max\{0, b_n\}}{a_n} = 0 \right).$$

We say that a sequence of intervals $\mathcal{J} = \{[a_n, b_n]\}_{n \in \mathbb{N}} \in \mathfrak{S}$, is **right-side (left-side) tending to zero** if there exists $n_0 \in \mathbb{N}$ such that $b_n > 0$ ($a_n < 0$) for $n \geq n_0$ and

$$\lim_{n \rightarrow \infty} \frac{\min\{0, a_n\}}{b_n} = 0 \quad \left(\lim_{n \rightarrow \infty} \frac{\max\{0, b_n\}}{a_n} = 0 \right).$$

Sequence of intervals $\mathcal{J} \in \mathfrak{S}$ is **one-side tending to zero** if it is right-side or left-side tending to zero.

We say that a sequence of intervals $\mathcal{J} = \{[a_n, b_n]\}_{n \in \mathbb{N}} \in \mathfrak{S}$, is **right-side (left-side) tending to zero** if there exists $n_0 \in \mathbb{N}$ such that $b_n > 0$ ($a_n < 0$) for $n \geq n_0$ and

$$\lim_{n \rightarrow \infty} \frac{\min\{0, a_n\}}{b_n} = 0 \quad \left(\lim_{n \rightarrow \infty} \frac{\max\{0, b_n\}}{a_n} = 0 \right).$$

Sequence of intervals $\mathcal{J} \in \mathfrak{S}$ is **one-side tending to zero** if it is right-side or left-side tending to zero.

Theorem 9.

Let $\mathcal{J} \in \mathfrak{S}$. Then $[a, b] \in \mathcal{T}_{\mathcal{J}}$ ($(a, b] \in \mathcal{T}_{\mathcal{J}}$) for $a < b$ if and only if the sequence \mathcal{J} is right-side (left-side) tending to zero.

Theorem 10.

If \mathcal{J} is a sequence of intervals one-side tending to zero, then:

- (i) $\mathcal{C}_{nat,nat} \setminus \mathcal{C}_{\mathcal{J},\mathcal{J}} \neq \emptyset$,
- (ii) $\mathcal{C}_{\mathcal{J},\mathcal{J}} \setminus \mathcal{C}_{nat,nat} \neq \emptyset$.

Theorem 10.

If \mathcal{J} is a sequence of intervals one-side tending to zero, then:

- (i) $\mathcal{C}_{nat,nat} \setminus \mathcal{C}_{\mathcal{J},\mathcal{J}} \neq \emptyset$,
- (ii) $\mathcal{C}_{\mathcal{J},\mathcal{J}} \setminus \mathcal{C}_{nat,nat} \neq \emptyset$.


Theorem 11.

Let \mathcal{J} be a sequence of intervals one-side tending to zero. Then

- (i) $\mathcal{C}_{\mathcal{J},\mathcal{J}} \subsetneq \mathcal{C}_{\mathcal{J},nat}$,
- (ii) $\mathcal{C}_{nat,nat} \subsetneq \mathcal{C}_{\mathcal{J},nat}$.


Let \mathcal{J} be the sequence right side-tending to zero.

$$f(x) = -|x|, f \in \mathcal{C}_{nat,nat} \subset \mathcal{C}_{\mathcal{J},nat}.$$



Let \mathcal{J} be the sequence right side-tending to zero.


$$f(x) = -|x|, f \in \mathcal{C}_{nat,nat} \subset \mathcal{C}_{\mathcal{J},nat}.$$



$[-1, 1) \in \mathcal{T}_{\mathcal{J}}$ but $f^{-1}([-1, 1)) = [-1, 1] \notin \mathcal{T}_{\mathcal{J}}$. Thus $f \notin \mathcal{C}_{\mathcal{J},\mathcal{J}}$.

Let \mathcal{J} be the sequence right side-tending to zero.


$$f(x) = -|x|, f \in \mathcal{C}_{nat,nat} \subset \mathcal{C}_{\mathcal{J},nat}.$$



$[-1, 1) \in \mathcal{T}_{\mathcal{J}}$ but $f^{-1}([-1, 1)) = [-1, 1] \notin \mathcal{T}_{\mathcal{J}}$. Thus $f \notin \mathcal{C}_{\mathcal{J},\mathcal{J}}$.
 $f \in \mathcal{C}_{nat,nat} \setminus \mathcal{C}_{\mathcal{J},\mathcal{J}}$, $f \in \mathcal{C}_{\mathcal{J},nat} \setminus \mathcal{C}_{\mathcal{J},\mathcal{J}}$.


Let \mathcal{J} be the sequence right side-tending to zero.

$$h(x) = x - k \quad \text{for } x \in [k, k + 1), k \in \mathbb{Z}.$$



Let \mathcal{J} be the sequence right side-tending to zero.


$$h(x) = x - k \quad \text{for } x \in [k, k+1), k \in \mathbb{Z}.$$



If $A \in \mathcal{T}_{\mathcal{J}}$, then $h^{-1}(A) = \bigcup_{k \in \mathbb{Z}} ((A \cap [0, 1)) + k) \in \mathcal{T}_{\mathcal{J}}$.
Hence $h \in \mathcal{C}_{\mathcal{J}, \mathcal{J}} \subset \mathcal{C}_{\mathcal{J}, \text{nat}}$.

Let \mathcal{J} be the sequence right side-tending to zero.

$$h(x) = x - k \quad \text{for } x \in [k, k + 1), k \in \mathbb{Z}.$$




If $A \in \mathcal{T}_{\mathcal{J}}$, then $h^{-1}(A) = \bigcup_{k \in \mathbb{Z}} ((A \cap [0, 1)) + k) \in \mathcal{T}_{\mathcal{J}}$.

Hence $h \in \mathcal{C}_{\mathcal{J}, \mathcal{J}} \subset \mathcal{C}_{\mathcal{J}, \text{nat}}$.

Since $h^{-1}((-1, \frac{1}{2})) = \bigcup_{k \in \mathbb{Z}} [k, k + \frac{1}{2}) \notin \mathcal{T}_{\text{nat}}$, we have $h \notin \mathcal{C}_{\text{nat}, \text{nat}}$.

Let \mathcal{J} be the sequence right side-tending to zero.

$$h(x) = x - k \quad \text{for } x \in [k, k+1), k \in \mathbb{Z}.$$



If $A \in \mathcal{T}_{\mathcal{J}}$, then $h^{-1}(A) = \bigcup_{k \in \mathbb{Z}} ((A \cap [0, 1)) + k) \in \mathcal{T}_{\mathcal{J}}$.

Hence $h \in \mathcal{C}_{\mathcal{J}, \mathcal{J}} \subset \mathcal{C}_{\mathcal{J}, \text{nat}}$.

Since $h^{-1}((-1, \frac{1}{2})) = \bigcup_{k \in \mathbb{Z}} [k, k + \frac{1}{2}) \notin \mathcal{T}_{\text{nat}}$, we have

$h \notin \mathcal{C}_{\text{nat}, \text{nat}}$.

$h \in \mathcal{C}_{\mathcal{J}, \mathcal{J}} \setminus \mathcal{C}_{\text{nat}, \text{nat}}, h \in \mathcal{C}_{\mathcal{J}, \text{nat}} \setminus \mathcal{C}_{\text{nat}, \text{nat}}$.

Property 3.

*For $\mathcal{J} \in \mathfrak{S}$ be a sequence of intervals one-side tending to zero.
Then the following inclusions holds:*

$$\begin{aligned} \mathcal{C}_{nat, \mathcal{J}} &\subsetneq \mathcal{C}_{nat, nat} \subsetneq \mathcal{C}_{\mathcal{J}, nat} \\ \mathcal{C}_{nat, \mathcal{J}} &\subsetneq \mathcal{C}_{\mathcal{J}, \mathcal{J}} \subsetneq \mathcal{C}_{\mathcal{J}, nat} \end{aligned}$$

Theorem 12.

*If $\mathcal{J} \in \mathfrak{S}$, then there exists $\mathcal{K} \in \mathfrak{S}$ such that $\mathcal{T}_{\mathcal{J}} \neq \mathcal{T}_{\mathcal{K}}$,
 $\mathcal{C}_{\mathcal{J},\mathcal{J}} \neq \mathcal{C}_{\mathcal{K},\mathcal{K}}$ and $\mathcal{C}_{\mathcal{J},\text{nat}} \neq \mathcal{C}_{\mathcal{K},\text{nat}}$.*

Theorem 12.

If $\mathcal{J} \in \mathfrak{S}$, then there exists $\mathcal{K} \in \mathfrak{S}$ such that $\mathcal{T}_{\mathcal{J}} \neq \mathcal{T}_{\mathcal{K}}$, $\mathcal{C}_{\mathcal{J},\mathcal{J}} \neq \mathcal{C}_{\mathcal{K},\mathcal{K}}$ and $\mathcal{C}_{\mathcal{J},\text{nat}} \neq \mathcal{C}_{\mathcal{K},\text{nat}}$.

Problem 1.

$$\mathcal{T}_{\mathcal{J}} \neq \mathcal{T}_{\mathcal{K}} \iff \mathcal{C}_{\mathcal{J},\text{nat}} \neq \mathcal{C}_{\mathcal{K},\text{nat}}.$$

Theorem 12.

If $\mathcal{J} \in \mathfrak{S}$, then there exists $\mathcal{K} \in \mathfrak{S}$ such that $\mathcal{T}_{\mathcal{J}} \neq \mathcal{T}_{\mathcal{K}}$, $\mathcal{C}_{\mathcal{J},\mathcal{J}} \neq \mathcal{C}_{\mathcal{K},\mathcal{K}}$ and $\mathcal{C}_{\mathcal{J},\text{nat}} \neq \mathcal{C}_{\mathcal{K},\text{nat}}$.

Problem 1.

$$\mathcal{T}_{\mathcal{J}} \neq \mathcal{T}_{\mathcal{K}} \iff \mathcal{C}_{\mathcal{J},\text{nat}} \neq \mathcal{C}_{\mathcal{K},\text{nat}}.$$

Problem 2.

$$\mathcal{T}_{\mathcal{J}} \neq \mathcal{T}_{\mathcal{K}} \iff \mathcal{C}_{\mathcal{J},\mathcal{J}} \neq \mathcal{C}_{\mathcal{K},\mathcal{K}}.$$

Let $\mathcal{J}, \mathcal{K} \in \mathfrak{S}$. Then the sequence ordered in an arbitrary fashion containing all intervals of the sequences \mathcal{J} and \mathcal{K} , denoted by $\mathcal{J} \cup \mathcal{K}$, is called **the union of sequences** \mathcal{J} and \mathcal{K} .

Let $\mathcal{J}, \mathcal{K} \in \mathfrak{S}$. Then the sequence ordered in an arbitrary fashion containing all intervals of the sequences \mathcal{J} and \mathcal{K} , denoted by $\mathcal{J} \cup \mathcal{K}$, is called **the union of sequences** \mathcal{J} and \mathcal{K} .

Property 4.

If $\mathcal{J} \in \mathfrak{S}$ and $\mathcal{K} \in \mathfrak{S}$, then

$$\mathcal{T}_{\mathcal{J} \cup \mathcal{K}} = \mathcal{T}_{\mathcal{J}} \cap \mathcal{T}_{\mathcal{K}}.$$

Let $\mathcal{J}, \mathcal{K} \in \mathfrak{S}$. Then the sequence ordered in an arbitrary fashion containing all intervals of the sequences \mathcal{J} and \mathcal{K} , denoted by $\mathcal{J} \cup \mathcal{K}$, is called **the union of sequences** \mathcal{J} and \mathcal{K} .

Property 4.

If $\mathcal{J} \in \mathfrak{S}$ and $\mathcal{K} \in \mathfrak{S}$, then

$$\mathcal{T}_{\mathcal{J} \cup \mathcal{K}} = \mathcal{T}_{\mathcal{J}} \cap \mathcal{T}_{\mathcal{K}}.$$

$$\Phi_{\mathcal{J} \cup \mathcal{K}}(A) = \Phi_{\mathcal{J}}(A) \cap \Phi_{\mathcal{K}}(A).$$

Property 5.

Let $\mathcal{J} \in \mathfrak{S}$ and $\mathcal{K} \in \mathfrak{S}$. Then

- (i) $\mathcal{C}_{\mathcal{J},nat} \cap \mathcal{C}_{\mathcal{K},nat} = \mathcal{C}_{\mathcal{J} \cup \mathcal{K},nat},$
- (ii) $\mathcal{C}_{nat,\mathcal{J}} \cap \mathcal{C}_{nat,\mathcal{K}} = \mathcal{C}_{nat,\mathcal{J} \cup \mathcal{K}},$
- (iii) $\mathcal{C}_{\mathcal{J},\mathcal{J}} \cap \mathcal{C}_{\mathcal{K},\mathcal{K}} \subset \mathcal{C}_{\mathcal{J} \cup \mathcal{K},\mathcal{J} \cup \mathcal{K}}.$

Property 5.

Let $\mathcal{J} \in \mathfrak{S}$ and $\mathcal{K} \in \mathfrak{S}$. Then

- (i) $\mathcal{C}_{\mathcal{J},nat} \cap \mathcal{C}_{\mathcal{K},nat} = \mathcal{C}_{\mathcal{J} \cup \mathcal{K},nat}$,
- (ii) $\mathcal{C}_{nat,\mathcal{J}} \cap \mathcal{C}_{nat,\mathcal{K}} = \mathcal{C}_{nat,\mathcal{J} \cup \mathcal{K}}$,
- (iii) $\mathcal{C}_{\mathcal{J},\mathcal{J}} \cap \mathcal{C}_{\mathcal{K},\mathcal{K}} \subset \mathcal{C}_{\mathcal{J} \cup \mathcal{K},\mathcal{J} \cup \mathcal{K}}$.

Let $\mathcal{J} = \{[-\frac{1}{n}, 0]\}_{n \in \mathbb{N}}$, $\mathcal{K} = \{[0, \frac{1}{n}]\}_{n \in \mathbb{N}}$.
 $\mathcal{T}_{\mathcal{J} \cup \mathcal{K}}$ is the density topology.

Property 5.

Let $\mathcal{J} \in \mathfrak{S}$ and $\mathcal{K} \in \mathfrak{S}$. Then

- (i) $\mathcal{C}_{\mathcal{J},nat} \cap \mathcal{C}_{\mathcal{K},nat} = \mathcal{C}_{\mathcal{J} \cup \mathcal{K},nat}$,
- (ii) $\mathcal{C}_{nat,\mathcal{J}} \cap \mathcal{C}_{nat,\mathcal{K}} = \mathcal{C}_{nat,\mathcal{J} \cup \mathcal{K}}$,
- (iii) $\mathcal{C}_{\mathcal{J},\mathcal{J}} \cap \mathcal{C}_{\mathcal{K},\mathcal{K}} \subset \mathcal{C}_{\mathcal{J} \cup \mathcal{K},\mathcal{J} \cup \mathcal{K}}$.

Let $\mathcal{J} = \{[-\frac{1}{n}, 0]\}_{n \in \mathbb{N}}$, $\mathcal{K} = \{[0, \frac{1}{n}]\}_{n \in \mathbb{N}}$.

$\mathcal{T}_{\mathcal{J} \cup \mathcal{K}}$ is the density topology.

$f(x) = -x$ belongs to the family $\mathcal{C}_{\mathcal{J} \cup \mathcal{K},\mathcal{J} \cup \mathcal{K}}$.

Property 5.

Let $\mathcal{J} \in \mathfrak{S}$ and $\mathcal{K} \in \mathfrak{S}$. Then

- (i) $\mathcal{C}_{\mathcal{J},nat} \cap \mathcal{C}_{\mathcal{K},nat} = \mathcal{C}_{\mathcal{J} \cup \mathcal{K},nat}$,
- (ii) $\mathcal{C}_{nat,\mathcal{J}} \cap \mathcal{C}_{nat,\mathcal{K}} = \mathcal{C}_{nat,\mathcal{J} \cup \mathcal{K}}$,
- (iii) $\mathcal{C}_{\mathcal{J},\mathcal{J}} \cap \mathcal{C}_{\mathcal{K},\mathcal{K}} \subset \mathcal{C}_{\mathcal{J} \cup \mathcal{K},\mathcal{J} \cup \mathcal{K}}$.

Let $\mathcal{J} = \{[-\frac{1}{n}, 0]\}_{n \in \mathbb{N}}$, $\mathcal{K} = \{[0, \frac{1}{n}]\}_{n \in \mathbb{N}}$.

$\mathcal{T}_{\mathcal{J} \cup \mathcal{K}}$ is the density topology.

$f(x) = -x$ belongs to the family $\mathcal{C}_{\mathcal{J} \cup \mathcal{K},\mathcal{J} \cup \mathcal{K}}$.

$[0, 1) \in \mathcal{T}_{\mathcal{K}}$, whereas $f^{-1}([0, 1)) = (-1, 0] \notin \mathcal{T}_{\mathcal{K}}$.

Property 5.

Let $\mathcal{J} \in \mathfrak{S}$ and $\mathcal{K} \in \mathfrak{S}$. Then

- (i) $\mathcal{C}_{\mathcal{J},nat} \cap \mathcal{C}_{\mathcal{K},nat} = \mathcal{C}_{\mathcal{J} \cup \mathcal{K},nat}$,
- (ii) $\mathcal{C}_{nat,\mathcal{J}} \cap \mathcal{C}_{nat,\mathcal{K}} = \mathcal{C}_{nat,\mathcal{J} \cup \mathcal{K}}$,
- (iii) $\mathcal{C}_{\mathcal{J},\mathcal{J}} \cap \mathcal{C}_{\mathcal{K},\mathcal{K}} \subset \mathcal{C}_{\mathcal{J} \cup \mathcal{K},\mathcal{J} \cup \mathcal{K}}$.

Let $\mathcal{J} = \{[-\frac{1}{n}, 0]\}_{n \in \mathbb{N}}$, $\mathcal{K} = \{[0, \frac{1}{n}]\}_{n \in \mathbb{N}}$.

$\mathcal{T}_{\mathcal{J} \cup \mathcal{K}}$ is the density topology.

$f(x) = -x$ belongs to the family $\mathcal{C}_{\mathcal{J} \cup \mathcal{K},\mathcal{J} \cup \mathcal{K}}$.

$[0, 1) \in \mathcal{T}_{\mathcal{K}}$, whereas $f^{-1}([0, 1)) = (-1, 0] \notin \mathcal{T}_{\mathcal{K}}$.

It implies that $f \notin \mathcal{C}_{\mathcal{K},\mathcal{K}}$.

- [1] M. Csörnyei, *Density theorems revisited*, Acta Sci. Math. **64** (1998), 59-65.
- [2] M. Filipczak, T. Filipczak, J. Hejduk, *On the comparison of the density type topologies*, Atti Sem. Mat. Fis. Univ. Modena **LII** (2004), 37-46.
- [3] M. Filipczak, J. Hejduk, *On topologies associated with the Lebesgue measure*, Tatra Mt. Math. Publ. **28** (2004), 187-197.
- [4] J. Hejduk, R. Wiertelak, *On the generalization of density topologies on the real line*, to appear in Math. Slovaca at 2014.
- [5] W. Wilczyński, *Density topologies*, Handbook of Measure Theory, Ed. E. Pap. Elsevier, chapter 15(2002), 675-702.