

# Resolvability of abstract density topologies in $\mathbb{R}^n$ generated by lower or almost lower density operators

Grażyna Horbaczewska  
University of Łódź

28th International Summer Conference  
on Real Functions Theory  
2014

## Resolvability

### Definition (1943, E. Hewitt)

A topological space  $(X, \tau)$  is *resolvable* if it contains two disjoint dense subsets.

### Definition

Let  $\alpha$  be an arbitrary cardinal number greater than one. We say that a space  $(X, \tau)$  is  $\alpha$ -resolvable if there is a family of  $\alpha$ -many pairwise disjoint dense sets each of which intersects each nonempty open subsets of  $X$  in at least  $\alpha$  points.

$$\Delta(X, \tau) := \min\{\text{card}(A) : A \in \tau \wedge A \neq \emptyset\}$$

### Definition

$(X, \tau)$  is *maximally resolvable* if it is  $\Delta(X, \tau)$ -resolvable.

## Resolvability

### Definition (1943, E. Hewitt)

A topological space  $(X, \tau)$  is *resolvable* if it contains two disjoint dense subsets.

### Definition

Let  $\alpha$  be an arbitrary cardinal number greater than one. We say that a space  $(X, \tau)$  is  $\alpha$ -resolvable if there is a family of  $\alpha$ -many pairwise disjoint dense sets each of which intersects each nonempty open subsets of  $X$  in at least  $\alpha$  points.

$$\Delta(X, \tau) := \min\{\text{card}(A) : A \in \tau \wedge A \neq \emptyset\}$$

### Definition

$(X, \tau)$  is *maximally resolvable* if it is  $\Delta(X, \tau)$ -resolvable.

## Resolvability

### Definition (1943, E. Hewitt)

A topological space  $(X, \tau)$  is *resolvable* if it contains two disjoint dense subsets.

### Definition

Let  $\alpha$  be an arbitrary cardinal number greater than one. We say that a space  $(X, \tau)$  is  $\alpha$ -*resolvable* if there is a family of  $\alpha$ -many pairwise disjoint dense sets each of which intersects each nonempty open subsets of  $X$  in at least  $\alpha$  points.

$$\Delta(X, \tau) := \min\{\text{card}(A) : A \in \tau \wedge A \neq \emptyset\}$$

### Definition

$(X, \tau)$  is *maximally resolvable* if it is  $\Delta(X, \tau)$ -resolvable.

## Resolvability

### Definition (1943, E. Hewitt)

A topological space  $(X, \tau)$  is resolvable if it contains two disjoint dense subsets.

### Definition

Let  $\alpha$  be an arbitrary cardinal number greater than one. We say that a space  $(X, \tau)$  is  $\alpha$ -resolvable if there is a family of  $\alpha$ -many pairwise disjoint dense sets each of which intersects each nonempty open subsets of  $X$  in at least  $\alpha$  points.

$$\Delta(X, \tau) := \min\{\text{card}(A) : A \in \tau \wedge A \neq \emptyset\}$$

### Definition

$(X, \tau)$  is maximally resolvable if it is  $\Delta(X, \tau)$ -resolvable.

## Maximal resolvability

Theorem (W.W.Comfort, S. Garcia-Ferreira)

If a Hausdorff space without isolated points is locally compact or metrizable then it is maximally resolvable.

## Extraresolvability

## Definition

$(X, \tau)$  is extraresolvable if there exists a family  $\mathcal{M}$  of dense subsets of  $X$  such that

$$\text{card}(\mathcal{M}) > \Delta(X, \tau)$$

and for every  $C, D \in \mathcal{M}$ ,  $C \neq D$ , the set  $C \cap D$  is nowhere dense.

J. Hejduk, R. Wiertelak,

*On the abstract density topologies generated by lower and almost lower density operators*

Traditional and present-day topics in real analysis, Łódź University Press, 2013.

Let  $X$  be a nonempty set,  $\mathcal{S}$  be a  $\sigma$ -algebra of sets from  $X$  and  $\mathcal{I} \subset \mathcal{S}$  be a proper  $\sigma$ -ideal.

### Definition

We say that an operator  $\Phi : \mathcal{S} \rightarrow \mathcal{S}$  is a lower density operator on a measurable space  $(X, \mathcal{S}, \mathcal{I})$  if

- (i)  $\Phi(\emptyset) = \emptyset$ ,  $\Phi(X) = X$ ;
- (ii)  $\forall_{A, B \in \mathcal{S}} \Phi(A \cap B) = \Phi(A) \cap \Phi(B)$ ;
- (iii)  $\forall_{A, B \in \mathcal{S}} (A \triangle B \in \mathcal{I} \Rightarrow \Phi(A) = \Phi(B))$ ;
- (iv)  $\forall_{A \in \mathcal{S}} A \triangle \Phi(A) \in \mathcal{I}$ .



J. Hejduk, R. Wiertelak,

*On the abstract density topologies generated by lower and almost lower density operators*

Traditional and present-day topics in real analysis, Łódź University Press, 2013.

Let  $X$  be a nonempty set,  $\mathcal{S}$  be a  $\sigma$ -algebra of sets from  $X$  and  $\mathcal{I} \subset \mathcal{S}$  be a proper  $\sigma$ -ideal.

### Definition

We say that an operator  $\Phi : \mathcal{S} \rightarrow \mathcal{S}$  is a lower density operator on a measurable space  $(X, \mathcal{S}, \mathcal{I})$  if

- (i)  $\Phi(\emptyset) = \emptyset$ ,  $\Phi(X) = X$ ;
- (ii)  $\forall_{A, B \in \mathcal{S}} \Phi(A \cap B) = \Phi(A) \cap \Phi(B)$ ;
- (iii)  $\forall_{A, B \in \mathcal{S}} (A \triangle B \in \mathcal{I} \Rightarrow \Phi(A) = \Phi(B))$ ;
- (iv)  $\forall_{A \in \mathcal{S}} A \triangle \Phi(A) \in \mathcal{I}$ .

J. Hejduk, R. Wiertelak,

*On the abstract density topologies generated by lower and almost lower density operators*

Traditional and present-day topics in real analysis, Łódź University Press, 2013.

Let  $X$  be a nonempty set,  $\mathcal{S}$  be a  $\sigma$ -algebra of sets from  $X$  and  $\mathcal{I} \subset \mathcal{S}$  be a proper  $\sigma$ -ideal.

### Definition

We say that an operator  $\Phi : \mathcal{S} \rightarrow \mathcal{S}$  is a lower density operator on a measurable space  $(X, \mathcal{S}, \mathcal{I})$  if

- (i)  $\Phi(\emptyset) = \emptyset$ ,  $\Phi(X) = X$ ;
- (ii)  $\forall_{A, B \in \mathcal{S}} \Phi(A \cap B) = \Phi(A) \cap \Phi(B)$ ;
- (iii)  $\forall_{A, B \in \mathcal{S}} (A \triangle B \in \mathcal{I} \Rightarrow \Phi(A) = \Phi(B))$ ;
- (iv)  $\forall_{A \in \mathcal{S}} A \triangle \Phi(A) \in \mathcal{I}$ .

## Abstract density topologies

### Definition

We say that a topology  $\tau$  is an abstract density topology on  $X$  if there exists a lower density operator  $\Phi$  on  $(X, \mathcal{S}, \mathcal{I})$  such that

$$\tau = \mathcal{T}_\Phi,$$

where  $\mathcal{T}_\Phi := \{A \in \mathcal{S} : A \subset \Phi(A)\}$ .

Topology  $\mathcal{T}_\Phi$  is called generated by a lower density operator on  $(X, \mathcal{S}, \mathcal{I})$ .

### Theorem

Let  $\Phi$  be a lower density operator on  $(X, \mathcal{S}, \mathcal{I})$ .

Then the family  $\mathcal{T}_\Phi$  is a topology on  $X$  if and only if the pair  $(\mathcal{S}, \mathcal{I})$  has the hull property.

## Abstract density topologies

### Definition

We say that a topology  $\tau$  is an abstract density topology on  $X$  if there exists a lower density operator  $\Phi$  on  $(X, \mathcal{S}, \mathcal{I})$  such that

$$\tau = \mathcal{T}_\Phi,$$

where  $\mathcal{T}_\Phi := \{A \in \mathcal{S} : A \subset \Phi(A)\}$ .

Topology  $\mathcal{T}_\Phi$  is called generated by a lower density operator on  $(X, \mathcal{S}, \mathcal{I})$ .

### Theorem

Let  $\Phi$  be a lower density operator on  $(X, \mathcal{S}, \mathcal{I})$ .

Then the family  $\mathcal{T}_\Phi$  is a topology on  $X$  if and only if the pair  $(\mathcal{S}, \mathcal{I})$  has the hull property.

## Abstract density topologies generated by LD

- density topology - O.Haupt, C.Pauc (1952)
- $\mathcal{I}$ -density topology - W.Poreda, E.Wagner-Bojakowska, W.Wilczyński (1985)
- topology involving measure and category - W.Wojdowski (1989)
- $\langle s \rangle$ -density topology with respect to category - J.Hejduk, G.H. (2003)
- $\langle s \rangle$ -density topology with respect to measure - M.Filipczak, J.Hejduk (2004), G.H. (2011)
- $\langle s \rangle$ - simple density topology with respect to category - V.Aversa, W.Wilczyński (2004)
- density topology related to category with respect to a sequence tending to zero - R.Wiertelak (2006)
- $\Psi_{\mathcal{I}}$ - density topology - E.Łazarow, A.Vizvary (2010)
- category  $\psi$ -density topology - W.Wojdowski, W.Wilczyński (2011)

## Abstract density topologies generated by LD

- density topology - O.Haupt, C.Pauc (1952)
- $\mathcal{I}$ -density topology - W.Poreda, E.Wagner-Bojakowska, W.Wilczyński (1985)
- topology involving measure and category - W.Wojdowski (1989)
- $< s >$ -density topology with respect to category - J.Hejduk, G.H. (2003)
- $< s >$ -density topology with respect to measure - M.Filipczak, J.Hejduk (2004), G.H. (2011)
- $< s >$ - simple density topology with respect to category - V.Aversa, W.Wilczyński (2004)
- density topology related to category with respect to a sequence tending to zero - R.Wiertelak (2006)
- $\Psi_{\mathcal{I}}$ - density topology - E.Łazarow, A.Vizvary (2010)
- category  $\psi$ -density topology - W.Wojdowski, W.Wilczyński (2011)

## Abstract density topologies generated by LD

- density topology - O.Haupt, C.Pauc (1952)
- $\mathcal{I}$ -density topology - W.Poreda, E.Wagner-Bojakowska, W.Wilczyński (1985)
- topology involving measure and category - W.Wojdowski (1989)
- $< s >$ -density topology with respect to category - J.Hejduk, G.H. (2003)
- $< s >$ -density topology with respect to measure - M.Filipczak, J.Hejduk (2004), G.H. (2011)
- $< s >$ - simple density topology with respect to category - V.Aversa, W.Wilczyński (2004)
- density topology related to category with respect to a sequence tending to zero - R.Wiertelak (2006)
- $\Psi_{\mathcal{I}}$ - density topology - E.Łazarow, A.Vizvary (2010)
- category  $\psi$ -density topology - W.Wojdowski, W.Wilczyński (2011)

## Abstract density topologies generated by LD

- density topology - O.Haupt, C.Pauc (1952)
- $\mathcal{I}$ -density topology - W.Poreda, E.Wagner-Bojakowska, W.Wilczyński (1985)
- topology involving measure and category - W.Wojdowski (1989)
- $< s >$ -density topology with respect to category - J.Hejduk, G.H. (2003)
- $< s >$ -density topology with respect to measure - M.Filipczak, J.Hejduk (2004), G.H. (2011)
- $< s >$ - simple density topology with respect to category - V.Aversa, W.Wilczyński (2004)
- density topology related to category with respect to a sequence tending to zero - R.Wiertelak (2006)
- $\Psi_{\mathcal{I}}$ - density topology - E.Łazarow, A.Vizvary (2010)
- category  $\psi$ -density topology - W.Wojdowski, W.Wilczyński (2011)



## Abstract density topologies generated by LD

- density topology - O.Haupt, C.Pauc (1952)
- $\mathcal{I}$ -density topology - W.Poreda, E.Wagner-Bojakowska, W.Wilczyński (1985)
- topology involving measure and category - W.Wojdowski (1989)
- $< s >$ -density topology with respect to category - J.Hejduk, G.H. (2003)
- $< s >$ -density topology with respect to measure - M.Filipczak, J.Hejduk (2004), G.H. (2011)
- $< s >$ - simple density topology with respect to category - V.Aversa, W.Wilczyński (2004)
- density topology related to category with respect to a sequence tending to zero - R.Wiertelak (2006)
- $\Psi_{\mathcal{I}}$ - density topology - E.Łazarow, A.Vizvary (2010)
- category  $\psi$ -density topology - W.Wojdowski, W.Wilczyński (2011)

## Abstract density topologies generated by LD

- density topology - O.Haupt, C.Pauc (1952)
- $\mathcal{I}$ -density topology - W.Poreda, E.Wagner-Bojakowska, W.Wilczyński (1985)
- topology involving measure and category - W.Wojdowski (1989)
- $< s >$ -density topology with respect to category - J.Hejduk, G.H. (2003)
- $< s >$ -density topology with respect to measure - M.Filipczak, J.Hejduk (2004), G.H. (2011)
- $< s >$ - simple density topology with respect to category - V.Aversa, W.Wilczyński (2004)
- density topology related to category with respect to a sequence tending to zero - R.Wiertelak (2006)
- $\Psi_{\mathcal{I}}$ - density topology - E.Łazarow, A.Vizvary (2010)
- category  $\psi$ -density topology - W.Wojdowski, W.Wilczyński (2011)

## Abstract density topologies generated by LD

- density topology - O.Haupt, C.Pauc (1952)
- $\mathcal{I}$ -density topology - W.Poreda, E.Wagner-Bojakowska, W.Wilczyński (1985)
- topology involving measure and category - W.Wojdowski (1989)
- $\langle s \rangle$ -density topology with respect to category - J.Hejduk, G.H. (2003)
- $\langle s \rangle$ -density topology with respect to measure - M.Filipczak, J.Hejduk (2004), G.H. (2011)
- $\langle s \rangle$ - simple density topology with respect to category - V.Aversa, W.Wilczyński (2004)
- density topology related to category with respect to a sequence tending to zero - R.Wiertelak (2006)
- $\Psi_{\mathcal{I}}$ - density topology - E.Łazarow, A.Vizvary (2010)
- category  $\psi$ -density topology - W.Wojdowski, W.Wilczyński (2011)

## Abstract density topologies generated by LD

- density topology - O.Haupt, C.Pauc (1952)
- $\mathcal{I}$ -density topology - W.Poreda, E.Wagner-Bojakowska, W.Wilczyński (1985)
- topology involving measure and category - W.Wojdowski (1989)
- $< s >$ -density topology with respect to category - J.Hejduk, G.H. (2003)
- $< s >$ -density topology with respect to measure - M.Filipczak, J.Hejduk (2004), G.H. (2011)
- $< s >$ - simple density topology with respect to category - V.Aversa, W.Wilczyński (2004)
- density topology related to category with respect to a sequence tending to zero - R.Wiertelak (2006)
- $\Psi_{\mathcal{I}}$ - density topology - E.Łazarow, A.Vizvary (2010)
- category  $\psi$ -density topology - W.Wojdowski, W.Wilczyński (2011)

## Abstract density topologies generated by LD

- density topology - O.Haupt, C.Pauc (1952)
- $\mathcal{I}$ -density topology - W.Poreda, E.Wagner-Bojakowska, W.Wilczyński (1985)
- topology involving measure and category - W.Wojdowski (1989)
- $< s >$ -density topology with respect to category - J.Hejduk, G.H. (2003)
- $< s >$ -density topology with respect to measure - M.Filipczak, J.Hejduk (2004), G.H. (2011)
- $< s >$ - simple density topology with respect to category - V.Aversa, W.Wilczyński (2004)
- density topology related to category with respect to a sequence tending to zero - R.Wiertelak (2006)
- $\Psi_{\mathcal{I}}$ - density topology - E.Łazarow, A.Vizvary (2010)
- category  $\psi$ -density topology - W.Wojdowski, W.Wilczyński (2011)

## Abstract density topologies

### Definition

We say that an operator  $\Phi : \mathcal{S} \rightarrow 2^X$  is an almost lower density operator on a measurable space  $(X, \mathcal{S}, \mathcal{I})$  if

- (i)  $\Phi(\emptyset) = \emptyset$ ,  $\Phi(X) = X$ ;
- (ii)  $\forall_{A, B \in \mathcal{S}} \Phi(A \cap B) = \Phi(A) \cap \Phi(B)$ ;
- (iii)  $\forall_{A, B \in \mathcal{S}} (A \triangle B \in \mathcal{I} \Rightarrow \Phi(A) = \Phi(B))$ ;
- (iv-a)  $\forall_{A \in \mathcal{S}} \Phi(A) \setminus A \in \mathcal{I}$ .

### Theorem

Let  $\Phi$  be an almost lower density operator on  $(X, \mathcal{S}, \mathcal{I})$ .  
If the pair  $(\mathcal{S}, \mathcal{I})$  has the hull property, then the family  
 $\mathcal{T}_\Phi = \{A \in \mathcal{S} : A \subset \Phi(A)\}$  is a topology on  $X$ .

In this case we say that topology  $\mathcal{T}_\Phi$  is generated by the almost lower density operator on  $(X, \mathcal{S}, \mathcal{I})$ .

## Abstract density topologies

### Definition

We say that an operator  $\Phi : \mathcal{S} \rightarrow 2^X$  is an almost lower density operator on a measurable space  $(X, \mathcal{S}, \mathcal{I})$  if

- (i)  $\Phi(\emptyset) = \emptyset$ ,  $\Phi(X) = X$ ;
- (ii)  $\forall_{A, B \in \mathcal{S}} \Phi(A \cap B) = \Phi(A) \cap \Phi(B)$ ;
- (iii)  $\forall_{A, B \in \mathcal{S}} (A \triangle B \in \mathcal{I} \Rightarrow \Phi(A) = \Phi(B))$ ;
- (iv-a)  $\forall_{A \in \mathcal{S}} \Phi(A) \setminus A \in \mathcal{I}$ .

### Theorem

Let  $\Phi$  be an almost lower density operator on  $(X, \mathcal{S}, \mathcal{I})$ .  
If the pair  $(\mathcal{S}, \mathcal{I})$  has the hull property, then the family  $\mathcal{T}_\Phi = \{A \in \mathcal{S} : A \subset \Phi(A)\}$  is a topology on  $X$ .

In this case we say that topology  $\mathcal{T}_\Phi$  is generated by the almost lower density operator on  $(X, \mathcal{S}, \mathcal{I})$ .

## Abstract density topologies generated by ALD

- **$\psi$ -density topology - M.Terepeta, E.Wagner-Bojakowska (1999)**
- density topology with respect to the O'Malley points - W.Poreda, W.Wilczyński (2001)
- density topology with respect to measure and category - J.Hejduk (2002)
- simple density topology - V.Aversa, W.Wilczyński (2004)
- complete density topology - W.Wilczyński, W.Wojdowski (2007)
- topologies connected with Hausdorff measures - G.H. (2008)
- $f$ -density topology - M.Filipczak, T.Filipczak (2008)
- $f$ -symmetrical density topology - J.Hejduk (2008)
- topology in the aspect of measure with respect to a sequence tending to zero - J.Hejduk, R.Wiertelak (2012)



## Abstract density topologies generated by ALD

- $\psi$ -density topology - M.Terepeta, E.Wagner-Bojakowska (1999)
- density topology with respect to the O'Malley points - W.Poreda, W.Wilczyński (2001)
- density topology with respect to measure and category - J.Hejduk (2002)
- simple density topology - V.Aversa, W.Wilczyński (2004)
- complete density topology - W.Wilczyński, W.Wojdowski (2007)
- topologies connected with Hausdorff measures - G.H. (2008)
- $f$ -density topology - M.Filipczak, T.Filipczak (2008)
- $f$ -symmetrical density topology - J.Hejduk (2008)
- topology in the aspect of measure with respect to a sequence tending to zero - J.Hejduk, R.Wiertelak (2012)

## Abstract density topologies generated by ALD

- $\psi$ -density topology - M.Terepeta, E.Wagner-Bojakowska (1999)
- density topology with respect to the O'Malley points - W.Poreda, W.Wilczyński (2001)
- density topology with respect to measure and category - J.Hejduk (2002)
- simple density topology - V.Aversa, W.Wilczyński (2004)
- complete density topology - W.Wilczyński, W.Wojdowski (2007)
- topologies connected with Hausdorff measures - G.H. (2008)
- $f$ -density topology - M.Filipczak, T.Filipczak (2008)
- $f$ -symmetrical density topology - J.Hejduk (2008)
- topology in the aspect of measure with respect to a sequence tending to zero - J.Hejduk, R.Wiertelak (2012)

## Abstract density topologies generated by ALD

- $\psi$ -density topology - M.Terepeta, E.Wagner-Bojakowska (1999)
- density topology with respect to the O'Malley points - W.Poreda, W.Wilczyński (2001)
- density topology with respect to measure and category - J.Hejduk (2002)
- simple density topology - V.Aversa, W.Wilczyński (2004)
- complete density topology - W.Wilczyński, W.Wojdowski (2007)
- topologies connected with Hausdorff measures - G.H. (2008)
- $f$ -density topology - M.Filipczak, T.Filipczak (2008)
- $f$ -symmetrical density topology - J.Hejduk (2008)
- topology in the aspect of measure with respect to a sequence tending to zero - J.Hejduk, R.Wiertelak (2012)

## Abstract density topologies generated by ALD

- $\psi$ -density topology - M.Terepeta, E.Wagner-Bojakowska (1999)
- density topology with respect to the O'Malley points - W.Poreda, W.Wilczyński (2001)
- density topology with respect to measure and category - J.Hejduk (2002)
- simple density topology - V.Aversa, W.Wilczyński (2004)
- complete density topology - W.Wilczyński, W.Wojdowski (2007)
- topologies connected with Hausdorff measures - G.H. (2008)
- $f$ -density topology - M.Filipczak, T.Filipczak (2008)
- $f$ -symmetrical density topology - J.Hejduk (2008)
- topology in the aspect of measure with respect to a sequence tending to zero - J.Hejduk, R.Wiertelak (2012)

## Abstract density topologies generated by ALD

- $\psi$ -density topology - M.Terepeta, E.Wagner-Bojakowska (1999)
- density topology with respect to the O'Malley points - W.Poreda, W.Wilczyński (2001)
- density topology with respect to measure and category - J.Hejduk (2002)
- simple density topology - V.Aversa, W.Wilczyński (2004)
- complete density topology - W.Wilczyński, W.Wojdowski (2007)
- topologies connected with Hausdorff measures - G.H. (2008)
- $f$ -density topology - M.Filipczak, T.Filipczak (2008)
- $f$ -symmetrical density topology - J.Hejduk (2008)
- topology in the aspect of measure with respect to a sequence tending to zero - J.Hejduk, R.Wiertelak (2012)

## Abstract density topologies generated by ALD

- $\psi$ -density topology - M.Terepeta, E.Wagner-Bojakowska (1999)
- density topology with respect to the O'Malley points - W.Poreda, W.Wilczyński (2001)
- density topology with respect to measure and category - J.Hejduk (2002)
- simple density topology - V.Aversa, W.Wilczyński (2004)
- complete density topology - W.Wilczyński, W.Wojdowski (2007)
- topologies connected with Hausdorff measures - G.H. (2008)
- $f$ -density topology - M.Filipczak, T.Filipczak (2008)
- $f$ -symmetrical density topology - J.Hejduk (2008)
- topology in the aspect of measure with respect to a sequence tending to zero - J.Hejduk, R.Wiertelak (2012)

## Abstract density topologies generated by ALD

- $\psi$ -density topology - M.Terepeta, E.Wagner-Bojakowska (1999)
- density topology with respect to the O'Malley points - W.Poreda, W.Wilczyński (2001)
- density topology with respect to measure and category - J.Hejduk (2002)
- simple density topology - V.Aversa, W.Wilczyński (2004)
- complete density topology - W.Wilczyński, W.Wojdowski (2007)
- topologies connected with Hausdorff measures - G.H. (2008)
- $f$ -density topology - M.Filipczak, T.Filipczak (2008)
- $f$ -symmetrical density topology - J.Hejduk (2008)
- topology in the aspect of measure with respect to a sequence tending to zero - J.Hejduk, R.Wiertelak (2012)

## Abstract density topologies generated by ALD

- $\psi$ -density topology - M.Terepeta, E.Wagner-Bojakowska (1999)
- density topology with respect to the O'Malley points - W.Poreda, W.Wilczyński (2001)
- density topology with respect to measure and category - J.Hejduk (2002)
- simple density topology - V.Aversa, W.Wilczyński (2004)
- complete density topology - W.Wilczyński, W.Wojdowski (2007)
- topologies connected with Hausdorff measures - G.H. (2008)
- $f$ -density topology - M.Filipczak, T.Filipczak (2008)
- $f$ -symmetrical density topology - J.Hejduk (2008)
- topology in the aspect of measure with respect to a sequence tending to zero - J.Hejduk, R.Wiertelak (2012)



$(\mathbb{R}^n, \mathcal{S}, \mathcal{I})$   $(\mathbb{R}^n, \mathcal{T}_\Phi)$ -generated by ALD

$$(A \in \mathcal{T}_\Phi \wedge A \neq \emptyset) \Rightarrow A \notin \mathcal{I}$$

### Theorem

If every set  $A \in \mathcal{S} \setminus \mathcal{I}$  contains a perfect set ...

$\mathcal{I}$  - Borel and  $\mathcal{S} = \mathcal{Bor} \Delta \mathcal{I}$  ( $\mathcal{I} = \mathcal{N} \cap \mathcal{K}$ )

$(\mathcal{L}, \mathcal{N})$

$(\mathcal{B}, \mathcal{K})$

$(\mathbb{R}^n, \mathcal{S}, \mathcal{I})$   $(\mathbb{R}^n, \mathcal{T}_\Phi)$ -generated by ALD

$$(A \in \mathcal{T}_\Phi \wedge A \neq \emptyset) \Rightarrow A \notin \mathcal{I}$$

### Theorem

If every set  $A \in \mathcal{S} \setminus \mathcal{I}$  contains a perfect set ...

$\mathcal{I}$  - Borel and  $\mathcal{S} = \mathcal{Bor} \Delta \mathcal{I}$  ( $\mathcal{I} = \mathcal{N} \cap \mathcal{K}$ )

$(\mathcal{L}, \mathcal{N})$

$(\mathcal{B}, \mathcal{K})$

$(\mathbb{R}^n, \mathcal{S}, \mathcal{I})$   $(\mathbb{R}^n, \mathcal{T}_\Phi)$ -generated by ALD

$$(A \in \mathcal{T}_\Phi \wedge A \neq \emptyset) \Rightarrow A \notin \mathcal{I}$$

### Theorem

If every set  $A \in \mathcal{S} \setminus \mathcal{I}$  contains a perfect set ...

$\mathcal{I}$  - Borel and  $\mathcal{S} = \mathcal{Bor} \Delta \mathcal{I}$  ( $\mathcal{I} = \mathcal{N} \cap \mathcal{K}$ )

$(\mathcal{L}, \mathcal{N})$

$(\mathcal{B}, \mathcal{K})$

$(\mathbb{R}^n, \mathcal{S}, \mathcal{I})$   $(\mathbb{R}^n, \mathcal{T}_\Phi)$ -generated by ALD

$$(A \in \mathcal{T}_\Phi \wedge A \neq \emptyset) \Rightarrow A \notin \mathcal{I}$$

### Theorem

If every set  $A \in \mathcal{S} \setminus \mathcal{I}$  contains a perfect set ...

$\mathcal{I}$  - Borel and  $\mathcal{S} = \mathcal{Bor} \Delta \mathcal{I}$  ( $\mathcal{I} = \mathcal{N} \cap \mathcal{K}$ )

$(\mathcal{L}, \mathcal{N})$

$(\mathcal{B}, \mathcal{K})$

$(\mathbb{R}^n, \mathcal{S}, \mathcal{I})$   $(\mathbb{R}^n, \mathcal{T}_\Phi)$ -generated by ALD

$$(A \in \mathcal{T}_\Phi \wedge A \neq \emptyset) \Rightarrow A \notin \mathcal{I}$$

### Theorem

If every set  $A \in \mathcal{S} \setminus \mathcal{I}$  contains a perfect set ...

$\mathcal{I}$  - Borel and  $\mathcal{S} = \mathcal{Bor} \triangle \mathcal{I}$  ( $\mathcal{I} = \mathcal{N} \cap \mathcal{K}$ )

$(\mathcal{L}, \mathcal{N})$

$(\mathcal{B}, \mathcal{K})$

$(\mathbb{R}^n, \mathcal{S}, \mathcal{I})$   $(\mathbb{R}^n, \mathcal{T}_\Phi)$ -generated by ALD

$$(A \in \mathcal{T}_\Phi \wedge A \neq \emptyset) \Rightarrow A \notin \mathcal{I}$$

### Theorem

If every set  $A \in \mathcal{S} \setminus \mathcal{I}$  contains a perfect set ...

$\mathcal{I}$  - Borel and  $\mathcal{S} = \mathcal{Bor} \triangle \mathcal{I}$  ( $\mathcal{I} = \mathcal{N} \cap \mathcal{K}$ )

$(\mathcal{L}, \mathcal{N})$

$(\mathcal{B}, \mathcal{K})$

$(\mathbb{R}^n, \mathcal{S}, \mathcal{I})$   $(\mathbb{R}^n, \mathcal{T}_\Phi)$ -generated by ALD

$$(A \in \mathcal{T}_\Phi \wedge A \neq \emptyset) \Rightarrow A \notin \mathcal{I}$$

### Theorem

If every set  $A \in \mathcal{S} \setminus \mathcal{I}$  contains a perfect set  
then

- $\Delta(\mathbb{R}^n, \mathcal{T}_\Phi) = \mathfrak{c}$ ,
- every Bernstein set is dense in  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ .

$(\mathbb{R}^n, \mathcal{S}, \mathcal{I})$   $(\mathbb{R}^n, \mathcal{T}_\Phi)$ -generated by ALD

$$(A \in \mathcal{T}_\Phi \wedge A \neq \emptyset) \Rightarrow A \notin \mathcal{I}$$

### Theorem

If every set  $A \in \mathcal{S} \setminus \mathcal{I}$  contains a perfect set  
then

- $\Delta(\mathbb{R}^n, \mathcal{T}_\Phi) = \mathfrak{c}$ ,
- every Bernstein set is dense in  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ .



## Maximal resolvability

There exists a partition of  $\mathbb{R}^n$  into continuum many Bernstein sets.

### Theorem

Any topological space  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ , where  $\mathcal{T}_\Phi$  is a topology generated by ALD on  $(\mathbb{R}^n, \mathcal{S}, \mathcal{I})$  with a property that every  $A \in \mathcal{S} \setminus \mathcal{I}$  contains a perfect set, is maximally resolvable.

- J. Luukkainen - density topology on  $\mathbb{R}$
- E. Wagner-Bojakowska - I-density topology on  $\mathbb{R}$
- G.H. -  $\psi$ -density topology on  $\mathbb{R}$
- D. Rose, B. Thurston - lower density spaces on  $\mathbb{R}$

## Maximal resolvability

There exists a partition of  $\mathbb{R}^n$  into continuum many Bernstein sets.

### Theorem

Any topological space  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ , where  $\mathcal{T}_\Phi$  is a topology generated by ALD on  $(\mathbb{R}^n, \mathcal{S}, \mathcal{I})$  with a property that every  $A \in \mathcal{S} \setminus \mathcal{I}$  contains a perfect set, is maximally resolvable.

- J. Luukkainen - density topology on  $\mathbb{R}$
- E. Wagner-Bojakowska - I-density topology on  $\mathbb{R}$
- G.H. -  $\psi$ -density topology on  $\mathbb{R}$
- D. Rose, B. Thurston - lower density spaces on  $\mathbb{R}$

## Maximal resolvability

There exists a partition of  $\mathbb{R}^n$  into continuum many Bernstein sets.

### Theorem

Any topological space  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ , where  $\mathcal{T}_\Phi$  is a topology generated by ALD on  $(\mathbb{R}^n, \mathcal{S}, \mathcal{I})$  with a property that every  $A \in \mathcal{S} \setminus \mathcal{I}$  contains a perfect set, is maximally resolvable.

- J. Luukkainen - density topology on  $\mathbb{R}$
- E. Wagner-Bojakowska - I-density topology on  $\mathbb{R}$
- G.H. -  $\psi$ -density topology on  $\mathbb{R}$
- D. Rose, B. Thurston - lower density spaces on  $\mathbb{R}$

## Maximal resolvability

There exists a partition of  $\mathbb{R}^n$  into continuum many Bernstein sets.

### Theorem

Any topological space  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ , where  $\mathcal{T}_\Phi$  is a topology generated by ALD on  $(\mathbb{R}^n, \mathcal{S}, \mathcal{I})$  with a property that every  $A \in \mathcal{S} \setminus \mathcal{I}$  contains a perfect set, is maximally resolvable.

- J. Luukkainen - density topology on  $\mathbb{R}$
- E. Wagner-Bojakowska - l-density topology on  $\mathbb{R}$
- G.H. -  $\psi$ -density topology on  $\mathbb{R}$
- D. Rose, B. Thurston - lower density spaces on  $\mathbb{R}$

## Maximal resolvability

There exists a partition of  $\mathbb{R}^n$  into continuum many Bernstein sets.

### Theorem

Any topological space  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ , where  $\mathcal{T}_\Phi$  is a topology generated by ALD on  $(\mathbb{R}^n, \mathcal{S}, \mathcal{I})$  with a property that every  $A \in \mathcal{S} \setminus \mathcal{I}$  contains a perfect set, is maximally resolvable.

- J. Luukkainen - density topology on  $\mathbb{R}$
- E. Wagner-Bojakowska - l-density topology on  $\mathbb{R}$
- G.H. -  $\psi$ -density topology on  $\mathbb{R}$
- D. Rose, B. Thurston - lower density spaces on  $\mathbb{R}$

## Maximal resolvability

There exists a partition of  $\mathbb{R}^n$  into continuum many Bernstein sets.

### Theorem

Any topological space  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ , where  $\mathcal{T}_\Phi$  is a topology generated by ALD on  $(\mathbb{R}^n, \mathcal{S}, \mathcal{I})$  with a property that every  $A \in \mathcal{S} \setminus \mathcal{I}$  contains a perfect set, is maximally resolvable.

- J. Luukkainen - density topology on  $\mathbb{R}$
- E. Wagner-Bojakowska - l-density topology on  $\mathbb{R}$
- G.H. -  $\psi$ -density topology on  $\mathbb{R}$
- D. Rose, B. Thurston - lower density spaces on  $\mathbb{R}$

## Extraresolvability

Under MA in  $\mathbb{R}^n$  there exists a family of cardinality  $2^c$  almost disjoint Bernstein sets.

### Theorem

(ZFC+MA) Any topological space  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ , where  $\mathcal{T}_\Phi$  is a topology generated by ALD on  $(\mathbb{R}^n, \mathcal{S}, \mathcal{I})$  with a property that every  $A \in \mathcal{S} \setminus \mathcal{I}$  contains a perfect set, is extraresolvable.

(ZFC+MA)

- A. Bella - density topology on  $\mathbb{R}$
- E. Wagner-Bojakowska - I-density topology on  $\mathbb{R}$
- G.H. -  $\psi$ -density topology on  $\mathbb{R}$
- D. Rose, B. Thurston - lower density spaces on  $\mathbb{R}$

T. Natkaniec, *The density topology can be not extraresolvable* Real Anal. Exchange 30(1) (2004-2005).

## Extraresolvability

Under MA in  $\mathbb{R}^n$  there exists a family of cardinality  $2^c$  almost disjoint Bernstein sets.

### Theorem

(ZFC+MA) Any topological space  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ , where  $\mathcal{T}_\Phi$  is a topology generated by ALD on  $(\mathbb{R}^n, \mathcal{S}, \mathcal{I})$  with a property that every  $A \in \mathcal{S} \setminus \mathcal{I}$  contains a perfect set, is extraresolvable.

(ZFC+MA)

- A. Bella - density topology on  $\mathbb{R}$
- E. Wagner-Bojakowska - I-density topology on  $\mathbb{R}$
- G.H. -  $\psi$ -density topology on  $\mathbb{R}$
- D. Rose, B. Thurston - lower density spaces on  $\mathbb{R}$

T. Natkaniec, *The density topology can be not extraresolvable* Real Anal. Exchange 30(1) (2004-2005).



## Extraresolvability

Under MA in  $\mathbb{R}^n$  there exists a family of cardinality  $2^c$  almost disjoint Bernstein sets.

### Theorem

(ZFC+MA) Any topological space  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ , where  $\mathcal{T}_\Phi$  is a topology generated by ALD on  $(\mathbb{R}^n, \mathcal{S}, \mathcal{I})$  with a property that every  $A \in \mathcal{S} \setminus \mathcal{I}$  contains a perfect set, is extraresolvable.

(ZFC+MA)

- A. Bella - density topology on  $\mathbb{R}$
- E. Wagner-Bojakowska - I-density topology on  $\mathbb{R}$
- G.H. -  $\psi$ -density topology on  $\mathbb{R}$
- D. Rose, B. Thurston - lower density spaces on  $\mathbb{R}$

T. Natkaniec, *The density topology can be not extraresolvable* Real Anal. Exchange 30(1) (2004-2005).

## Extraresolvability

Under MA in  $\mathbb{R}^n$  there exists a family of cardinality  $2^c$  almost disjoint Bernstein sets.

### Theorem

(ZFC+MA) Any topological space  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ , where  $\mathcal{T}_\Phi$  is a topology generated by ALD on  $(\mathbb{R}^n, \mathcal{S}, \mathcal{I})$  with a property that every  $A \in \mathcal{S} \setminus \mathcal{I}$  contains a perfect set, is extraresolvable.

(ZFC+MA)

- A. Bella - density topology on  $\mathbb{R}$
- E. Wagner-Bojakowska - I-density topology on  $\mathbb{R}$
- G.H. -  $\psi$ -density topology on  $\mathbb{R}$
- D. Rose, B. Thurston - lower density spaces on  $\mathbb{R}$

T. Natkaniec, *The density topology can be not extraresolvable* Real Anal. Exchange 30(1) (2004-2005).

## Extraresolvability

Under MA in  $\mathbb{R}^n$  there exists a family of cardinality  $2^c$  almost disjoint Bernstein sets.

### Theorem

(ZFC+MA) Any topological space  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ , where  $\mathcal{T}_\Phi$  is a topology generated by ALD on  $(\mathbb{R}^n, \mathcal{S}, \mathcal{I})$  with a property that every  $A \in \mathcal{S} \setminus \mathcal{I}$  contains a perfect set, is extraresolvable.

(ZFC+MA)

- A. Bella - density topology on  $\mathbb{R}$
- E. Wagner-Bojakowska - I-density topology on  $\mathbb{R}$
- G.H. -  $\psi$ -density topology on  $\mathbb{R}$
- D. Rose, B. Thurston - lower density spaces on  $\mathbb{R}$

T. Natkaniec, *The density topology can be not extraresolvable* Real Anal. Exchange 30(1) (2004-2005).

## Extraresolvability

Under MA in  $\mathbb{R}^n$  there exists a family of cardinality  $2^c$  almost disjoint Bernstein sets.

### Theorem

(ZFC+MA) Any topological space  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ , where  $\mathcal{T}_\Phi$  is a topology generated by ALD on  $(\mathbb{R}^n, \mathcal{S}, \mathcal{I})$  with a property that every  $A \in \mathcal{S} \setminus \mathcal{I}$  contains a perfect set, is extraresolvable.

(ZFC+MA)

- A. Bella - density topology on  $\mathbb{R}$
- E. Wagner-Bojakowska - I-density topology on  $\mathbb{R}$
- G.H. -  $\psi$ -density topology on  $\mathbb{R}$
- D. Rose, B. Thurston - lower density spaces on  $\mathbb{R}$

T. Natkaniec, *The density topology can be not extraresolvable* Real Anal. Exchange 30(1) (2004-2005).

## Extraresolvability

Under MA in  $\mathbb{R}^n$  there exists a family of cardinality  $2^c$  almost disjoint Bernstein sets.

### Theorem

(ZFC+MA) Any topological space  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ , where  $\mathcal{T}_\Phi$  is a topology generated by ALD on  $(\mathbb{R}^n, \mathcal{S}, \mathcal{I})$  with a property that every  $A \in \mathcal{S} \setminus \mathcal{I}$  contains a perfect set, is extraresolvable.

(ZFC+MA)

- A. Bella - density topology on  $\mathbb{R}$
- E. Wagner-Bojakowska - I-density topology on  $\mathbb{R}$
- G.H. -  $\psi$ -density topology on  $\mathbb{R}$
- D. Rose, B. Thurston - lower density spaces on  $\mathbb{R}$

T. Natkaniec, *The density topology can be not extraresolvable* Real Anal. Exchange 30(1) (2004-2005).

## Extraresolvability

Under MA in  $\mathbb{R}^n$  there exists a family of cardinality  $2^c$  almost disjoint Bernstein sets.

### Theorem

(ZFC+MA) Any topological space  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ , where  $\mathcal{T}_\Phi$  is a topology generated by ALD on  $(\mathbb{R}^n, \mathcal{S}, \mathcal{I})$  with a property that every  $A \in \mathcal{S} \setminus \mathcal{I}$  contains a perfect set, is extraresolvable.

(ZFC+MA)

- A. Bella - density topology on  $\mathbb{R}$
- E. Wagner-Bojakowska - I-density topology on  $\mathbb{R}$
- G.H. -  $\psi$ -density topology on  $\mathbb{R}$
- D. Rose, B. Thurston - lower density spaces on  $\mathbb{R}$

T. Natkaniec, *The density topology can be not extraresolvable* Real Anal. Exchange 30(1) (2004-2005).

## Extraresolvability - negative results for LD

## Theorem

There exists a model of ZFC in which the space  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ , where  $\mathcal{T}_\Phi$  is a topology generated by a lower density operator on  $(\mathbb{R}^n, \mathcal{L}, \mathcal{N})$  or on  $(\mathbb{R}^n, \mathcal{B}, \mathcal{K})$ , is not extraresolvable.

ZFC+CPA (Covering Property Axiom) and an additional assumption:  
 $2^{\omega_1} = \omega_2$ .

Consequences of CPA:

- $\mathfrak{c} = \omega_2$
- $cf(\mathcal{N}) = \omega_1$

$$cf(\mathcal{I}) := \min\{\text{card}(\mathcal{I}_0) : \mathcal{I}_0 \subset \mathcal{I} \wedge \forall J \in \mathcal{I} \exists J_0 \in \mathcal{I}_0 J \subset J_0\}$$

$$cf(\mathcal{I}) = \omega_1$$

$$NWD_{\mathcal{T}_\Phi} = \mathcal{I}$$

## Extraresolvability - negative results for LD

### Theorem

There exists a model of ZFC in which the space  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ , where  $\mathcal{T}_\Phi$  is a topology generated by a lower density operator on  $(\mathbb{R}^n, \mathcal{L}, \mathcal{N})$  or on  $(\mathbb{R}^n, \mathcal{B}, \mathcal{K})$ , is not extraresolvable.

ZFC+CPA (Covering Property Axiom) and an additional assumption:  
 $2^{\omega_1} = \omega_2$ .

Consequences of CPA:

- $\mathfrak{c} = \omega_2$
- $cf(\mathcal{N}) = \omega_1$

$$cf(\mathcal{I}) := \min\{\text{card}(\mathcal{I}_0) : \mathcal{I}_0 \subset \mathcal{I} \wedge \forall J \in \mathcal{I} \exists J_0 \in \mathcal{I}_0 J \subset J_0\}$$

$$cf(\mathcal{I}) = \omega_1$$

$$NWD_{\mathcal{T}_\Phi} = \mathcal{I}$$



## Extraresolvability - negative results for LD

## Theorem

There exists a model of ZFC in which the space  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ , where  $\mathcal{T}_\Phi$  is a topology generated by a lower density operator on  $(\mathbb{R}^n, \mathcal{L}, \mathcal{N})$  or on  $(\mathbb{R}^n, \mathcal{B}, \mathcal{K})$ , is not extraresolvable.

ZFC+CPA (Covering Property Axiom) and an additional assumption:  
 $2^{\omega_1} = \omega_2$ .

Consequences of CPA:

- $\mathfrak{c} = \omega_2$
- $cf(\mathcal{N}) = \omega_1$

$$cf(\mathcal{I}) := \min\{\text{card}(\mathcal{I}_0) : \mathcal{I}_0 \subset \mathcal{I} \wedge \forall J \in \mathcal{I} \exists J_0 \in \mathcal{I}_0 J \subset J_0\}$$

$$cf(\mathcal{I}) = \omega_1$$

$$NWD_{\mathcal{T}_\Phi} = \mathcal{I}$$

## Extraresolvability - negative results for LD

## Theorem

There exists a model of ZFC in which the space  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ , where  $\mathcal{T}_\Phi$  is a topology generated by a lower density operator on  $(\mathbb{R}^n, \mathcal{L}, \mathcal{N})$  or on  $(\mathbb{R}^n, \mathcal{B}, \mathcal{K})$ , is not extraresolvable.

ZFC+CPA (Covering Property Axiom) and an additional assumption:  
 $2^{\omega_1} = \omega_2$ .

Consequences of CPA:

- $\mathfrak{c} = \omega_2$
- $cf(\mathcal{N}) = \omega_1$

$$cf(\mathcal{I}) := \min\{\text{card}(\mathcal{I}_0) : \mathcal{I}_0 \subset \mathcal{I} \wedge \forall J \in \mathcal{I} \exists J_0 \in \mathcal{I}_0 J \subset J_0\}$$

$$cf(\mathcal{I}) = \omega_1$$

$$NWD_{\mathcal{T}_\Phi} = \mathcal{I}$$

## Extraresolvability - negative results for LD

### Theorem

There exists a model of ZFC in which the space  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ , where  $\mathcal{T}_\Phi$  is a topology generated by a lower density operator on  $(\mathbb{R}^n, \mathcal{L}, \mathcal{N})$  or on  $(\mathbb{R}^n, \mathcal{B}, \mathcal{K})$ , is not extraresolvable.

ZFC+CPA (Covering Property Axiom) and an additional assumption:  
 $2^{\omega_1} = \omega_2$ .

Consequences of CPA:

- $\mathfrak{c} = \omega_2$
- $cf(\mathcal{N}) = \omega_1$

$$cf(\mathcal{I}) := \min\{\text{card}(\mathcal{I}_0) : \mathcal{I}_0 \subset \mathcal{I} \wedge \forall J \in \mathcal{I} \exists J_0 \in \mathcal{I}_0 J \subset J_0\}$$

$$cf(\mathcal{I}) = \omega_1$$

$$NWD_{\mathcal{T}_\Phi} = \mathcal{I}$$

## Extraresolvability - negative results for LD

### Theorem

There exists a model of ZFC in which the space  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ , where  $\mathcal{T}_\Phi$  is a topology generated by a lower density operator on  $(\mathbb{R}^n, \mathcal{L}, \mathcal{N})$  or on  $(\mathbb{R}^n, \mathcal{B}, \mathcal{K})$ , is not extraresolvable.

ZFC+CPA (Covering Property Axiom) and an additional assumption:  
 $2^{\omega_1} = \omega_2$ .

Consequences of CPA:

- $\mathfrak{c} = \omega_2$
- $cf(\mathcal{N}) = \omega_1$

$$cf(\mathcal{I}) := \min\{\text{card}(\mathcal{I}_0) : \mathcal{I}_0 \subset \mathcal{I} \wedge \forall J \in \mathcal{I} \exists J_0 \in \mathcal{I}_0 J \subset J_0\}$$

$$cf(\mathcal{I}) = \omega_1$$

$$NWD_{\mathcal{T}_\Phi} = \mathcal{I}$$

## Extraresolvability - negative results for LD

## Theorem

There exists a model of ZFC in which the space  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ , where  $\mathcal{T}_\Phi$  is a topology generated by a lower density operator on  $(\mathbb{R}^n, \mathcal{L}, \mathcal{N})$  or on  $(\mathbb{R}^n, \mathcal{B}, \mathcal{K})$ , is not extraresolvable.

ZFC+CPA (Covering Property Axiom) and an additional assumption:  
 $2^{\omega_1} = \omega_2$ .

Consequences of CPA:

- $\mathfrak{c} = \omega_2$
- $cf(\mathcal{N}) = \omega_1$

$$cf(\mathcal{I}) := \min\{\text{card}(\mathcal{I}_0) : \mathcal{I}_0 \subset \mathcal{I} \wedge \forall J \in \mathcal{I} \exists J_0 \in \mathcal{I}_0 J \subset J_0\}$$

$$cf(\mathcal{I}) = \omega_1$$

$$NWD_{\mathcal{T}_\Phi} = \mathcal{I}$$

?

Does a model of ZFC exist in which the space  $(\mathbb{R}^n, \mathcal{T}_\Phi)$ , where  $\mathcal{T}_\Phi$  is a topology generated by an almost lower density operator on  $(\mathbb{R}^n, \mathcal{L}, \mathcal{N})$  or on  $(\mathbb{R}^n, \mathcal{B}, \mathcal{K})$ , is not extraresolvable?

## Bibliography

- Bella A., *The Density topology is extraresolvable* Atti Sem. Mat. Fis. Univ. Modena 48 (2000), 495-498.
- Comfort W.W., Negrepontis S., *The Theory of Ultrafilters* Grundle Math. Wissensch. 211, Springer-Verlag, Berlin 1974.
- Ciesielski K., Pawlikowski J., *Covering Property Axiom CPA. A combinatorial core of the iterated perfect set model* Cambridge Tracts in Mathematics 164, Cambridge Univ. Press, 2004.
- Hejduk J., Wiertelak R., *On the abstract density topologies generated by lower and almost lower density operators* Traditional and present-day topics in real analysis, Łódź University Press, 2013.
- Hewitt E., *A problem of set-theoretic topology* Duke Math. J. 10 (1943), 309-333.
- Horbaczewska G., *Resolvability of  $\psi$ -density topology* Bull. de la Societe des sciences et des letters de Łódź LII (2002), 5-9.
- Luukkainen J., *The density topology is maximally resolvable* Real Anal. Exchange 25(1) (1999-2000), 419-420.
- Natkaniec T., *The density topology can be not extraresolvable* Real Anal. Exchange 30(1) (2004-2005), 393-396.
- Rose D., Thurston B., *Maximally resolvable lower density spaces* Real Anal. Exchange 33(2) (2007-2008), 475-482.
- Wagner-Bojakowska E., *Resolvability of  $l$ -density topology* Zeszyty Naukowe WSIn 1(1) (2001), 35-37.