

# Generalizations of Urysohn's lemma for the family of extra strong Świątkowski functions

Paulina Szczuka

Department of Mathematics  
Kazimierz Wielki University  
Bydgoszcz, POLAND

Stara Lesna 2014

# Introduction

Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be a function.

## Definition

We say that  $f$  is *Darboux*, if it maps connected sets onto connected sets; i.e., if whenever  $a < b$  and  $y$  is a number between  $f(a)$  and  $f(b)$ , there is  $x_0 \in (a, b)$  such that  $f(x_0) = y$ .

## Definition

We say that  $f$  is *quasi-continuous* in the sense of Kempisty, if for all  $x \in \mathbb{R}$  and open sets  $U \ni x$  and  $V \ni f(x)$ , the set  $\text{int}(U \cap f^{-1}(V)) \neq \emptyset$ .

# Introduction

Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be a function.

## Definition

We say that  $f$  is *Darboux*, if it maps connected sets onto connected sets; i.e., if whenever  $a < b$  and  $y$  is a number between  $f(a)$  and  $f(b)$ , there is  $x_0 \in (a, b)$  such that  $f(x_0) = y$ .

## Definition

We say that  $f$  is *quasi-continuous* in the sense of Kempisty, if for all  $x \in \mathbb{R}$  and open sets  $U \ni x$  and  $V \ni f(x)$ , the set  $\text{int}(U \cap f^{-1}(V)) \neq \emptyset$ .

# Introduction

Let  $f: \mathbb{R} \rightarrow \mathbb{R}$  be a function.

## Definition

We say that  $f$  is *Darboux*, if it maps connected sets onto connected sets; i.e., if whenever  $a < b$  and  $y$  is a number between  $f(a)$  and  $f(b)$ , there is  $x_0 \in (a, b)$  such that  $f(x_0) = y$ .

## Definition

We say that  $f$  is *quasi-continuous* in the sense of Kempisty, if for all  $x \in \mathbb{R}$  and open sets  $U \ni x$  and  $V \ni f(x)$ , the set  $\text{int}(U \cap f^{-1}(V)) \neq \emptyset$ .

# Introduction

The symbol  $\mathcal{C}(f)$  denotes the set of points of continuity of  $f$ .

## Definition

We say that  $f$  is *strong Świątkowski*, if whenever  $a < b$  and  $y$  is a number between  $f(a)$  and  $f(b)$ , there is  $x_0 \in (a, b) \cap \mathcal{C}(f)$  such that  $f(x_0) = y$ .

## Definition

We say that  $f$  is *extra strong Świątkowski*, if whenever  $a < b$  and  $y$  belongs to the closed interval with endpoints  $f(a)$  and  $f(b)$ , there is  $x_0 \in [a, b] \cap \mathcal{C}(f)$  such that  $f(x_0) = y$ .

# Introduction

The symbol  $\mathcal{C}(f)$  denotes the set of points of continuity of  $f$ .

## Definition

We say that  $f$  is *strong Świątkowski*, if whenever  $a < b$  and  $y$  is a number between  $f(a)$  and  $f(b)$ , there is  $x_0 \in (a, b) \cap \mathcal{C}(f)$  such that  $f(x_0) = y$ .

## Definition

We say that  $f$  is *extra strong Świątkowski*, if whenever  $a < b$  and  $y$  belongs to the closed interval with endpoints  $f(a)$  and  $f(b)$ , there is  $x_0 \in [a, b] \cap \mathcal{C}(f)$  such that  $f(x_0) = y$ .

# Introduction

The symbol  $\mathcal{C}(f)$  denotes the set of points of continuity of  $f$ .

## Definition

We say that  $f$  is *strong Świątkowski*, if whenever  $a < b$  and  $y$  is a number between  $f(a)$  and  $f(b)$ , there is  $x_0 \in (a, b) \cap \mathcal{C}(f)$  such that  $f(x_0) = y$ .

## Definition

We say that  $f$  is *extra strong Świątkowski*, if whenever  $a < b$  and  $y$  belongs to the closed interval with endpoints  $f(a)$  and  $f(b)$ , there is  $x_0 \in [a, b] \cap \mathcal{C}(f)$  such that  $f(x_0) = y$ .

# Introduction

The symbols below denote the following families of real functions:

- $\mathcal{C}$  – the class of all continuous functions,
- $\mathcal{D}$  – the class of all Darboux functions,
- $\mathcal{Q}$  – the class of all quasi-continuous functions,
- $\mathcal{DQ}$  – the class of all Darboux quasi-continuous functions,
- $\acute{S}_s$  – the class of all strong Świątkowski functions,
- $\acute{S}_{es}$  – the class of all extra strong Świątkowski functions.



# Introduction

The symbols below denote the following families of real functions:

- $\mathcal{C}$  – the class of all continuous functions,
- $\mathcal{D}$  – the class of all Darboux functions,
- $\mathcal{Q}$  – the class of all quasi-continuous functions,
- $\mathcal{DQ}$  – the class of all Darboux quasi-continuous functions,
- $\mathcal{S}_s$  – the class of all strong Świątkowski functions,
- $\mathcal{S}_{es}$  – the class of all extra strong Świątkowski functions.

# Introduction

The symbols below denote the following families of real functions:

- $\mathcal{C}$  – the class of all continuous functions,
- $\mathcal{D}$  – the class of all Darboux functions,
- $\mathcal{Q}$  – the class of all quasi-continuous functions,
- $\mathcal{DQ}$  – the class of all Darboux quasi-continuous functions,
- $\mathcal{S}_s$  – the class of all strong Świątkowski functions,
- $\mathcal{S}_{es}$  – the class of all extra strong Świątkowski functions.

# Introduction

The symbols below denote the following families of real functions:

- $\mathcal{C}$  – the class of all continuous functions,
- $\mathcal{D}$  – the class of all Darboux functions,
- $\mathcal{Q}$  – the class of all quasi-continuous functions,
- $\mathcal{DQ}$  – the class of all Darboux quasi-continuous functions,
- $\acute{S}_s$  – the class of all strong Świątkowski functions,
- $\acute{S}_{es}$  – the class of all extra strong Świątkowski functions.

# Introduction

The symbols below denote the following families of real functions:

- $\mathcal{C}$  – the class of all continuous functions,
- $\mathcal{D}$  – the class of all Darboux functions,
- $\mathcal{Q}$  – the class of all quasi-continuous functions,
- $\mathcal{DQ}$  – the class of all Darboux quasi-continuous functions,
- $\mathcal{S}_s$  – the class of all strong Świątkowski functions,
- $\mathcal{S}_{es}$  – the class of all extra strong Świątkowski functions.

# Introduction

The symbols below denote the following families of real functions:

- $\mathcal{C}$  – the class of all continuous functions,
- $\mathcal{D}$  – the class of all Darboux functions,
- $\mathcal{Q}$  – the class of all quasi-continuous functions,
- $\mathcal{DQ}$  – the class of all Darboux quasi-continuous functions,
- $\acute{S}_s$  – the class of all strong Świątkowski functions,
- $\acute{S}_{es}$  – the class of all extra strong Świątkowski functions.

# Introduction

The symbols below denote the following families of real functions:

- $\mathcal{C}$  – the class of all continuous functions,
- $\mathcal{D}$  – the class of all Darboux functions,
- $\mathcal{Q}$  – the class of all quasi-continuous functions,
- $\mathcal{DQ}$  – the class of all Darboux quasi-continuous functions,
- $\acute{\mathcal{S}}_s$  – the class of all strong Świątkowski functions,
- $\acute{\mathcal{S}}_{es}$  – the class of all extra strong Świątkowski functions.

# Introduction

It can be readily verified that we have the following proper inclusions:

- $\mathcal{C} \subsetneq \dot{\mathcal{S}}_{es} \subsetneq \dot{\mathcal{S}}_s \subsetneq \mathcal{DQ} \subsetneq \mathcal{D},$
- $\mathcal{C} \subsetneq \dot{\mathcal{S}}_{es} \subsetneq \dot{\mathcal{S}}_s \subsetneq \mathcal{DQ} \subsetneq \mathcal{Q}.$

## Definition

We say that a set  $A \subset \mathbb{R}$  is *semi-open*, if  $A \subset \text{cl int } A$ , and it is *semi-closed*, if  $\text{int cl } A \subset A$ .

# Introduction

It can be readily verified that we have the following proper inclusions:

$$\bullet \mathcal{C} \subsetneq \mathcal{S}_{es} \subsetneq \mathcal{S}_s \subsetneq \mathcal{DQ} \subsetneq \mathcal{D},$$

$$\bullet \mathcal{C} \subsetneq \mathcal{S}_{es} \subsetneq \mathcal{S}_s \subsetneq \mathcal{DQ} \subsetneq \mathcal{Q}.$$

## Definition

We say that a set  $A \subset \mathbb{R}$  is *semi-open*, if  $A \subset \text{cl int } A$ , and it is *semi-closed*, if  $\text{int cl } A \subset A$ .



# Introduction

It can be readily verified that we have the following proper inclusions:

- $\mathcal{C} \subsetneq \mathcal{S}_{es} \subsetneq \mathcal{S}_s \subsetneq \mathcal{DQ} \subsetneq \mathcal{D}$ ,
- $\mathcal{C} \subsetneq \mathcal{S}_{es} \subsetneq \mathcal{S}_s \subsetneq \mathcal{DQ} \subsetneq \mathcal{Q}$ .

## Definition

We say that a set  $A \subset \mathbb{R}$  is *semi-open*, if  $A \subset \text{cl int } A$ , and it is *semi-closed*, if  $\text{int cl } A \subset A$ .

# Introduction

It can be readily verified that we have the following proper inclusions:

- $\mathcal{C} \subsetneq \mathcal{S}_{es} \subsetneq \mathcal{S}_s \subsetneq \mathcal{DQ} \subsetneq \mathcal{D}$ ,
- $\mathcal{C} \subsetneq \mathcal{S}_{es} \subsetneq \mathcal{S}_s \subsetneq \mathcal{DQ} \subsetneq \mathcal{Q}$ .

## Definition

We say that a set  $A \subset \mathbb{R}$  is *semi-open*, if  $A \subset \text{cl int } A$ , and it is *semi-closed*, if  $\text{int cl } A \subset A$ .

# Introduction

It can be readily verified that we have the following proper inclusions:

- $\mathcal{C} \subsetneq \mathcal{S}_{es} \subsetneq \mathcal{S}_s \subsetneq \mathcal{DQ} \subsetneq \mathcal{D}$ ,
- $\mathcal{C} \subsetneq \mathcal{S}_{es} \subsetneq \mathcal{S}_s \subsetneq \mathcal{DQ} \subsetneq \mathcal{Q}$ .

## Definition

We say that a set  $A \subset \mathbb{R}$  is *semi-open*, if  $A \subset \text{cl int } A$ , and it is *semi-closed*, if  $\text{int cl } A \subset A$ .

Notice that a set  $A \subset \mathbb{R}$  is semi-open if and only if its complement is semi-closed.

# Classical separation property

## Lemma (Urysohn 1925)

*Let  $X$  be a normal topological space and sets  $A_0, A_1 \subset X$  be disjoint and closed. Then there exists a continuous function  $f: X \rightarrow \mathbb{R}$  such that  $f = 0$  on  $A_0$  and  $f = 1$  on  $A_1$ . Moreover, if  $A_0$  and  $A_1$  are  $G_\delta$ -sets, then we can require that  $f(x) \in (0, 1)$  for each  $x \in X \setminus (A_0 \cup A_1)$ .*

# Classical separation property

The symbol  $[f = a]$  stands for the set  $\{x \in \mathbb{R} ; f(x) = a\}$ .

## Theorem

*Let  $X$  be a perfectly normal topological space and sets  $A_0, A_1 \subset X$  be disjoint. There is a continuous function  $f: X \rightarrow \mathbb{R}$  such that  $A_0 = [f = 0]$  and  $A_1 = [f = 1]$  if and only if sets  $A_0$  and  $A_1$  are closed.*

## Theorem

*Let  $X$  be a normal topological space and sets  $A_0, A_1 \subset X$ . There is a continuous function  $f: X \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$  if and only if sets  $\text{cl } A_0 \cap \text{cl } A_1 = \emptyset$ .*

# Classical separation property

The symbol  $[f = a]$  stands for the set  $\{x \in \mathbb{R} ; f(x) = a\}$ .  
Similarly we define  $[f > a]$  and  $[f < a]$ .

## Theorem

*Let  $X$  be a perfectly normal topological space and sets  $A_0, A_1 \subset X$  be disjoint. There is a continuous function  $f: X \rightarrow \mathbb{R}$  such that  $A_0 = [f = 0]$  and  $A_1 = [f = 1]$  if and only if sets  $A_0$  and  $A_1$  are closed.*

## Theorem

*Let  $X$  be a normal topological space and sets  $A_0, A_1 \subset X$ . There is a continuous function  $f: X \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$  if and only if sets  $\text{cl } A_0 \cap \text{cl } A_1 = \emptyset$ .*

# Classical separation property

The symbol  $[f = a]$  stands for the set  $\{x \in \mathbb{R} ; f(x) = a\}$ .  
Similarly we define  $[f > a]$  and  $[f < a]$ .

## Theorem

*Let  $X$  be a perfectly normal topological space and sets  $A_0, A_1 \subset X$  be disjoint. There is a continuous function  $f: X \rightarrow \mathbb{R}$  such that  $A_0 = [f = 0]$  and  $A_1 = [f = 1]$  if and only if sets  $A_0$  and  $A_1$  are closed.*

## Theorem

*Let  $X$  be a normal topological space and sets  $A_0, A_1 \subset X$ . There is a continuous function  $f: X \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$  if and only if sets  $\text{cl } A_0 \cap \text{cl } A_1 = \emptyset$ .*

# Classical separation property

The symbol  $[f = a]$  stands for the set  $\{x \in \mathbb{R} ; f(x) = a\}$ .  
Similarly we define  $[f > a]$  and  $[f < a]$ .

## Theorem

*Let  $X$  be a perfectly normal topological space and sets  $A_0, A_1 \subset X$  be disjoint. There is a continuous function  $f: X \rightarrow \mathbb{R}$  such that  $A_0 = [f = 0]$  and  $A_1 = [f = 1]$  if and only if sets  $A_0$  and  $A_1$  are closed.*

## Theorem

*Let  $X$  be a normal topological space and sets  $A_0, A_1 \subset X$ . There is a continuous function  $f: X \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$  if and only if sets  $\text{cl } A_0 \cap \text{cl } A_1 = \emptyset$ .*



# Classical separation property

## Theorem (Maliszewski 2002)

*Let  $A_0, A_1 \subset \mathbb{R}$  be disjoint. There is a Darboux function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 = [f = 0]$  and  $A_1 = [f = 1]$  if and only if the set  $\mathbb{R} \setminus (A_0 \cup A_1)$  is bilaterally  $\mathfrak{c}$ -dense in itself and  $\text{card}([ \alpha, \beta ] \setminus (A_0 \cup A_1)) = \mathfrak{c}$  for each  $\alpha \in A_0$  and  $\beta \in A_1$ .*

## Theorem (Maliszewski 2002)

*Let  $A_0, A_1 \subset \mathbb{R}$  be disjoint. There is a Darboux function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$  if and only if  $\text{card}([ \alpha, \beta ] \setminus (A_0 \cup A_1)) = \mathfrak{c}$  for each  $\alpha \in A_0$  and  $\beta \in A_1$ .*

# Classical separation property

## Theorem (Maliszewski 2002)

*Let  $A_0, A_1 \subset \mathbb{R}$  be disjoint. There is a Darboux function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 = [f = 0]$  and  $A_1 = [f = 1]$  if and only if the set  $\mathbb{R} \setminus (A_0 \cup A_1)$  is bilaterally  $\mathfrak{c}$ -dense in itself and  $\text{card}([ \alpha, \beta ] \setminus (A_0 \cup A_1)) = \mathfrak{c}$  for each  $\alpha \in A_0$  and  $\beta \in A_1$ .*

## Theorem (Maliszewski 2002)

*Let  $A_0, A_1 \subset \mathbb{R}$  be disjoint. There is a Darboux function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$  if and only if  $\text{card}([ \alpha, \beta ] \setminus (A_0 \cup A_1)) = \mathfrak{c}$  for each  $\alpha \in A_0$  and  $\beta \in A_1$ .*

# Classical separation property

## Theorem (Kowalewski 2005)

*Let  $A_0, A_1 \subset \mathbb{R}$  be disjoint. There is a quasi-continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 = [f = 0]$  and  $A_1 = [f = 1]$  if and only if sets  $A_0$ ,  $A_1$  and  $A_0 \cup A_1$  are semi-closed.*

## Theorem (Kowalewski 2005)

*Let  $A_0, A_1 \subset \mathbb{R}$ . There is a quasi-continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$  if and only if  $A_0 \cap A_1 = \text{int cl } A_0 \cap A_1 = A_0 \cap \text{int cl } A_1 = \emptyset$ .*

# Classical separation property

## Theorem (Kowalewski 2005)

*Let  $A_0, A_1 \subset \mathbb{R}$  be disjoint. There is a quasi-continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 = [f = 0]$  and  $A_1 = [f = 1]$  if and only if sets  $A_0$ ,  $A_1$  and  $A_0 \cup A_1$  are semi-closed.*

## Theorem (Kowalewski 2005)

*Let  $A_0, A_1 \subset \mathbb{R}$ . There is a quasi-continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$  if and only if  $A_0 \cap A_1 = \text{int cl } A_0 \cap A_1 = A_0 \cap \text{int cl } A_1 = \emptyset$ .*

# Classical separation property

## Theorem (Kowalewski and Maliszewski 2008)

*Let  $A_0, A_1 \subset \mathbb{R}$  be disjoint. The following conditions are equivalent:*

- 1 *there is a strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 = [f = 0]$  and  $A_1 = [f = 1]$ ,*
- 2 *there is a Darboux quasi-continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 = [f = 0]$  and  $A_1 = [f = 1]$ ,*
- 3 *sets  $A_0$  and  $A_1$  are semi-closed,  $\mathbb{R} \setminus (A_0 \cup A_1)$  is bilaterally dense in itself and  $I(\alpha, \beta) \setminus \text{cl}(A_0 \cup A_1) \neq \emptyset$  for each  $\alpha \in A_0$  and  $\beta \in A_1$ .*

# Classical separation property

## Theorem (Kowalewski and Maliszewski 2008)

*Let  $A_0, A_1 \subset \mathbb{R}$  be disjoint. The following conditions are equivalent:*

- ❶ *there is a strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 = [f = 0]$  and  $A_1 = [f = 1]$ ,*
- ❷ *there is a Darboux quasi-continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 = [f = 0]$  and  $A_1 = [f = 1]$ ,*
- ❸ *sets  $A_0$  and  $A_1$  are semi-closed,  $\mathbb{R} \setminus (A_0 \cup A_1)$  is bilaterally dense in itself and  $I(\alpha, \beta) \setminus \text{cl}(A_0 \cup A_1) \neq \emptyset$  for each  $\alpha \in A_0$  and  $\beta \in A_1$ .*

# Classical separation property

## Theorem (Kowalewski and Maliszewski 2008)

*Let  $A_0, A_1 \subset \mathbb{R}$  be disjoint. The following conditions are equivalent:*

- ❶ *there is a strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 = [f = 0]$  and  $A_1 = [f = 1]$ ,*
- ❷ *there is a Darboux quasi-continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 = [f = 0]$  and  $A_1 = [f = 1]$ ,*
- ❸ *sets  $A_0$  and  $A_1$  are semi-closed,  $\mathbb{R} \setminus (A_0 \cup A_1)$  is bilaterally dense in itself and  $I(\alpha, \beta) \setminus \text{cl}(A_0 \cup A_1) \neq \emptyset$  for each  $\alpha \in A_0$  and  $\beta \in A_1$ .*

# Classical separation property

## Theorem (Kowalewski and Maliszewski 2008)

*Let  $A_0, A_1 \subset \mathbb{R}$  be disjoint. The following conditions are equivalent:*

- ❶ *there is a strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 = [f = 0]$  and  $A_1 = [f = 1]$ ,*
- ❷ *there is a Darboux quasi-continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 = [f = 0]$  and  $A_1 = [f = 1]$ ,*
- ❸ *sets  $A_0$  and  $A_1$  are semi-closed,  $\mathbb{R} \setminus (A_0 \cup A_1)$  is bilaterally dense in itself and  $I(\alpha, \beta) \setminus \text{cl}(A_0 \cup A_1) \neq \emptyset$  for each  $\alpha \in A_0$  and  $\beta \in A_1$ .*



# Classical separation property

Theorem (Kowalewski and Maliszewski 2008; Szczuka 2014)

*Let  $A_0, A_1 \subset \mathbb{R}$  be disjoint. The following conditions are equivalent:*

- 1 *there is an extra strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$ ,*
- 2 *there is a strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$ ,*
- 3 *there is a Darboux quasi-continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$ ,*
- 4  *$I(\alpha, \beta) \setminus \text{cl}(A_0 \cup A_1) \neq \emptyset$  for each  $\alpha \in A_0$  and  $\beta \in A_1$ .*

# Classical separation property

Theorem (Kowalewski and Maliszewski 2008; Szczuka 2014)

*Let  $A_0, A_1 \subset \mathbb{R}$  be disjoint. The following conditions are equivalent:*

- ❶ *there is an extra strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$ ,*
- ❷ *there is a strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$ ,*
- ❸ *there is a Darboux quasi-continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$ ,*
- ❹  *$I(\alpha, \beta) \setminus \text{cl}(A_0 \cup A_1) \neq \emptyset$  for each  $\alpha \in A_0$  and  $\beta \in A_1$ .*

# Classical separation property

Theorem (Kowalewski and Maliszewski 2008; Szczuka 2014)

*Let  $A_0, A_1 \subset \mathbb{R}$  be disjoint. The following conditions are equivalent:*

- ❶ *there is an extra strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$ ,*
- ❷ *there is a strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$ ,*
- ❸ *there is a Darboux quasi-continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$ ,*
- ❹  *$I(\alpha, \beta) \setminus \text{cl}(A_0 \cup A_1) \neq \emptyset$  for each  $\alpha \in A_0$  and  $\beta \in A_1$ .*

# Classical separation property

Theorem (Kowalewski and Maliszewski 2008; Szczuka 2014)

*Let  $A_0, A_1 \subset \mathbb{R}$  be disjoint. The following conditions are equivalent:*

- ❶ *there is an extra strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$ ,*
- ❷ *there is a strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$ ,*
- ❸ *there is a Darboux quasi-continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$ ,*
- ❹  *$I(\alpha, \beta) \setminus \text{cl}(A_0 \cup A_1) \neq \emptyset$  for each  $\alpha \in A_0$  and  $\beta \in A_1$ .*

# Classical separation property

Theorem (Kowalewski and Maliszewski 2008; Szczuka 2014)

*Let  $A_0, A_1 \subset \mathbb{R}$  be disjoint. The following conditions are equivalent:*

- ❶ *there is an extra strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$ ,*
- ❷ *there is a strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$ ,*
- ❸ *there is a Darboux quasi-continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 \subset [f = 0]$  and  $A_1 \subset [f = 1]$ ,*
- ❹  *$I(\alpha, \beta) \setminus \text{cl}(A_0 \cup A_1) \neq \emptyset$  for each  $\alpha \in A_0$  and  $\beta \in A_1$ .*

# Classical separation property

## Theorem (Szczuka 2014)

*Let  $A_0, A_1 \subset \mathbb{R}$  be disjoint. There is an extra strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 = [f = 0]$  and  $A_1 = [f = 1]$  if and only if the following conditions are true:*

- 1  $A_0$  and  $A_1$  are semi-closed,
- 2  $\mathbb{R} \setminus (A_0 \cup A_1)$  is bilaterally dense in itself,
- 3  $I(\alpha, \beta) \setminus \text{cl}(A_0 \cup A_1) \neq \emptyset$  for each  $\alpha \in A_0$  and  $\beta \in A_1$ ,
- 4 for  $i \in \{0, 1\}$  there is a  $G_\delta$ -set  $B_i \subset A_i$  such that  $\text{cl } A_{1-i} \cap B_i = \emptyset$  and  $I[x, t) \cap B_i \neq \emptyset$  for each  $x \in A_i$  and  $t \in \mathbb{R}$ .

# Classical separation property

## Theorem (Szczuka 2014)

*Let  $A_0, A_1 \subset \mathbb{R}$  be disjoint. There is an extra strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 = [f = 0]$  and  $A_1 = [f = 1]$  if and only if the following conditions are true:*

- 1  $A_0$  and  $A_1$  are semi-closed,
- 2  $\mathbb{R} \setminus (A_0 \cup A_1)$  is bilaterally dense in itself,
- 3  $I(\alpha, \beta) \setminus \text{cl}(A_0 \cup A_1) \neq \emptyset$  for each  $\alpha \in A_0$  and  $\beta \in A_1$ ,
- 4 for  $i \in \{0, 1\}$  there is a  $G_\delta$ -set  $B_i \subset A_i$  such that  $\text{cl } A_{1-i} \cap B_i = \emptyset$  and  $I[x, t) \cap B_i \neq \emptyset$  for each  $x \in A_i$  and  $t \in \mathbb{R}$ .

# Classical separation property

## Theorem (Szczuka 2014)

*Let  $A_0, A_1 \subset \mathbb{R}$  be disjoint. There is an extra strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 = [f = 0]$  and  $A_1 = [f = 1]$  if and only if the following conditions are true:*

- 1  $A_0$  and  $A_1$  are semi-closed,
- 2  $\mathbb{R} \setminus (A_0 \cup A_1)$  is bilaterally dense in itself,
- 3  $I(\alpha, \beta) \setminus \text{cl}(A_0 \cup A_1) \neq \emptyset$  for each  $\alpha \in A_0$  and  $\beta \in A_1$ ,
- 4 for  $i \in \{0, 1\}$  there is a  $G_\delta$ -set  $B_i \subset A_i$  such that  $\text{cl } A_{1-i} \cap B_i = \emptyset$  and  $I[x, t) \cap B_i \neq \emptyset$  for each  $x \in A_i$  and  $t \in \mathbb{R}$ .



# Classical separation property

## Theorem (Szczuka 2014)

*Let  $A_0, A_1 \subset \mathbb{R}$  be disjoint. There is an extra strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 = [f = 0]$  and  $A_1 = [f = 1]$  if and only if the following conditions are true:*

- 1  $A_0$  and  $A_1$  are semi-closed,
- 2  $\mathbb{R} \setminus (A_0 \cup A_1)$  is bilaterally dense in itself,
- 3  $I(\alpha, \beta) \setminus \text{cl}(A_0 \cup A_1) \neq \emptyset$  for each  $\alpha \in A_0$  and  $\beta \in A_1$ ,
- 4 for  $i \in \{0, 1\}$  there is a  $G_\delta$ -set  $B_i \subset A_i$  such that  $\text{cl } A_{1-i} \cap B_i = \emptyset$  and  $I[x, t) \cap B_i \neq \emptyset$  for each  $x \in A_i$  and  $t \in \mathbb{R}$ .

# Classical separation property

## Theorem (Szczuka 2014)

*Let  $A_0, A_1 \subset \mathbb{R}$  be disjoint. There is an extra strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A_0 = [f = 0]$  and  $A_1 = [f = 1]$  if and only if the following conditions are true:*

- 1  $A_0$  and  $A_1$  are semi-closed,
- 2  $\mathbb{R} \setminus (A_0 \cup A_1)$  is bilaterally dense in itself,
- 3  $I(\alpha, \beta) \setminus \text{cl}(A_0 \cup A_1) \neq \emptyset$  for each  $\alpha \in A_0$  and  $\beta \in A_1$ ,
- 4 for  $i \in \{0, 1\}$  there is a  $G_\delta$ -set  $B_i \subset A_i$  such that  $\text{cl } A_{1-i} \cap B_i = \emptyset$  and  $I[x, t) \cap B_i \neq \emptyset$  for each  $x \in A_i$  and  $t \in \mathbb{R}$ .

# New separation property

## Theorem

*Let  $X$  be a perfectly normal topological space and sets  $A^+, A^- \subset X$  be disjoint. There is a continuous function  $f: X \rightarrow \mathbb{R}$  such that  $A^+ = [f > 0]$  and  $A^- = [f < 0]$  if and only if sets  $A^+$  and  $A^-$  are open.*

## Theorem

*Let  $X$  be a perfectly normal topological space and sets  $A^+, A^- \subset X$ . There is a continuous function  $f: X \rightarrow \mathbb{R}$  such that  $A^+ \subset [f > 0]$  and  $A^- \subset [f < 0]$  if and only if  $\text{cl } A^+ \cap A^- = \text{cl } A^- \cap A^+ = \emptyset$ .*

# New separation property

## Theorem

*Let  $X$  be a perfectly normal topological space and sets  $A^+, A^- \subset X$  be disjoint. There is a continuous function  $f: X \rightarrow \mathbb{R}$  such that  $A^+ = [f > 0]$  and  $A^- = [f < 0]$  if and only if sets  $A^+$  and  $A^-$  are open.*

## Theorem

*Let  $X$  be a perfectly normal topological space and sets  $A^+, A^- \subset X$ . There is a continuous function  $f: X \rightarrow \mathbb{R}$  such that  $A^+ \subset [f > 0]$  and  $A^- \subset [f < 0]$  if and only if  $\text{cl } A^+ \cap A^- = \text{cl } A^- \cap A^+ = \emptyset$ .*

# New separation property

## Theorem (Maliszewski 2002)

*Let  $A^+, A^- \subset \mathbb{R}$  be disjoint. There is a Darboux function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A^+ = [f > 0]$  and  $A^- = [f < 0]$  if and only if sets  $A^+$  and  $A^-$  are bilaterally  $\mathfrak{c}$ -dense in themselves and  $I[\alpha, \beta] \setminus (A^- \cup A^+) \neq \emptyset$  for each  $\alpha \in A^-$  and  $\beta \in A^+$ .*

## Theorem (Maliszewski 2002)

*Let  $A^+, A^- \subset \mathbb{R}$  be disjoint. There is a Darboux function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A^+ \subset [f > 0]$  and  $A^- \subset [f < 0]$  if and only if every point from  $A^+$  is bilateral  $\mathfrak{c}$ -limit point of  $\mathbb{R} \setminus A^-$ , every point from  $A^-$  is bilateral  $\mathfrak{c}$ -limit point of  $\mathbb{R} \setminus A^+$  and  $I[\alpha, \beta] \setminus (A^- \cup A^+) \neq \emptyset$  for each  $\alpha \in A^-$  and  $\beta \in A^+$ .*

# New separation property

## Theorem (Maliszewski 2002)

*Let  $A^+, A^- \subset \mathbb{R}$  be disjoint. There is a Darboux function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A^+ = [f > 0]$  and  $A^- = [f < 0]$  if and only if sets  $A^+$  and  $A^-$  are bilaterally  $\mathfrak{c}$ -dense in themselves and  $I[\alpha, \beta] \setminus (A^- \cup A^+) \neq \emptyset$  for each  $\alpha \in A^-$  and  $\beta \in A^+$ .*

## Theorem (Maliszewski 2002)

*Let  $A^+, A^- \subset \mathbb{R}$  be disjoint. There is a Darboux function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A^+ \subset [f > 0]$  and  $A^- \subset [f < 0]$  if and only if every point from  $A^+$  is bilateral  $\mathfrak{c}$ -limit point of  $\mathbb{R} \setminus A^-$ , every point from  $A^-$  is bilateral  $\mathfrak{c}$ -limit point of  $\mathbb{R} \setminus A^+$  and  $I[\alpha, \beta] \setminus (A^- \cup A^+) \neq \emptyset$  for each  $\alpha \in A^-$  and  $\beta \in A^+$ .*

# New separation property

## Theorem (Kowalewski 2005)

*Let  $A^+, A^- \subset \mathbb{R}$  be disjoint. There is a quasi-continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A^+ = [f > 0]$  and  $A^- = [f < 0]$  if and only if sets  $A^+$  and  $A^-$  are semi-open.*

## Theorem (Kowalewski 2005)

*Let  $A^+, A^- \subset \mathbb{R}$ . There is a quasi-continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A^+ \subset [f > 0]$  and  $A^- \subset [f < 0]$  if and only if  $A^+ \cap A^- = \text{int cl } A^+ \cap A^- = A^+ \cap \text{int cl } A^- = \emptyset$ .*

# New separation property

## Theorem (Kowalewski 2005)

*Let  $A^+, A^- \subset \mathbb{R}$  be disjoint. There is a quasi-continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A^+ = [f > 0]$  and  $A^- = [f < 0]$  if and only if sets  $A^+$  and  $A^-$  are semi-open.*

## Theorem (Kowalewski 2005)

*Let  $A^+, A^- \subset \mathbb{R}$ . There is a quasi-continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A^+ \subset [f > 0]$  and  $A^- \subset [f < 0]$  if and only if  $A^+ \cap A^- = \text{int cl } A^+ \cap A^- = A^+ \cap \text{int cl } A^- = \emptyset$ .*



# New separation property

## Theorem (Kowalewski and Maliszewski 2008)

*Let  $A^+, A^- \subset \mathbb{R}$  be disjoint. There is a Darboux quasi-continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A^+ = [f > 0]$  and  $A^- = [f < 0]$  if and only if sets  $A^+$  and  $A^-$  are semi-open and bilaterally dense in themselves and  $I[\alpha, \beta] \setminus (A^- \cup A^+) \neq \emptyset$  for each  $\alpha \in A^-$  and  $\beta \in A^+$ .*

## Theorem (Kowalewski and Maliszewski 2008)

*Let  $A^+, A^- \subset \mathbb{R}$  be disjoint. There is a Darboux quasi-continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A^+ \subset [f > 0]$  and  $A^- \subset [f < 0]$  if and only if  $I[\alpha, \beta] \setminus (A^- \cup A^+) \neq \emptyset$  and  $I(\alpha, \beta) \setminus \text{cl } A^- \neq \emptyset \neq I(\alpha, \beta) \setminus \text{cl } A^+$  for each  $\alpha \in A^-$  and  $\beta \in A^+$ .*

# New separation property

## Theorem (Kowalewski and Maliszewski 2008)

*Let  $A^+, A^- \subset \mathbb{R}$  be disjoint. There is a Darboux quasi-continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A^+ = [f > 0]$  and  $A^- = [f < 0]$  if and only if sets  $A^+$  and  $A^-$  are semi-open and bilaterally dense in themselves and  $I[\alpha, \beta] \setminus (A^- \cup A^+) \neq \emptyset$  for each  $\alpha \in A^-$  and  $\beta \in A^+$ .*

## Theorem (Kowalewski and Maliszewski 2008)

*Let  $A^+, A^- \subset \mathbb{R}$  be disjoint. There is a Darboux quasi-continuous function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A^+ \subset [f > 0]$  and  $A^- \subset [f < 0]$  if and only if  $I[\alpha, \beta] \setminus (A^- \cup A^+) \neq \emptyset$  and  $I(\alpha, \beta) \setminus \text{cl } A^- \neq \emptyset \neq I(\alpha, \beta) \setminus \text{cl } A^+$  for each  $\alpha \in A^-$  and  $\beta \in A^+$ .*

# New separation property

## Theorem (Kowalewski 2010)

*Let  $A^+, A^- \subset \mathbb{R}$  be disjoint. There is a strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A^+ = [f > 0]$  and  $A^- = [f < 0]$  if and only if sets  $A^+$  and  $A^-$  are semi-open and bilaterally dense in themselves and there is a  $G_\delta$ -set  $D \subset \mathbb{R} \setminus (A^+ \cup A^-)$  such that  $I(\alpha, \beta) \cap D \neq \emptyset$  for each  $\alpha \in A^-$  and  $\beta \in A^+$ .*

## Theorem (Kowalewski 2010)

*Let  $A^+, A^- \subset \mathbb{R}$  be disjoint. There is a strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A^+ \subset [f > 0]$  and  $A^- \subset [f < 0]$  if and only if  $I(\alpha, \beta) \setminus \text{cl } A^- \neq \emptyset \neq I(\alpha, \beta) \setminus \text{cl } A^+$  for each  $\alpha \in A^-$  and  $\beta \in A^+$  and there is a  $G_\delta$ -set  $D \subset \mathbb{R} \setminus (A^+ \cup A^-)$  such that  $I(\alpha, \beta) \cap D \neq \emptyset$  for each  $\alpha \in A^-$  and  $\beta \in A^+$ .*

# New separation property

## Theorem (Kowalewski 2010)

*Let  $A^+, A^- \subset \mathbb{R}$  be disjoint. There is a strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A^+ = [f > 0]$  and  $A^- = [f < 0]$  if and only if sets  $A^+$  and  $A^-$  are semi-open and bilaterally dense in themselves and there is a  $G_\delta$ -set  $D \subset \mathbb{R} \setminus (A^+ \cup A^-)$  such that  $I(\alpha, \beta) \cap D \neq \emptyset$  for each  $\alpha \in A^-$  and  $\beta \in A^+$ .*

## Theorem (Kowalewski 2010)

*Let  $A^+, A^- \subset \mathbb{R}$  be disjoint. There is a strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A^+ \subset [f > 0]$  and  $A^- \subset [f < 0]$  if and only if  $I(\alpha, \beta) \setminus \text{cl } A^- \neq \emptyset \neq I(\alpha, \beta) \setminus \text{cl } A^+$  for each  $\alpha \in A^-$  and  $\beta \in A^+$  and there is a  $G_\delta$ -set  $D \subset \mathbb{R} \setminus (A^+ \cup A^-)$  such that  $I(\alpha, \beta) \cap D \neq \emptyset$  for each  $\alpha \in A^-$  and  $\beta \in A^+$ .*

# New separation property

## Problem

Characterize a pair  $(A^+, A^-)$  of disjoint sets for which there is an extra strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A^+ = [f > 0]$  and  $A^- = [f < 0]$ .

## Problem

Characterize a pair  $(A^+, A^-)$  of disjoint sets for which there is an extra strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A^+ \subset [f > 0]$  and  $A^- \subset [f < 0]$ .

# New separation property





## Problem

Characterize a pair  $(A^+, A^-)$  of disjoint sets for which there is an extra strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A^+ = [f > 0]$  and  $A^- = [f < 0]$ .





## Problem

Characterize a pair  $(A^+, A^-)$  of disjoint sets for which there is an extra strong Świątkowski function  $f: \mathbb{R} \rightarrow \mathbb{R}$  such that  $A^+ \subset [f > 0]$  and  $A^- \subset [f < 0]$ .

# The bibliography





-  S. Kempisty, *Sur les fonctions quasicontinues*, Fund. Math. **19** (1932), 184–197.
-  M. Kowalewski, *Separating sets by quasi-continuous functions*, Atti Semin. Mat. Fis. Univ. Modena Reggio Emilia, **53** (2005), 299–303.
-  M. Kowalewski, *Separating sets by quasi-continuous functions*, Atti Semin. Mat. Fis. Univ. Modena Reggio Emilia, **53** (2005), 299–303.
-  M. Kowalewski, A. Maliszewski, *Separating sets by Darboux quasi-continuous functions*, Topology Appl., **155** (2008), 1990–1995.

# The bibliography





-  S. Kempisty, *Sur les fonctions quasicontinues*, Fund. Math. **19** (1932), 184–197.
-  M. Kowalewski, *Separating sets by quasi-continuous functions*, Atti Semin. Mat. Fis. Univ. Modena Reggio Emilia, **53** (2005), 299–303.
-  M. Kowalewski, *Separating sets by quasi-continuous functions*, Atti Semin. Mat. Fis. Univ. Modena Reggio Emilia, **53** (2005), 299–303.
-  M. Kowalewski, A. Maliszewski, *Separating sets by Darboux quasi-continuous functions*, Topology Appl., **155** (2008), 1990–1995.








# The bibliography

-  S. Kempisty, *Sur les fonctions quasicontinues*, Fund. Math. **19** (1932), 184–197.
-  M. Kowalewski, *Separating sets by quasi-continuous functions*, Atti Semin. Mat. Fis. Univ. Modena Reggio Emilia, **53** (2005), 299–303.
-  M. Kowalewski, *Separating sets by quasi-continuous functions*, Atti Semin. Mat. Fis. Univ. Modena Reggio Emilia, **53** (2005), 299–303.
-  M. Kowalewski, A. Maliszewski, *Separating sets by Darboux quasi-continuous functions*, Topology Appl., **155** (2008), 1990–1995.






# The bibliography

-  S. Kempisty, *Sur les fonctions quasicontinues*, Fund. Math. **19** (1932), 184–197.
-  M. Kowalewski, *Separating sets by quasi-continuous functions*, Atti Semin. Mat. Fis. Univ. Modena Reggio Emilia, **53** (2005), 299–303.
-  M. Kowalewski, *Separating sets by quasi-continuous functions*, Atti Semin. Mat. Fis. Univ. Modena Reggio Emilia, **53** (2005), 299–303.
-  M. Kowalewski, A. Maliszewski, *Separating sets by Darboux quasi-continuous functions*, Topology Appl., **155** (2008), 1990–1995.






# The bibliography

-  M. Kowalewski, *Rozdzielanie zbiorów przy użyciu funkcji z zadanej klasy*, PhD thesis, Łódź 2010.
-  N. Levine, *Semi-open sets and semi-continuity in topological spaces*, Amer. Math. Monthly, **70** (1963), 36–41.
-  A. Maliszewski, *On the limits of Świątkowski functions*, Zeszyty Nauk. Politech. Łódz. Mat., **27(719)** (1995), 87–93.
-  A. Maliszewski, *Separating sets by Darboux-like functions*, Fund. Math. **175** (2002), 271–283.
-  P. Szczuka, *Sums and products of extra strong Świątkowski functions*, Tatra Mt. Math. Publ., **49** (2011), 71–79.






# The bibliography

-  M. Kowalewski, *Rozdzielanie zbiorów przy użyciu funkcji z zadanej klasy*, PhD thesis, Łódź 2010.
-  N. Levine, *Semi-open sets and semi-continuity in topological spaces*, Amer. Math. Monthly, **70** (1963), 36–41.
-  A. Maliszewski, *On the limits of Świątkowski functions*, Zeszyty Nauk. Politech. Łódz. Mat., **27(719)** (1995), 87–93.
-  A. Maliszewski, *Separating sets by Darboux-like functions*, Fund. Math. **175** (2002), 271–283.
-  P. Szczuka, *Sums and products of extra strong Świątkowski functions*, Tatra Mt. Math. Publ., **49** (2011), 71–79.






# The bibliography

-  M. Kowalewski, *Rozdzielanie zbiorów przy użyciu funkcji z zadanej klasy*, PhD thesis, Łódź 2010.
-  N. Levine, *Semi-open sets and semi-continuity in topological spaces*, Amer. Math. Monthly, **70** (1963), 36–41.
-  A. Maliszewski, *On the limits of Świątkowski functions*, Zeszyty Nauk. Politech. Łódz. Mat., **27(719)** (1995), 87–93.
-  A. Maliszewski, *Separating sets by Darboux-like functions*, Fund. Math. **175** (2002), 271–283.
-  P. Szczuka, *Sums and products of extra strong Świątkowski functions*, Tatra Mt. Math. Publ., **49** (2011), 71–79.





# The bibliography

-  M. Kowalewski, *Rozdzielanie zbiorów przy użyciu funkcji z zadanej klasy*, PhD thesis, Łódź 2010.
-  N. Levine, *Semi-open sets and semi-continuity in topological spaces*, Amer. Math. Monthly, **70** (1963), 36–41.
-  A. Maliszewski, *On the limits of Świątkowski functions*, Zeszyty Nauk. Politech. Łódz. Mat., **27(719)** (1995), 87–93.
-  A. Maliszewski, *Separating sets by Darboux-like functions*, Fund. Math. **175** (2002), 271–283.
-  P. Szczuka, *Sums and products of extra strong Świątkowski functions*, Tatra Mt. Math. Publ., **49** (2011), 71–79.

# The bibliography





-  M. Kowalewski, *Rozdzielanie zbiorów przy użyciu funkcji z zadanej klasy*, PhD thesis, Łódź 2010.
-  N. Levine, *Semi-open sets and semi-continuity in topological spaces*, Amer. Math. Monthly, **70** (1963), 36–41.
-  A. Maliszewski, *On the limits of Świątkowski functions*, Zeszyty Nauk. Politech. Łódz. Mat., **27(719)** (1995), 87–93.
-  A. Maliszewski, *Separating sets by Darboux-like functions*, Fund. Math. **175** (2002), 271–283.
-  P. Szczuka, *Sums and products of extra strong Świątkowski functions*, Tatra Mt. Math. Publ., **49** (2011), 71–79.

# The bibliography





-  P. Szczuka, *Generalizations of Urysohn's lemma for some subclasses of Darboux functions* - preprint.
-  P. Szczuka, *Generalizations of Urysohn's lemma for the family of extra strong Świątkowski functions* - preprint.
-  H. P. Thielman, *Types of functions*, Amer. Math. Monthly **60** (1953), no. 3, 156–161.
-  P. Urysohn, *Über die Mächtigkeit der zusammenhängenden Mengen*, Math. Ann. **94** (1925), no. 1, 262–295.







# The bibliography

-  P. Szczuka, *Generalizations of Urysohn's lemma for some subclasses of Darboux functions* - preprint.
-  P. Szczuka, *Generalizations of Urysohn's lemma for the family of extra strong Świątkowski functions* - preprint.
-  H. P. Thielman, *Types of functions*, Amer. Math. Monthly **60** (1953), no. 3, 156–161.
-  P. Urysohn, *Über die Mächtigkeit der zusammenhängenden Mengen*, Math. Ann. **94** (1925), no. 1, 262–295.

# The bibliography

-  P. Szczuka, *Generalizations of Urysohn's lemma for some subclasses of Darboux functions* - preprint.
-  P. Szczuka, *Generalizations of Urysohn's lemma for the family of extra strong Świątkowski functions* - preprint.
-  H. P. Thielman, *Types of functions*, Amer. Math. Monthly **60** (1953), no. 3, 156–161.
-  P. Urysohn, *Über die Mächtigkeit der zusammenhängenden Mengen*, Math. Ann. **94** (1925), no. 1, 262–295.

# The bibliography

-  P. Szczuka, *Generalizations of Urysohn's lemma for some subclasses of Darboux functions* - preprint.
-  P. Szczuka, *Generalizations of Urysohn's lemma for the family of extra strong Świątkowski functions* - preprint.
-  H. P. Thielman, *Types of functions*, Amer. Math. Monthly **60** (1953), no. 3, 156–161.
-  P. Urysohn, *Über die Mächtigkeit der zusammenhängenden Mengen*, Math. Ann. **94** (1925), no. 1, 262–295.