

On M -sets for series with respect to characters of compact zero-dimensional groups

Valentin Skvortsov

Institute of Mathematics, Casimirus the Great University,
Poland

Department of Mathematics, Moscow State University,
Russia

28th INTERNATIONAL SUMMER
CONFERENCE ON REAL FUNCTIONS
THEORY August 31 – September 5, 2014,
Stara Lesna

U -set and M -set

A set E is said to be U -set for a system of functions (especially for orthogonal system) if the convergence of a series with respect to this system to zero outside the set E implies that all coefficients of the series are zero.

U -set and M -set

A set E is said to be U -set for a system of functions (especially for orthogonal system) if the convergence of a series with respect to this system to zero outside the set E implies that all coefficients of the series are zero.

Otherwise the set is called M -set, i.e., a set E is said to be M -set for a system of functions if there exists a non-trivial series with respect to this system which converges to zero outside the set E .

U-set and M-set

A set E is said to be **U-set** for a system of functions (especially for orthogonal system) if the convergence of a series with respect to this system to zero outside the set E implies that all coefficients of the series are zero.

Otherwise the set is called **M-set**, i.e., a set E is said to be M -set for a system of functions if there exists a non-trivial series with respect to this system which converges to zero outside the set E .

This series is called **null-series** if $\mu E = 0$.

U -set and M -set

A set E is said to be U -set for a system of functions (especially for orthogonal system) if the convergence of a series with respect to this system to zero outside the set E implies that all coefficients of the series are zero.

Otherwise the set is called M -set, i.e., a set E is said to be M -set for a system of functions if there exists a non-trivial series with respect to this system which converges to zero outside the set E .

This series is called $null$ -series if $\mu E = 0$.

A set is called M_0 -set, or a set of $strict\ multiplicity$, if it is an M -set and the corresponding null-series is a Fourier-Stieltjes series of a probability measure concentrated on E .

U -set and M -set

A set E is said to be U -set for a system of functions (especially for orthogonal system) if the convergence of a series with respect to this system to zero outside the set E implies that all coefficients of the series are zero.

Otherwise the set is called M -set, i.e., a set E is said to be M -set for a system of functions if there exists a non-trivial series with respect to this system which converges to zero outside the set E .

This series is called $null$ -series if $\mu E = 0$.

A set is called M_0 -set, or a set of $strict$ $multiplicity$, if it is an M -set and the corresponding null-series is a Fourier-Stieltjes series of a probability measure concentrated on E .

Survey: A. S. Kechris, A. Louveau , Descriptive set theory and the structure of sets of uniqueness, London Mathematical Society lecture series 128, Cambridge University Press, 1987.

Zero-dimensional compact abelian group

Let G be a zero-dimensional compact abelian group.

Zero-dimensional compact abelian group

Let G be a **zero-dimensional compact abelian group**.

Topology in such a group can be given by a chain of subgroups

$$G = G_0 \supset G_1 \supset G_2 \dots \supset G_n \supset \dots$$

with $0 = \bigcap_{n=0}^{+\infty} G_n$.

Zero-dimensional compact abelian group

Let G be a **zero-dimensional compact abelian group**.

Topology in such a group can be given by a chain of subgroups

$$G = G_0 \supset G_1 \supset G_2 \dots \supset G_n \supset \dots$$

with $0 = \bigcap_{n=0}^{+\infty} G_n$.

The subgroups G_n are **clopen** sets with respect to this topology.

Zero-dimensional compact abelian group

Let G be a **zero-dimensional compact abelian group**.

Topology in such a group can be given by a chain of subgroups

$$G = G_0 \supset G_1 \supset G_2 \dots \supset G_n \supset \dots$$

with $0 = \bigcap_{n=0}^{+\infty} G_n$.

The subgroups G_n are **clopen** sets with respect to this topology.

As G is compact the factor group G_n/G_{n+1} is finite for each n . Let its order be p_n .

Zero-dimensional compact abelian group

Let G be a **zero-dimensional compact abelian group**.

Topology in such a group can be given by a chain of subgroups

$$G = G_0 \supset G_1 \supset G_2 \dots \supset G_n \supset \dots$$

with $0 = \bigcap_{n=0}^{+\infty} G_n$.

The subgroups G_n are **clopen** sets with respect to this topology.

As G is compact the factor group G_n/G_{n+1} is finite for each n . Let its order be p_n .

Then the order of G_0/G_1 is p_0 , and the order of G_0/G_n , $n = 1, 2, \dots$, is

$$m_n := p_0 \cdot p_1 \cdot \dots \cdot p_{n-1},$$

with $p_i \geq 2$ for all i (put $m_0 := 1$).

Zero-dimensional compact abelian group

Denote by K_n any coset of the subgroup G_n and by $K_n(g)$ the coset of the subgroup G_n which contains the element g , i.e.,

$$K_n(g) = g + G_n.$$

Zero-dimensional compact abelian group

Denote by K_n any coset of the subgroup G_n and by $K_n(g)$ the coset of the subgroup G_n which contains the element g , i.e.,

$$K_n(g) = g + G_n.$$

For each $g \in G$ the sequence $\{K_n(g)\}$ is decreasing and

$$\{g\} = \bigcap_n K_n(g).$$

Zero-dimensional compact abelian group

Denote by K_n any coset of the subgroup G_n and by $K_n(g)$ the coset of the subgroup G_n which contains the element g , i.e.,

$$K_n(g) = g + G_n.$$

For each $g \in G$ the sequence $\{K_n(g)\}$ is decreasing and

$$\{g\} = \bigcap_n K_n(g).$$

Examples

1. Dyadic Cantor group G

Examples

1. Dyadic Cantor group G

The group of sequences of integers of the form $\tilde{x} = \{x_j\}_{j=0}^{\infty}$ where $0 \leq x_j \leq 1$ with group operation defined as component-wise addition modulo 2 for each component.

Examples

1. Dyadic Cantor group G

The group of sequences of integers of the form $\tilde{x} = \{x_j\}_{j=0}^{\infty}$ where $0 \leq x_j \leq 1$ with group operation defined as component-wise addition modulo 2 for each component.

Topology is defined by subgroups

$$G_n = \{\tilde{x} \in G : x_j = 0 \text{ for } j = 0, \dots, n-1\}$$

Examples

1. Dyadic Cantor group G

The group of sequences of integers of the form $\tilde{x} = \{x_j\}_{j=0}^{\infty}$ where $0 \leq x_j \leq 1$ with group operation defined as component-wise addition modulo 2 for each component.

Topology is defined by subgroups

$$G_n = \{\tilde{x} \in G : x_j = 0 \text{ for } j = 0, \dots, n-1\}$$

2. Vilenkin group G is defined by a sequence of prime numbers $\{p_j\}$, $p_j \geq 2$

Examples

1. Dyadic Cantor group G

The group of sequences of integers of the form $\tilde{x} = \{x_j\}_{j=0}^{\infty}$ where $0 \leq x_j \leq 1$ with group operation defined as component-wise addition modulo 2 for each component.

Topology is defined by subgroups

$$G_n = \{\tilde{x} \in G : x_j = 0 \text{ for } j = 0, \dots, n-1\}$$

2. Vilenkin group G is defined by a sequence of prime numbers $\{p_j\}$, $p_j \geq 2$

Then G is the group of sequences of integers of the form $\tilde{x} = \{x_j\}_{j=0}^{\infty}$ where $0 \leq x_j \leq p_j$ with group operation defined as component-wise addition modulo p_j for j th component.

Examples

1. Dyadic Cantor group G

The group of sequences of integers of the form $\tilde{x} = \{x_j\}_{j=0}^{\infty}$ where $0 \leq x_j \leq 1$ with group operation defined as component-wise addition modulo 2 for each component.

Topology is defined by subgroups

$$G_n = \{\tilde{x} \in G : x_j = 0 \text{ for } j = 0, \dots, n-1\}$$

2. Vilenkin group G is defined by a sequence of prime numbers $\{p_j\}$, $p_j \geq 2$

Then G is the group of sequences of integers of the form $\tilde{x} = \{x_j\}_{j=0}^{\infty}$ where $0 \leq x_j \leq p_j$ with group operation defined as component-wise addition modulo p_j for j th component.

Topology is defined by subgroups

$$G_n = \{\tilde{x} \in G : x_j = 0 \text{ for } j = 0, \dots, n-1\}$$

Examples

3. An important example - group of p -adic integers.

Examples

3. An important example - **group of p -adic integers**.

The additive group of p -adic integers (with p being a fixed prime number) is the set of formal sums:

$$\sum_{j=0}^{\infty} x_j p^j$$

where $0 \leq x_i \leq p - 1$ and the addition is done with **carries**,

Examples

3. An important example - group of p -adic integers.

The additive group of p -adic integers (with p being a fixed prime number) is the set of formal sums:

$$\sum_{j=0}^{\infty} x_j p^j$$

where $0 \leq x_i \leq p-1$ and the addition is done with carries,

i.e., to add two sequences $\sum_{j=0}^{\infty} x_j p^j$ and $\sum_{j=0}^{\infty} y_j p^j$, we add coordinate-wise and if any of the sums is p or more, we take a carry of 1 to the sum in next coordinate.

Examples

3. An important example - group of p -adic integers.

The additive group of p -adic integers (with p being a fixed prime number) is the set of formal sums:

$$\sum_{j=0}^{\infty} x_j p^j$$

where $0 \leq x_i \leq p-1$ and the addition is done with carries,

i.e., to add two sequences $\sum_{j=0}^{\infty} x_j p^j$ and $\sum_{j=0}^{\infty} y_j p^j$, we add coordinate-wise and if any of the sums is p or more, we take a carry of 1 to the sum in next coordinate.

Topology is given by subgroups

$$G_n = \left\{ \sum_{j=0}^{\infty} x_j p^j : x_j = 0 \text{ for } j = 1, \dots, n-1 \right\}.$$

Group of characters

Let Γ denotes the dual group of G , i.e., the **group of characters** of the group G .

Group of characters

Let Γ denotes the dual group of G , i.e., the **group of characters** of the group G .

The group Γ is discrete and it can be represented as a sum of increasing chain of finite subgroups

$$\Gamma_0 \subset \Gamma_1 \subset \Gamma_2 \subset \dots \subset \Gamma_n \subset \dots$$

where $\Gamma_0 = \{\gamma^{(0)}\}$ with $\gamma^{(0)}(g) = 1$ for all $g \in G$.

Group of characters

Let Γ denotes the dual group of G , i.e., the **group of characters** of the group G .

The group Γ is discrete and it can be represented as a sum of increasing chain of finite subgroups

$$\Gamma_0 \subset \Gamma_1 \subset \Gamma_2 \subset \dots \subset \Gamma_n \subset \dots$$

where $\Gamma_0 = \{\gamma^{(0)}\}$ with $\gamma^{(0)}(g) = 1$ for all $g \in G$.

For each n the group Γ_n is the **annihilator** of G_n , i.e.,

$$\Gamma_n = G_n^\perp := \{\gamma \in \Gamma : (\gamma(g) = 1 \text{ for all } g \in G_n)\}.$$

Group of characters

The characters γ constitute a countable **orthogonal system** on G with respect to normalized Haar measure μ_G and we can consider a series

$$\sum_{\gamma \in \Gamma} a_{\gamma} \gamma$$

with respect to this system.

Group of characters

The characters γ constitute a countable **orthogonal system** on G with respect to normalized Haar measure μ_G and we can consider a series

$$\sum_{\gamma \in \Gamma} a_{\gamma} \gamma$$

with respect to this system.

Important subsequence of partial sums of this series are

$$\sum_{\gamma \in \Gamma_n} a_{\gamma} \gamma.$$

Properties of the characters

For each coset K_n of G_n choose and fix an element g_{K_n} . Then for each $n \in \mathbb{Z}$ we can represent any element $g \in G$ in the form:

$$g = g_{K_n} + \{g\}_n$$

where $\{g\}_n \in G_n$.

Properties of the characters

For each coset K_n of G_n choose and fix an element g_{K_n} . Then for each $n \in \mathbb{Z}$ we can represent any element $g \in G$ in the form:

$$g = g_{K_n} + \{g\}_n$$

where $\{g\}_n \in G_n$.

This representation, properties of a character and the definition of the annihilator imply

$$\gamma(g) = \gamma(g_{K_n})\gamma(\{g\}_n) = \gamma(g_{K_n}).$$

Properties of the characters

For each coset K_n of G_n choose and fix an element g_{K_n} . Then for each $n \in \mathbb{Z}$ we can represent any element $g \in G$ in the form:

$$g = g_{K_n} + \{g\}_n$$

where $\{g\}_n \in G_n$.

This representation, properties of a character and the definition of the annihilator imply

$$\gamma(g) = \gamma(g_{K_n})\gamma(\{g\}_n) = \gamma(g_{K_n}).$$

So with a fixed element g_{K_n} , the value $\gamma(g)$ is constant for all $g \in K_n$.
So we get:

Properties of the characters

For each coset K_n of G_n choose and fix an element g_{K_n} . Then for each $n \in \mathbb{Z}$ we can represent any element $g \in G$ in the form:

$$g = g_{K_n} + \{g\}_n$$

where $\{g\}_n \in G_n$.

This representation, properties of a character and the definition of the annihilator imply

$$\gamma(g) = \gamma(g_{K_n})\gamma(\{g\}_n) = \gamma(g_{K_n}).$$

So with a fixed element g_{K_n} , the value $\gamma(g)$ is constant for all $g \in K_n$.
So we get:

Lemma

If $\gamma \in \Gamma_n$ then γ is constant on each coset K_n of G_n .

Numeration of the characters

The factor groups $\Gamma_{n+1}/\Gamma_n = G_{n+1}^\perp/G_n^\perp$ and G_n/G_{n+1} are isomorphic and so they are of the same order p_n for each $n \in \mathbb{N}$. This implies that the group Γ_n has $m_n = p_0 \cdot p_1 \cdot \dots \cdot p_{n-1}$ elements.

Numeration of the characters

The factor groups $\Gamma_{n+1}/\Gamma_n = G_{n+1}^\perp/G_n^\perp$ and G_n/G_{n+1} are isomorphic and so they are of the same order p_n for each $n \in \mathbb{N}$. This implies that the group Γ_n has $m_n = p_0 \cdot p_1 \cdot \dots \cdot p_{n-1}$ elements.

Put $\gamma_0 = 0$. Pick up a character $\gamma \in \Gamma_{n+1} \setminus \Gamma_n$ for each $n = 0, 1, 2, \dots$ and denote it by γ_{m_n} .

Numeration of the characters

The factor groups $\Gamma_{n+1}/\Gamma_n = G_{n+1}^\perp/G_n^\perp$ and G_n/G_{n+1} are isomorphic and so they are of the same order p_n for each $n \in \mathbb{N}$. This implies that the group Γ_n has $m_n = p_0 \cdot p_1 \cdot \dots \cdot p_{n-1}$ elements.

Put $\gamma_0 = 0$. Pick up a character $\gamma \in \Gamma_{n+1} \setminus \Gamma_n$ for each $n = 0, 1, 2, \dots$ and denote it by γ_{m_n} .

For

$$n = \sum_{k=0}^s t_k m_k, \quad 0 \leq t_k \leq p_k - 1$$

define

$$\gamma_n := \prod_{k=0}^s (\gamma_{m_k})^{t_k}.$$

Now we can consider a usual series in this numeration: $\sum_{i=0}^{\infty} a_i \gamma_i$.

Numeration of the characters

The factor groups $\Gamma_{n+1}/\Gamma_n = G_{n+1}^\perp/G_n^\perp$ and G_n/G_{n+1} are isomorphic and so they are of the same order p_n for each $n \in \mathbb{N}$. This implies that the group Γ_n has $m_n = p_0 \cdot p_1 \cdot \dots \cdot p_{n-1}$ elements.

Put $\gamma_0 = 0$. Pick up a character $\gamma \in \Gamma_{n+1} \setminus \Gamma_n$ for each $n = 0, 1, 2, \dots$ and denote it by γ_{m_n} .

For

$$n = \sum_{k=0}^s t_k m_k, \quad 0 \leq t_k \leq p_k - 1$$

define

$$\gamma_n := \prod_{k=0}^s (\gamma_{m_k})^{t_k}.$$

Now we can consider a usual series in this numeration: $\sum_{i=0}^{\infty} a_i \gamma_i$. In particular

$$\sum_{\gamma \in \Gamma_n} a_\gamma \gamma = S_{m_n} = \sum_{i=0}^{m_n-1} a_i \gamma_i$$

Examples

If G is dyadic Cantor group, then Γ is **Walsh system**.

Examples

If G is dyadic Cantor group, then Γ is **Walsh system**.

For Vilenkin group it is **Price-Vilenkin system**.

Examples

If G is dyadic Cantor group, then Γ is **Walsh system**.

For Vilenkin group it is **Price-Vilenkin system**.

G is the group of p -adic integers with p being a fixed prime number.

Examples

If G is dyadic Cantor group, then Γ is **Walsh system**.

For Vilenkin group it is **Price-Vilenkin system**.

G is the group of p -adic integers with p being a fixed prime number.

At a point

$$\sum_{j=0}^{\infty} x_j p^j$$

put

$$\gamma_{p^n}(x) = \prod_{j=0}^n \exp\left(\frac{2\pi i x_j}{p^{n+1}}\right)$$

.

Examples

If G is dyadic Cantor group, then Γ is **Walsh system**.

For Vilenkin group it is **Price-Vilenkin system**.

G is the group of p -adic integers with p being a fixed prime number.

At a point

$$\sum_{j=0}^{\infty} x_j p^j$$

put

$$\gamma_{p^n}(x) = \prod_{j=0}^n \exp\left(\frac{2\pi i x_j}{p^{n+1}}\right)$$

.

For example for $p = 2$ γ_{2^2} assumes the values

$$1, \quad -1, \quad i, \quad -i, \quad \exp\left(\frac{\pi i}{4}\right), \quad -\exp\left(\frac{\pi i}{4}\right), \quad \exp\left(\frac{3\pi i}{4}\right), \quad -\exp\left(\frac{3\pi i}{4}\right).$$

Examples

If G is dyadic Cantor group, then Γ is **Walsh system**.

For Vilenkin group it is **Price-Vilenkin system**.

G is the group of p -adic integers with p being a fixed prime number.

At a point

$$\sum_{j=0}^{\infty} x_j p^j$$

put

$$\gamma_{p^n}(x) = \prod_{j=0}^n \exp\left(\frac{2\pi i x_j}{p^{n+1}}\right)$$

.

For example for $p = 2$ γ_{2^2} assumes the values

$$1, \quad -1, \quad i, \quad -i, \quad \exp\left(\frac{\pi i}{4}\right), \quad -\exp\left(\frac{\pi i}{4}\right), \quad \exp\left(\frac{3\pi i}{4}\right), \quad -\exp\left(\frac{3\pi i}{4}\right).$$

For $m = \sum_{n=0}^{n_m} t_n p^n$ γ_m is a product of γ_{p^n} .

QUASI-MEASURE

The integral $\int_{K_n} S_{m_n}$ defines an additive function ψ on the family \mathcal{I} of all cosets called **quasi-measure**.

QUASI-MEASURE

The integral $\int_{K_n} S_{m_n}$ defines an additive function ψ on the family \mathcal{I} of all cosets called **quasi-measure**.

Since the sum S_{m_n} is constant on each K_n we get

QUASI-MEASURE

The integral $\int_{K_n} S_{m_n}$ defines an additive function ψ on the family \mathcal{I} of all cosets called **quasi-measure**.

Since the sum S_{m_n} is constant on each K_n we get

$$S_{m_n}(g) = \frac{1}{|K_n|} \int_{K_n} S_{m_n} d\mu = \frac{\psi(K_n)}{|K_n|} \quad (1)$$

for any point $g \in K_n$.

QUASI-MEASURE

The integral $\int_{K_n} S_{m_n}$ defines an additive function ψ on the family \mathcal{I} of all cosets called **quasi-measure**.

Since the sum S_{m_n} is constant on each K_n we get

$$S_{m_n}(g) = \frac{1}{|K_n|} \int_{K_n} S_{m_n} d\mu = \frac{\psi(K_n)}{|K_n|} \quad (1)$$

for any point $g \in K_n$.

So at any point $g \in G$ we get

$$\lim_{n \rightarrow \infty} S_{m_n}(g) = D_B \psi(g).$$

QUASI-MEASURE

Theorem

Any series w.r. to Γ is Fourier-Stieltjes series w.r. to quasi-measure, so that

$$a_\gamma = \hat{\psi}(\gamma) = \int_G \gamma d\psi.$$

A set $E \subset G$ is M -set w.r. to Γ iff there exists a μ -singular quasi-measure ψ supported by E with

$$\lim_{n \rightarrow \infty} \hat{\psi}(n) = 0.$$

QUASI-MEASURE

Theorem

Any series w.r. to Γ is Fourier-Stieltjes series w.r. to quasi-measure, so that

$$a_\gamma = \hat{\psi}(\gamma) = \int_G \gamma d\psi.$$

A set $E \subset G$ is M -set w.r. to Γ iff there exists a μ -singular quasi-measure ψ supported by E with

$$\lim_{n \rightarrow \infty} \hat{\psi}(n) = 0.$$

Theorem

Let some integration process \mathcal{A} be given which produces an integral additive on \mathcal{I} . Let a \mathcal{B} -interval function ψ be the quasi-measure generated by the series and (1) holds. Then this series is the Fourier series of an \mathcal{A} -integrable function f if and only if $\psi(I) = (\mathcal{A}) \int_I f$ for any \mathcal{B} -interval I .

Localization theorem

Localization Theorem.

Let the series

$$\sum_n c_n \gamma_n \quad (2)$$

is a formal product of the series

$$\sum_n a_n \gamma_n, \quad a_n \rightarrow 0,$$

and a polynomial $P = \sum_{n=0}^k b_n \gamma_n$. Then the series (2) and $P \sum_n a_n \gamma_n$ are uniformly equiconvergent, i.e., their difference is a series that is uniformly convergent on G with sum zero.

Localization theorem

Corollary.

For a series

$$\sum_n a_n \gamma_n, \quad a_n \rightarrow 0, \quad (3)$$

and a coset K_n there exists a series

$$\sum_n c_n \gamma_n, \quad c_n \rightarrow 0,$$

which is uniformly equiconvergent with (3) on K_n and is uniformly convergent to zero on $G \setminus K_n$.

Localization theorem

Corollary.

For a series

$$\sum_n a_n \gamma_n, \quad a_n \rightarrow 0, \quad (3)$$

and a coset K_n there exists a series

$$\sum_n c_n \gamma_n, \quad c_n \rightarrow 0,$$

which is uniformly equiconvergent with (3) on K_n and is uniformly convergent to zero on $G \setminus K_n$.

Lemma.

If for the series $\sum_n a_n \gamma_n$ with respect to the system Γ satisfying the condition $\lim_{n \rightarrow \infty} a_n = 0$, some subsequence of partial sums of the form $S_{m_{n_k}}(g)$ converges to zero for all g in some open set O , then this series converges to zero on O .

Construction of M_0 -set

Let h be a nondecreasing, right-continuous function, $h(0) = 0$. The Hausdorff h -measure \mathcal{H}^h is defined by

$$\mathcal{H}^h(A) = \liminf_{\delta \rightarrow 0} \sum \{h(\text{diam}(E_i)) : E_i \text{ open, } \cup E_i \supset A, \text{diam}(E_i) \leq \delta\}.$$

Construction of M_0 -set

Let h be a nondecreasing, right-continuous function, $h(0) = 0$. The Hausdorff h -measure \mathcal{H}^h is defined by

$$\mathcal{H}^h(A) = \liminf_{\delta \rightarrow 0} \sum \{h(\text{diam}(E_i)) : E_i \text{ open}, \cup E_i \supset A, \text{diam}(E_i) \leq \delta\}.$$

Theorem

For any h there exists a perfect M -set E for the system Γ whose Hausdorff h -measure equals zero. Moreover it is M_0 -set, i.e., a set of *strict multiplicity* (corresponding quasi-measure is a probability measure concentrated on E).

Construction of M -set

Lemma (main step in construction).

For any $A > 0$, $\varepsilon > 0$ there exists $N \in \mathbb{N}$ such that for any K_n , $n \geq N$, any $\delta > 0$, B , $0 < |B| \leq A$ and any natural λ there exists a polynomial

$$P(g) = \sum_{j=m_\lambda}^{m_l-1} a_j \gamma_j(g)$$

with properties:

1. $|a_j| < \varepsilon$, $m_\lambda \leq j \leq m_l - 1$;
2. $P(g) = 0$ if $g \in G \setminus K_n$;
3. $|P(g)| \geq |B|$ if $g \in K_n$;
4. $sh(1/m_l) < \delta$ where s is the number of those K_l in K_n for which $|P(g)| \neq |B|$ with $g \in K_l$.

Construction of M -set

Now using this Lemma we construct by induction a sequence of polynomial Q_k and a sequence of sets $E_k, E_{k-1} \subset E_k$, such that

$$\sum_{j=1}^k Q_j(t) = 0$$

outside E_{k-1} and $\mu_h E = 0$ where $E = \bigcap_{k=1}^{\infty} E_k$.

Construction of M -set

Now using this Lemma we construct by induction a sequence of polynomial Q_k and a sequence of sets $E_k, E_{k-1} \subset E_k$, such that

$$\sum_{j=1}^k Q_j(t) = 0$$

outside E_{k-1} and $\mu_h E = 0$ where $E = \bigcap_{k=1}^{\infty} E_k$.

It can be proved that required null-series is

$$\sum_{j=1}^{\infty} Q_j(t)$$

and E is required M_0 -set.