Proceedings of the DIS'2004, Štrbské Pleso, Slovakia

PENTAQUARK SEARCHES AT CDF

IGOR V. GORELOV
(FOR THE CDF COLLABORATION)
Department of Physics and Astronomy,
University of New Mexico,
800 Yale Blvd. NE, Albuquerque, NM 87131, USA
E-mail:gorelov@fnal.gov

Experimental results of a search for the $\Xi_{3/2}(1860)$ cascade pentaquark state in data collected with the CDF 2 Detector in Run II at the Tevatron are presented. No evidence for these states in the neutral $\Xi^-\pi^+$ and doubly charged $\Xi^-\pi^-$ modes has been found. Preliminary upper limits on yields at 1862 MeV/c^2 relative to the well established resonance $\Xi^*(1530)^0$ are presented.

1 Introduction

Evidence for an exotic pentaquark state $\Theta^+(1540)$ with strangeness $S = +1$ has been claimed by a number of experimental groups. Enhancements with a significance of 4.4 to 7.0 standard deviations have been observed in the invariant mass of K^+N in photoproduction [1] and in pK^0_s [2]. The signals seen have been assigned to a $\Theta^+(1540)$ state. These results have inspired a search for other exotic baryon states. The NA49 Collaboration has reported [3] the observation of a strangeness $S = -2$, isospin I =3/2 state $\Xi^{--}_{3/2} \to \Xi^-\pi^-$. An indication of a neutral mode in $M(\Xi^-\pi^+)$ has been demonstrated [3] as well. Recently the H1 Collaboration at HERA has published [4] an observation of a narrow anti-charmed baryon state in the mode $D^{*+}\bar{p}$ at ∼3099 MeV/c^2 and interpreted this as a heavy pentaquark Θ^0_c.

The pentaquark state Θ^+, according to the chiral soliton model [5], is considered as a bound state of five quarks. Experimental evidence for pentaquark Θ^+ suggested the existence of other pentaquark partners classified within the antidecuplet $\overline{10}$ representation [5] or the $\overline{10}_f \oplus 8_f$ multiplet as predicted by the constituent quark model approach in [6].

The experimental status of pentaquark baryons includes some controversy. The signal of Θ^+ claimed by [1,2] is not confirmed by [7]. The cascade pentaquark claimed by [3] has not been seen by [8]. Negative results on both Θ^+ and $\Xi^{--}_{3/2}$ have been reported by large statistics experiments [9]. These experiments exploit their excellent mass resolution and large data samples to calibrate the mass spectra of interest by well established states like $\Lambda(1520) \to pK^-$ and $\Lambda^+_c \to pK^0_s$ (for Θ^+ searches) and $\Xi^*(1530)^0 \to \Xi^-\pi^+$ (for $\Xi_{3/2}$ searches).

Recently the CDF Collaboration undertook a comprehensive search for several pentaquark states using its Run II 220 pb^{-1} of data taken with the upgraded CDF 2 Detector. We present here the particular search for the cascade pentaquark $\Xi_{3/2}$ through its modes $\Xi^0_{3/2} \to \Xi^-\pi^+$ and $\Xi^{--}_{3/2} \to \Xi^-\pi^-$, both of which involve the

© Institute of Experimental Physics SAS, Košice, Slovakia

doubly strange cascade baryon Ξ^- in the final state. The analysis is based on two data samples. The first one was collected by a trigger selecting events with at least two tracks of opposite charge each, having a momentum above 2.0 GeV/c and an impact parameter measured by the CDF silicon detector to be larger than 100 μm. The total momentum of both tracks was required to be larger than 5.5 GeV/c. This "two displaced track trigger" sample is enriched by events with heavy quarks decaying via hadronic modes. A complementary dataset was taken with a trigger requiring an inclusive jet of transverse energy E_T >20 GeV.

2 Cascades in the CDF 2 Detector

The final state cascade baryon Ξ^- [a] decays almost 100% of time into $\Lambda^0\pi^-$, with a subsequent decay of the Λ^0. Since the days of bubble chamber experiments this spectacular mode has been identified and reconstructed as a single track vertex (we call it here `VERTEX_1`) formed by a kinked track, presumably the π^-, followed in the fiducial volume by another, V^0-vertex (`VERTEX_2`), presumably the Λ^0 decaying to $p\pi^-$, as sketched at Fig.1. In our analysis `VERTEX_2` was subjected to a 3-dimensional fit. Then the three tracks $p\pi^-$ and $\pi^{+,-}$ were fitted to a common 3-dimensional vertex with the constraint that $M(p\pi^-) = M^{\Lambda^0}_{PDG}$ and that `VERTEX_2` points back to `VERTEX_1`, see also Fig.1.



Figure 1: Sketch of the cascade decay topology. $M(p\pi^-) \in M^{\Lambda}_{PDG}\pm 5$ MeV/c^2 and the 2-dimensional $\chi^2_{r\phi} < 49.0$ cut were applied. $M(p\pi^-\pi_{kink}) \in M^{\Xi}_{PDG}\pm 60$MeV/c^2 and no $p\pi^-\pi_{kink}$ vertex fit quality χ^2 cut were required. The Ξ and Λ vertexes are separated by more than 1 cm in a transverse plane and an impact parameter $d_0(\Xi) <$ 150 μm defined in a transverse plane as well.

The resulting invariant mass spectrum of cascade candidates $M(\Xi^- \to \Lambda^0\pi^-)$ is shown on Fig.2. The clear signal of a cascade is seen on top of a large combinatorial background.



Figure 2: The invariant mass spectrum of $M(\Lambda^0\pi^-)$ after vertex fits described above. The cascade signal is present with a large combinatorial background. The shaded histogram corresponds to the wrong charge combinations $M(\Lambda^0\pi^+)$.

The long lifetime of Ξ^- hyperons ($c\tau$=4.91 cm) permits reconstruction of their tracks from hits in the CDF silicon tracker (SVX II). A novel technique developed

[a] Unless otherwise stated all references to the specific charge combination imply the charge conjugate combination as well.

by CDF uses the vertex position and momentum of a cascade hyperon fitted in the CDF outer tracker to seed the hyperon tracking in SVX II. This procedure results in a substantial background reduction and improved vertex and impact parameter resolution of the Ξ^-, see Fig. 3. The overall relative efficiency of the hyperon reconstruction with SVX II hits is ∼40%. The yields of cascades in the CDF 2 Detector are larger by a factor of ∼20 than the ones found by the NA49 experiment [3].



Figure 3: The invariant mass spectrum of Ξ^- hyperons which have tracks successfully reconstructed in the silicon tracker. A cut on impact parameter $d_0(\Xi)$ <150 μm was applied, selecting hyperons produced promptly in the primary vertex region. The very clean signal based on the data sample of integrated $\mathcal{L}$=220 pb^{-1} taken by the two displaced track trigger contains ∼36000 events. The analogous signal (not shown here) for the inclusive jet E_T >20.0 GeV dataset of the same $\mathcal{L}$ contains ∼4870 events.

3 Pentaquarks in the $\Xi^-\pi^+$ and $\Xi^-\pi^-$ Decay Modes

The hyperon tracks reconstructed in SVX II with mass $M(\Xi) \in M^{\Xi}_{PDG} \pm 10$ MeV/c^2 (see Fig. 3) were combined with all remaining tracks with P_T >400 MeV/c and 3 or more hits in the SVX II tracker. Then the track pairs $\Xi^-\pi^{+,-}$ were subjected to a vertex fit constrained by the requirement for the secondary vertex to point to the primary one. The invariant mass spectra $M(\Xi^-\pi^{+,-})$ are shown in Fig. 4 and 5.



Figure 4: The invariant mass spectrum for $\Xi^-\pi^+$ whose hyperons have tracks successfully reconstructed in the SVX II. The track pairs form the vertex fitted with two-dimensional $\chi^2_{r\phi} < 36.0$. The two displaced track trigger data sample of integrated luminosity $\mathcal{L}$=220 pb^{-1} is used. The fit to a Breit-Wigner convoluted with a Gaussian finds 2182±92 events in the peak for $\Xi^*(1530)^0 \to \Xi^-\pi^+$. The peak is used as a gauge signal for pentaquark searches. The similar spectrum (not shown here) in the inclusive jet E_T >20 GeV sample yields 387±34 events for $\Xi^*(1530)^0$. The pentaquark signal region at 1862 MeV/c^2 is fitted with a Gaussian of a fixed width σ =8 MeV/c^2 predicted by Monte-Carlo simulation.



Figure 5: The invariant mass spectrum for $\Xi^-\pi^-$ whose hyperons have tracks successfully reconstructed in the SVX II. The two displaced track trigger data sample of $\mathcal{L}$=220 pb^{-1} is used. The spectrum is fitted by a polynomial background shaped by a square-root threshold function. The NA49 signal region at $M(\Xi^-\pi^-)$ = 1862 MeV/c^2 is fitted with a Gaussian of fixed width σ =8 MeV/c^2. A similar spectrum (not shown here) is observed for the inclusive jet E_T >20 GeV sample.

The spectra $M(\Xi^-\pi^+)$ (Fig. 4) and $M(\Xi^-\pi^-)$ (Fig. 5) corresponding to neutral and doubly charged cascade pentaquark modes do not reveal any enhancement around M=1862 MeV/c^2 [3]. Similar results have been obtained with the inclusive jet E_T >20 GeV sample. We have set upper limits on the production of pentaquarks decaying via both modes. These are shown in a Table 1 below.

Mode	@90%$C.L.$	@95%$C.L.$
Two Displaced Track Trigger Sample		
$\sigma\cdot$Br$(\Xi^-\pi^+)/\sigma\cdot$Br$(\Xi^*(1530)^0)$	0.06	0.07
$\sigma\cdot$Br$(\Xi^-\pi^-)/\sigma\cdot$Br$(\Xi^*(1530)^0)$	0.03	0.04
combined statistics	0.07	0.08
Inclusive Jet E_T >20 GeV Sample		
$\sigma\cdot$Br$(\Xi^-\pi^+)/\sigma\cdot$Br$(\Xi^*(1530)^0)$	0.06	0.08
$\sigma\cdot$Br$(\Xi^-\pi^-)/\sigma\cdot$Br$(\Xi^*(1530)^0)$	0.07	0.09
combined statistics	0.09	0.11

Table 1: Upper limits set for a $\Xi^0_{3/2}$ and $\Xi^{--}_{3/2}$ pentaquark states. The yields were calculated relative to the calibrating signal of $\Xi^*(1530)^0$ seen in both data samples.

4 Summary

The CDF Collaboration conducted a search for doubly strange $S = -2$ pentaquark states in the $\Xi^-\pi^+$ and $\Xi^-\pi^-$ decay modes. The signals of the basic hyperon Ξ^- state comprised ~36000 events in the two displaced trigger dataset and ~4900 events in the inclusive jet E_T >20 GeV dataset. The well established resonance $\Xi^*(1530)^0 \to \Xi^-\pi^+$ was used as a calibrating signal and yielded 2182±92 events from the two displaced trigger sample and 387±34 events from the inclusive jet E_T >20 GeV sample. No evidence of exotic baryon states produced in CDF Detector at Tevatron has been found. Upper limits on production of states in the mass range of ~1862 MeV/c^2 have been set. CDF Collaboration is pursuing a vigorous program of searches for possible pentaquark production at the Tevatron.

Acknowledgments

The author is grateful to his colleagues from the CDF Pentaquark Working Group, especially to Dr. D. Litvintsev for useful suggestions and comments made during

preparation of this talk. The author would like to thank Prof. Sally C. Seidel for support of this work, fruitful discussions, and comments.

References

1. T. Nakano *et al.*, Phys. Rev. Lett. **91**, 012002(2003).
   T. Stepanyan *et al.*, CLAS Collab., Phys. Rev. Lett. **91**, 252001(2003), [hep-ex/0307018].
   T. Barth *et al.*, SAPHIR Collab., Phys. Lett. B **572**, 127(2003), [hep-ex/0307083].
   V. Kubarovsky *et al.*, SAPHIR Collab., Phys. Rev. Lett. **92**, 032001(2004); Erratum ibid. **92**, 049902(2004).
2. V. Barmin *et al.*, DIANA Collab., Phys. Atom. Nucl. **66**, 1715(2003); Yad. Fiz. **66**, 1763(2003).
   T. Airapetian *et al.*, HERMES Collab., hep-ex/0312044.
   A.E. Asratyan, A.G. Dolgolenko, and M.A. Kubantsev, hep-ex/0309042.
   T. Aleev *et al.*, SVD Collab., hep-ex/0401024.
   S. Chekanov, *et al.*, ZEUS Collab., Phys. Lett. B **591**, 7(2004), [hep-ex/0403051].
   M. Abdel-Bary *et al.*, COSY-TOF Collab., hep-ex/0403011.
3. C. Alt, *et al.*, NA49 Collab., Phys. Rev. Lett. **92**, 042003(2004), [hep-ex/0310014].
4. A. Aktas, H1 Collab., hep-ex/0403017.
5. D. Diakonov, V. Petrov, M. Polyakov, Z. Phys. A **359**, 309(1997), [hep-ph/9703373].
6. M. Karliner and H.J. Lipkin, [hep-ph/0307243].
   R.I. Jaffe and F. Wilczek, Phys. Rev. Lett. **91**, 232003(2003), [hep-ph/0307341].
7. J.Z. Bai *et al.*, BES Collab., hep-ph/0312269.
   K.T. Knöpfle *et al.*, HERA-B Collab., hep-ex/0403020.
   C. Pinkenburg *et al.*, PHENIX Collab., nucl-ex/0404001.
8. K.T. Knöpfle *et al.*, HERA-B Collab., hep-ex/0403020.
   S. Chekanov, for the ZEUS Collab., Proceedings of this conference, [hep-ex/0405013].
9. A. Dzierba, `http://www.qnp2004.org/5q_talks/dzierba_5q_qnp2004.pdf`, talk at "Quarks and Nuclear Physics 2004," May 23-28, 2004, Bloomington, IN, USA.